


# Atlantic States Marine Fisheries Commission

## *2022 Atlantic Striped Bass Stock Assessment Update Report*



Accepted for Management Use  
by the Atlantic Striped Bass Management Board  
November 2022



*Sustainable and Cooperative Management of Atlantic Coastal Fisheries*

**Atlantic States Marine Fisheries Commission**

*Atlantic Striped Bass Stock Assessment Update*

Prepared by the

ASMFC Striped Bass Stock Assessment Subcommittee  
Mike Celestino, Chair, New Jersey Department of Environmental Protection  
Margaret Conroy, Delaware Division of Fish and Wildlife  
Hank Liao, Virginia Marine Resources Commission  
Gary Nelson, Massachusetts Division of Marine Fisheries  
Alexei Sharov, Maryland Department of Natural Resources  
John Sweka, US Fish and Wildlife Service  
Katie Drew, Atlantic States Marine Fisheries Commission

and the

ASMFC Striped Bass Technical Committee  
Nicole Lengyel Costa, Chair, Rhode Island Department of Environmental Management  
Ingrid Braun, Potomac River Fisheries Commission  
Michael Brown, Maine Department of Marine Resources  
Margaret Conroy, Delaware Division of Fish and Wildlife  
Caitlin Craig, New York Department of Environmental Conservation  
John Ellis, US Fish and Wildlife Service  
Charlton Holloman Godwin, North Carolina Division of Marine Fisheries  
Kurt Gottschall, Connecticut Department of Energy and Environmental Protection  
Tyler Grabowski, Pennsylvania Fish and Boat Commission  
Brendan Harrison, New Jersey Department of Environmental Protection  
Luke Lyon, District of Columbia Department of the Environment  
Jeremy McCargo, North Carolina Wildlife Resources Commission  
Joshua McGilly, Virginia Marine Resources Commission  
Steve Minkinen, US Fish and Wildlife Service  
Gary Nelson, Massachusetts Division of Marine Fisheries  
Alexei Sharov, Maryland Department of Natural Resources  
Kevin Sullivan, New Hampshire Fish and Game Department  
Tony Wood, NOAA Fisheries

A publication of the Atlantic States Marine Fisheries Commission pursuant to National Oceanic and Atmospheric Administration Award No. NA20NMF4740012.



## EXECUTIVE SUMMARY

The time series of striped bass removals and indices from the 2018 benchmark assessment was updated to include data from 2018-2021. Total removals from 2018-2021 averaged 5.37 million fish annually, a 24% decrease from 2017, the terminal year of the last assessment when the stock was experiencing overfishing. From 2018-2021, recreational release mortality made up 50% of total removals, with recreational harvest making up 37%, commercial harvest making up 11%, and commercial discards making up 2% of the total.

COVID-19 affected fishery-dependent and fishery-independent sampling for state surveys and the MRIP dockside intercept program, although the level of impact varied from state to state. The assessment model was able to accommodate the missing index data, but overall, COVID-19 increased uncertainty in the 2020 and 2021 data.

The single-stock statistical catch-at-age (SCA) model was updated through 2021. The model parameterization was the same as in the benchmark assessment, with the exception of a new selectivity block from 2020-2021 in the Bay and Ocean fleets, to account for the regulation changes from Addendum VI. Sensitivity runs were conducted to look at the effect of only including a new selectivity block in the Ocean fleet and the effect of not including any new selectivity blocks.

Because the recruitment trigger in Amendment 7 was tripped in 2021 for the Maryland juvenile abundance index, the biological reference points were updated using the low recruitment regime assumption. This resulted in a lower  $F$  target and  $F$  threshold compared to the benchmark assessment.

In 2021, the Atlantic striped bass stock was overfished but was not experiencing overfishing. Female spawning stock biomass 2021 was estimated at 64,805 metric tons (143 million pounds) which is below the updated SSB threshold of 85,457 metric tons (188 million pounds), and below the updated SSB target of 106,820 metric tons (235 million pounds). Total fishing mortality in 2021 was estimated at 0.14 which is below the updated  $F$  threshold of 0.20 per year, and below the updated  $F$  target of 0.17 per year.

The sensitivity run with the new selectivity block for the Ocean fleet only produced very similar results to the base run, while the sensitivity run with no new selectivity blocks produced higher estimates of  $F$  and lower estimates of SSB in 2020-2021. However, stock status was the same for all three runs.

The retrospective pattern remained moderate to low in magnitude for the assessment update, but reversed direction compared to the benchmark; the model underestimated  $F$  and overestimated SSB in the most recent peels. The retrospective-adjusted estimates of  $F$  and SSB were within the 90% confidence intervals of the unadjusted estimates, so correcting for retrospective pattern was not necessary for status determination or projections.

Projections were run to determine the probability of SSB being at or above the SSB target by 2029, the rebuilding deadline. Under the current  $F$ , there is a 78.6% chance the stock will be rebuilt by 2029, indicating a reduction in catch is not necessary at this time.

The sensitivity run with a new selectivity block in the Ocean fleet only produced very similar results to the base model, but the run with no new selectivity blocks was more pessimistic about rebuilding, requiring an 8.6% reduction in removals to have a 50% chance of being at or above the SSB target in 2029. However, there was a greater than 50% chance of being above the SSB threshold by 2029 for all three runs.

	<b>Target</b>	<b>Threshold</b>	<b>2021 Value</b>	<b>Status</b>
<b>Fishing Mortality</b>	0.17	0.20	0.14	Not overfishing
<b>Female SSB</b>	106,820 mt (235 million lbs)	85,457 mt (188 million lbs)	64,805 mt (143 million lbs)	Overfished

## TABLE OF CONTENTS

EXECUTIVE SUMMARY .....	ii
TERMS OF REFERENCE (TOR) .....	1
TOR 1. Update fishery-dependent data (landings, discards, catch-at-age, etc.) that were used in the previous peer-reviewed and accepted benchmark stock assessment.....	1
TOR 2. Update fishery-independent data (abundance indices, age-length data, etc.) that were used in the previous peer-reviewed and accepted benchmark stock assessment.....	1
TOR 3. Tabulate or list the life history information used in the assessment and/or model parameterization (M, age plus group, start year, maturity, sex ratio, etc.) and note any differences (e.g., new selectivity block, revised M value) from benchmark. ....	2
TOR 4. Update accepted model(s) or trend analyses and estimate uncertainty. Include sensitivity runs and retrospective analysis if possible and compare with the benchmark assessment results. Include bridge runs to sequentially document each change from the previously accepted model to the updated model.....	3
TOR 5. Update the biological reference points or trend-based indicators/metrics for the stock. Determine stock status. ....	5
TOR 6. Conduct short term projections when appropriate. Discuss assumptions if different from the benchmark and describe alternate runs.....	6
TOR 7. Comment on research recommendations from the benchmark stock assessment and note which have been addressed or initiated. Indicate which improvements should be made before the stock undergoes a benchmark assessment. ....	8
Literature Cited.....	8
List of Appendices.....	8
TABLES .....	9
FIGURES .....	16

## TERMS OF REFERENCE (TOR)

### **TOR 1. Update fishery-dependent data (landings, discards, catch-at-age, etc.) that were used in the previous peer-reviewed and accepted benchmark stock assessment.**

The time series of striped bass recreational and commercial removals from the 2018 benchmark assessment (NEFSC 2019) was updated to include data from 2018-2021. This included recreational harvest, recreational release mortalities, commercial harvest, and commercial discards.

Total removals from 2018-2021 averaged 5.37 million fish annually, a 24% decrease from 2017, the terminal year of the last assessment when the stock was experiencing overfishing (Table 1, Figure 2). Approximately 62% of the removals came from the ocean fleet over that time period, while 38% came from the Chesapeake Bay fleet, consistent with the overall percentages for the whole time series (Table 1, Figure 1).

From 2018-2021, recreational release mortality made up 50% of total removals, with recreational harvest making up 37% and commercial harvest making of 11% of the total (Figure 2). Commercial dead discards made up approximately 2% of the total removals.

COVID-19 had an impact on fishery-dependent data collection during 2020. Biological sampling levels for the recreational and commercial fisheries were reduced, which increased uncertainty somewhat in the catch-at-age for both fisheries. The MRIP effort survey continued uninterrupted, but the Access Point Angler Intercept Survey (APAIS) was suspended for part of 2020. Data from 2018 and 2019 were used to impute total recreational catch rates for 2020 where necessary. Overall, 29% of recreational harvest rate information and 15% of released alive rate information was attributed to imputed catch data for 2020 (Table 2). The percentage of imputed information in 2020 recreational catch rates varied from state to state, depending on the length of time that APAIS was suspended. Although COVID likely affected the overall harvest from the commercial fishery, it did not significantly impact reporting the catch.

The MRIP CPUE index of abundance was updated with data through 2021. The index was developed using the same species associations identified in the previous benchmark. Imputed records were excluded from the intercept data pull for 2020. The index declined somewhat from 2018-2021.

### **TOR 2. Update fishery-independent data (abundance indices, age-length data, etc.) that were used in the previous peer-reviewed and accepted benchmark stock assessment.**

Where possible, the fishery independent age-1+ and recruitment indices used in the most recent benchmark assessment (Table 3) were updated through 2021. Several surveys were impacted by COVID and other issues in the most recent years (Table 4 and Table 5).

The assessment used seven fishery independent indices of age-1+ abundance: the Chesapeake Bay Multispecies Monitoring and Assessment Program (ChesMMAP), the Maryland Spawning

Stock Survey (MDSSN), the Delaware Spawning Stock Electrofishing Survey (DESSN), the Delaware 30' Bottom Trawl Survey (DE30), the New York Ocean Haul Seine (NYOHS), the New Jersey Bottom Trawl Survey (NJTRL), and the Connecticut Long Island Sound Trawl Survey (CTLISTS). The NJ Trawl did not operate from 2019-2021 due to COVID and vessel issues. ChesMMAP changed vessels in 2018 and the calibration process has not been finished, so calibrated estimates were not available for 2019-2021 in time for this update. The DE SSN and CT LIST surveys did not operate in 2020 due to COVID. The MD SSN was interrupted for two weeks in 2021 due to COVID. Age-1+ surveys with data through 2021 showed mixed trends, with some surveys increasing since 2017 and some decreasing (Figure 3).

The assessment uses four age-0 juvenile abundance indices (JAI) and two age-1 indices as recruitment indices: the MD, VA, NJ, and NY JAIs and the MD and NY age-1 indices. The MD and VA JAIs were combined into a single composite JAI for Chesapeake Bay using the Conn (2010) method. The NJ JAI was the only survey that did not occur in 2020 due to COVID, although the start of the NY Age-1 survey was delayed. 2018 values indicated a strong year class in most indices, but 2021 was generally low (Figure 4 and Figure 5). The MD JAI tripped the recruitment trigger in 2021, with three consecutive years below the Amendment 7 recruitment threshold.

**TOR 3. Tabulate or list the life history information used in the assessment and/or model parameterization (M, age plus group, start year, maturity, sex ratio, etc.) and note any differences (e.g., new selectivity block, revised M value) from benchmark.**

Model equations are shown in Appendix 1 Table 1. The model parameterization was the same as used in the benchmark assessment (NEFSC 2019), with the exception of a new selectivity block from 2020-2021 in the Bay and Ocean fleets, to account for the regulation changes from Addendum VI (Table 6). In initial runs, the exponential-logistic and double-logistic selectivity equations were used to explore if the selectivity during 2020-2021 changed to dome-shaped due to changes in size-limits, particularly in the Ocean. Initial results showed that the 2020-2021 selectivity pattern in the Bay remained dome-shaped, and the 2020-2021 selectivity pattern in the Ocean remained flat-topped. Therefore, the exponential-logistic and Gompertz functions were used to model selectivity for 2020-2021.

Re-weighting of survey indices was required with the addition of four years of removal data and missing index data for several surveys. Survey CVs were adjusted to bring the RMSE close to one and effective sample sizes were adjusted once by using the Francis multipliers (Francis 2011). The RMSEs, CV weights and effective samples from the 2018 benchmark and 2022 assessment models are given in Table 2 in Appendix 1. The largest change in CV weight occurred for the NJ Trawl survey, where the correct CV time series was substituted for the incorrect values input in the benchmark.

No changes were made to the life history information used in the assessment (Table 7).

**TOR 4. Update accepted model(s) or trend analyses and estimate uncertainty. Include sensitivity runs and retrospective analysis if possible and compare with the benchmark assessment results. Include bridge runs to sequentially document each change from the previously accepted model to the updated model.**

### **Model Fit**

The model fit the observed total catches and catch age compositions of all fleets well (Appendix 2). The model fit the MDYOY (1970-1981) and MD & VA composite indices very well and the MD Age1, NYOHS, and MDSSN poorly. It fit the other indices reasonably well (Appendix 2). The predicted trends matched the observed trends in age composition of survey indices reasonably well for NYOHS, MDSSN, MRIP, CTLIST, and ChesMMAP. The model fit the age composition of NJTrawl, DESSN, and DE30FT survey adequately. Resulting contributions to total likelihood are listed in Table 3 of Appendix 1. Estimates of fully-recruited fishing mortality for each fleet and total fishing mortality, recruitment, parameters of the selectivity functions for the selectivity periods, catchability coefficients for all surveys, and parameters of the survey selectivity functions are given in Table 4 of Appendix 1.

Estimates of the catch selectivity patterns for each fleet showed that, although the patterns varied over time with changes in regulation, selectivity was dome-shaped for Chesapeake Bay and primarily flat-topped for the Ocean over time (Figure 6). There was a steep shift in the descending limb of the selectivity pattern in 2020-2021 for Chesapeake Bay compared to the previous selectivity block, and a shift in the selectivity in 2020-2021 for the Ocean to lower ages (Figure 6).

### **Fishing Mortality**

Fully-recruited annual fishing mortality in 2021 for the Bay and Ocean was 0.05 and 0.10 (Figure 7), and peaked at ages 6 and 10-15, respectively. Total fully-recruited  $F$  in 2021 was 0.14 (Table 8, Figure 7) and peaked at age 6. Coefficients of variation indicated region-specific and total fishing mortality estimates were precise (CVs mostly less than 0.20) (Table 4 of Appendix 1).

### **Recruitment**

Recruit numbers increased steadily through 1993 (Figure 8). Large recruitment events occurred in 1994, 1997, 2002, and 2004 as the large Chesapeake Bay 1993, 1996, 2001 and 2003 year-classes became age-1. Average to below-average year-classes were produced during 2004-2010, which resulted in a decline of age-1 numbers. Subsequently, strong year-classes were produced in 2011 and 2015. After 2016, recruit abundance fluctuated slightly and has averaged 123.5 million fish (Table 8, Figure 8). Four of the last five year-classes since 2015 have been below average, although not as low as the levels seen from 2004-2010; the 2018 year-class was above average (Table 8, Figure 8). The below-average 2020 and 2021 recruits will start contributing to SSB in 2027 and 2028 as those fish approach full maturity.

### **Population Abundance (January 1)**

Striped bass abundance (1+) increased steadily from 1982 through 1997 when it peaked



around 422.4 million fish (Table 8, Figure 9). Total abundance fluctuated without trend through 2004. From 2005-2009, age 1+ abundance declined to about 181.2 million fish. Thereafter, total abundance peaked in 2012 and 2016 as a result of two large year-classes (2011 and 2015) entering the age-1+ population (Table 8, Figure 9). From 2017-2019, total abundance averaged 243.3 million fish. Abundance declined slightly through 2021 to 218.9 million fish (Figure 9).

Abundance of striped bass age 8+ increased steadily through 2004 to 16.6 million fish, but then declined to 11.4 million fish through 2010 (Table 8, Figure 9). A small increase in 8+ abundance occurred in 2011 as the 2003 year-class became age 8 (Table 8, Figure 9). Abundance of age 8+ fish declined steadily through 2018 but has increased recently to an average of 6.7 million fish as the 2011 aged recruited to the age-8+ group (Table 8, Figure 9).

### **Spawning Stock Biomass and Total Biomass**

Female SSB grew steadily from 1982 through 2003 when it peaked at about 113,000 metric tons (Table 8, Figure 10). Female SSB declined steadily from 104,749 metric tons in 2010 to 55,120 metric tons in 2018, but in recent years, has steadily increased (Table 8, Figure 10). Estimates of female spawning stock biomass were very precise (CVs less than 0.14; Table 10 of Appendix 1).

Exploitable biomass (January 1) increased from 36,985 metric tons in 1982 to its peak at 333,000 metric tons in 1999 but declined steadily through 2015 (Figure 10). Since 2016, exploitable biomass steadily increased albeit at a slow pace.

### **Retrospective Analysis**

Moderate retrospective patterning (<15%) was evident in the more recent estimates of fully-recruited total  $F$  and female SSB (Figure 11). The retrospective pattern suggested that fishing mortality is likely slightly under-estimated (<12%) and female spawning biomass is over-estimated by 5-17%. Recruitment appeared to be over-estimated in most years, although underestimation did occur in a few years (Figure 11). The Mohn's rho values for fishing mortality, female SSB and recruitment were estimated to be -0.087, 0.103 and 0.156, respectively.

The current retrospective trends are different from what was observed in the 2018 benchmark and earlier assessments (NEFSC 2019). The past retrospective patterns showed that female SSB was typically under-estimated and fishing mortality was over-estimated. Exploratory analyses indicated that the change was due, in part, to the addition of new data and changes in index weighting. When the index CV weightings from the 2018 benchmark assessment was used in the current assessment, the past retrospective pattern was reproduced through the 2016 peel and then changed to what is observed currently, albeit at a lower level of percent difference (Appendix 1).

### **Sensitivity Runs**

The NY Age-1 seine survey and MD SSN survey were completed in all years, but the timing of each was affected by the COVID pandemic: the NY Age-1 survey started later than usual in 2020 and the MD SSN survey was suspended for two weeks in 2021. To determine if these potentially

biased values influenced the results of the assessment, a run was made in which those index values were coded as missing. Comparison of results (Figure 12) showed that the missing values had little influence on the time series of  $F$  and SSB estimates.

Two additional runs were made to explore the influence of using the new selectivity blocks in 2020-2021. One run was made in which a new 2020-2021 selectivity block was created only for the Ocean region and a second was made in which no new selectivity periods were created. Full results and diagnostics for these sensitivity runs are presented in Appendix 3 and 4.

Comparison of residual plots, particularly for the fleet age composition, showed that the base run produced the smallest residuals in 2020-2021 (Appendices 2-4). Based on Mohn’s rho, the base model had the lowest retrospective pattern ( $F=-0.087$ ;  $SSB=0.103$ ) compared to the Ocean only run ( $F=-0.094$ ;  $SSB=0.121$ ) and the no new selectivity blocks run ( $F=-0.107$ ;  $SSB=0.177$ ).

The run with the new selectivity for the Ocean fleet only produced very similar results to the base run, but the run with no new selectivity blocks produced higher estimates of  $F$  and lower estimates of SSB in 2020-2021 (Figure 13).

**Comparison of Results from the 2018 Benchmark Assessment with 2022 Update Assessment**

Fully-recruited fishing mortality and female spawning stock biomass estimates from the update and benchmarks assessments are shown in Figure 14. The updated assessment produced higher estimates of fishing mortality in 2012-2017 and lower estimates of female spawning stock biomass from 1992-2001 and 2012-2017.

**TOR 5. Update the biological reference points or trend-based indicators/metrics for the stock. Determine stock status.**

The fishing mortality and spawning stock biomass reference points were updated using the same methods as the benchmark assessment (NEFSC 2019). The spawning stock biomass threshold is the 1995 estimate of SSB from the current assessment and the SSB target is 125% of the threshold. Using a stochastic projection drawing recruitment from empirical estimates and a distribution of starting population abundance at age, fishing mortalities associated with the SSB target and threshold were determined. Empirical estimates of recruitment, selectivity, and the starting population came from the SCA model results. The selectivity pattern used in the projections was calculated as the geometric mean of the 2020-2021 total  $F$ -at-age, scaled to the highest  $F$ -at-age (Figure 15). Estimates of recruitment were restricted to 2008-2021 to represent the “low” recruitment regime. The population was projected for 100 years and fully-recruited  $F$  was adjusted until the median of the projected SSB reached the SSB target or threshold.

The updated SSB reference points and associated fishing mortalities are:

<b>SSB<sub>threshold</sub> = 85,457 metric tons</b>	<b>F<sub>threshold</sub> = 0.20</b>
<b>SSB<sub>target</sub> = 106,820 metric tons</b>	<b>F<sub>target</sub> = 0.17</b>

## Status of the Stock

Before stock status can proceed, analyses must be done to determine if the estimates of  $F$  and SSB in 2021 should be corrected for the apparent pattern observed in the retrospective analyses. Here we used the National Marine Fisheries Service standard procedure in which the estimates are adjusted for the retrospective pattern using Mohn's rho values (average of proportion differences over seven-year peels) and then compared to the unadjusted estimates and their associated 90% confidence intervals. If either retrospective-adjusted value falls outside an unadjusted value's 90% confidence intervals, then the retrospective-adjusted values are used. If not, the unadjusted values are sufficient for stock determination. Figure 16 shows a bivariate plot of the unadjusted estimates and their associated 90% confidence interval along with the retrospective-adjusted values. Because the retrospective-adjusted values fall within the 90% confidence intervals, retrospective adjustment is not needed.

In 2021, the Atlantic striped bass stock was overfished but was not experiencing overfishing based on the point estimates of fully-recruited fishing mortality and female spawning stock biomass relative to the reference points defined in this assessment. Female spawning stock biomass in 2021 was estimated at 64,805 metric tons (143 million pounds) which is below the SSB threshold of 85,457 metric tons (188 million pounds), and below the SSB target of 106,820 metric tons (235 million pounds) (Table 9, Figure 17). However, because of error associated with these estimates, there is a 0.9% probability that the 2021 female SSB estimate is above or equal to the SSB threshold and a 0% probability that the 2021 estimate is above the target.

Total fishing mortality in 2021 was estimated at 0.14 which is below the  $F$  threshold of 0.20 and the  $F$  target of 0.17 (Table 9, Figure 17). There is a 99.6% probability that the 2021 fully-recruited fishing mortality is below the fishing mortality threshold, and a 91% probability that the value is below the  $F$  target.

Although the estimate of  $F$  in 2021 was higher for the sensitivity run with no new selectivity blocks, stock status was the same for all three sensitivity runs: overfishing was not occurring and the stock was overfished.

### **TOR 6. Conduct short term projections when appropriate. Discuss assumptions if different from the benchmark and describe alternate runs.**

Three scenarios were run to determine when female SSB is expected to reach the SSB target under the "low" recruitment regime. In the first run, the population was projected over ten years assuming the  $F$  observed in 2021 (0.14) was the same in 2022-2030. In the second and third runs, the population was projected assuming fishing mortality in 2022-2030 was equal to  $F$  associated with the  $F$  target and  $F$  threshold values. Because the retrospective adjusted values of  $F$  and SSB fell within the 90% confidence intervals of the unadjusted estimates, retrospective-adjustment was not needed.

The projections used the same methods as the benchmark assessment (NEFSC 2019). For each scenario, the model begins in year 2021 with the estimates of January-1 abundance-at-age and associated standard errors from the SCA assessment model. The fully-recruited  $F$  estimate and

associated standard errors in 2021 ( $F=0.14$ ), selectivity-at-age in 2021, Rivard weights in 2021, natural mortality, female sex proportions-at-age, and female maturity-at-age are used to calculate female SSB as modeled in the SCA model. For 2022, the January-1 abundance-at-age is calculated from the known values of 2021 abundance-at-age, 2021 selectivity and fully-recruited  $F$  for 2021. For the remaining years, the January-1 abundance-at-age is projected and is calculated by using the previous year's abundance-at-age, the scenario fully-recruited  $F$ , and natural mortality following the standard exponential decay model. Female spawning stock biomass is calculated using the average Rivard weights-at-age from 2017-2021 along with proportion of female by age and maturity-at-age.

For each iteration of the simulation, the abundance and fishing mortality-at-age values in 2021 are randomly drawn from a normal distribution parameterized with the associated standard errors from the SCA assessment model. For the remaining years, abundance of age-1 recruits is randomly drawn from 2008-2021 recruitment estimates. An age-15 plus-group is assumed. For years 2022-2030, selectivity-at-age is assumed equal to the geometric mean selectivity for years 2020-2021. Female spawning stock biomass was calculated by using geometric mean Rivard weight estimates from 2017-2021, sex proportions-at-age, and female maturity-at-age. For each year of the projection, the probability of SSB being above the SSB target and threshold reference points was calculated from 10,000 simulations using function *pgen* in R package *fishmethods*.

## Results

Under current fully-recruited fishing mortality ( $F=0.14$ ), female SSB is expected to reach or exceed the SSB threshold by 2023 with a probability of 70.2%, and exceed or reach the SSB target by 2025 with a probability of 56.1% (Table 10, Figure 18). By the rebuilding deadline of 2029, there is a 78.6% chance the stock will be at or above the SSB target and a 96.7% chance the stock will be at or above the SSB threshold. Under  $F$  target ( $F=0.17$ ), female SSB is expected to reach or exceed the SSB threshold by 2023 with a probability of 61.9%, and exceed or reach the SSB target by 2028 with a probability of 52.0% (Table 10, Figure 18). Under  $F$  threshold ( $F=0.20$ ), female SSB is expected to reach or exceed the SSB threshold by 2023 with a probability of 53.2%, but has a less than 50% probability of reaching the SSB target in any year (Table 10, Figure 18).

The sensitivity run with a new selectivity block in the Ocean fleet only produced very similar results to the base model, but the run with no new selectivity blocks was more pessimistic about rebuilding, with the stock having a less than 20% chance of rebuilding under current  $F$  by 2029 (Appendix 4). An 8.6% reduction in removals would be required to have a 50% chance of being at or above the SSB target in 2029 under that model configuration. However, the stock did have a greater than 50% chance of being above the SSB threshold by 2029 in all three runs.

**TOR 7. Comment on research recommendations from the benchmark stock assessment and note which have been addressed or initiated. Indicate which improvements should be made before the stock undergoes a benchmark assessment.**

The research recommendations identified in the benchmark assessment (NEFSC 2019) remain relevant, particularly the research recommendations on enhanced collection of life history and biological information including paired scale-otolith samples, migration rates, and sex ratio data. Additional work on refining migration rates and stock composition estimates as well as incorporating tagging data into the spatial statistical catch-at-age model will be required before the next benchmark assessment.

**Literature Cited**

Conn, P.B. 2010. Hierarchical analysis of multiple noisy abundance indices. *Canadian Journal of Fisheries and Aquatic Sciences* 67(1), 108-120.

Francis, R.I.C.C. 2011. Data weighting in statistical fisheries stock assessment models. *Canadian Journal of Fisheries and Aquatic Sciences*. 68(6): 1124-1138.

Northeast Fisheries Science Center (NEFSC). 2019. 66th Northeast Regional Stock Assessment Workshop (66th SAW) Assessment Report. US Dept. Commer., Northeast Fish. Sci. Cent. Ref. Doc. 19-08; 1170 p.

**List of Appendices (which can be found [here](#))**

Appendix 1: Model structure and detailed results for the base model run.

Appendix 2. Diagnostic plots for the base model in which new 2020-2021 selectivity blocks were added for the Bay and Ocean regions.

Appendix 3. Diagnostic plots and results for a model run in which a new 2020-2021 selectivity block was added for the Ocean region only.

Appendix 4. Diagnostic plots and results from the SCA model with no new selectivity blocks added to the model.

**TABLES**

**Table 1. Total removals by fleet in numbers of fish**

<b>Year</b>	<b>Bay Fleet</b>	<b>Ocean Fleet</b>	<b>Total Removals</b>
<b>1982</b>	229,161	677,600	906,761
<b>1983</b>	339,515	709,879	1,049,394
<b>1984</b>	479,009	357,555	836,564
<b>1985</b>	48,686	853,917	902,603
<b>1986</b>	100,649	307,312	407,961
<b>1987</b>	44,939	231,939	276,878
<b>1988</b>	124,365	332,720	457,085
<b>1989</b>	85,092	521,339	606,431
<b>1990</b>	663,884	574,713	1,238,597
<b>1991</b>	790,833	927,478	1,718,311
<b>1992</b>	986,955	1,243,234	2,230,189
<b>1993</b>	941,415	1,088,947	2,030,362
<b>1994</b>	1,326,775	1,585,122	2,911,897
<b>1995</b>	1,978,738	3,049,239	5,027,977
<b>1996</b>	2,514,266	3,749,942	6,264,208
<b>1997</b>	3,166,575	4,214,559	7,381,134
<b>1998</b>	2,949,332	4,961,986	7,911,318
<b>1999</b>	3,195,145	4,867,163	8,062,308
<b>2000</b>	3,432,148	4,955,360	8,387,508
<b>2001</b>	2,586,938	5,184,845	7,771,783
<b>2002</b>	2,673,581	5,513,147	8,186,728
<b>2003</b>	3,333,975	5,528,236	8,862,211
<b>2004</b>	3,327,387	6,195,000	9,522,387
<b>2005</b>	2,971,213	6,137,340	9,108,553
<b>2006</b>	4,083,679	6,983,996	11,067,675
<b>2007</b>	3,162,774	5,132,018	8,294,792
<b>2008</b>	2,630,471	5,592,223	8,222,694
<b>2009</b>	3,151,161	4,880,287	8,031,448
<b>2010</b>	2,936,586	5,433,285	8,369,871
<b>2011</b>	2,520,001	5,037,736	7,557,737
<b>2012</b>	2,671,307	4,411,580	7,082,887
<b>2013</b>	2,752,138	5,754,205	8,506,343
<b>2014</b>	3,231,424	3,839,183	7,070,607
<b>2015</b>	2,788,075	3,315,477	6,103,552
<b>2016</b>	3,589,860	3,601,305	7,191,165
<b>2017</b>	2,495,418	4,553,797	7,049,215
<b>2018</b>	2,367,605	3,420,077	5,787,682
<b>2019</b>	2,114,336	3,344,764	5,459,100
<b>2020</b>	2,006,072	3,080,791	5,086,863
<b>2021</b>	1,633,797	3,510,737	5,144,534

**Table 2. Contribution of imputed data to 2020 MRIP catch rate estimates by state. Imputed data were 2018 and 2019 intercepts that were used to supplement 2020 APAIS data in strata that were under-sampled due to COVID-19.**

<b>State</b>	<b>Harvest Rate (A+B1)</b>	<b>Released Alive Rate (B2)</b>
<b>Maine</b>	0%	0%
<b>New Hampshire</b>	15%	7%
<b>Massachusetts</b>	3%	3%
<b>Rhode Island</b>	0%	13%
<b>Connecticut</b>	77%	56%
<b>New York</b>	53%	9%
<b>New Jersey</b>	51%	32%
<b>Delaware</b>	49%	13%
<b>Maryland</b>	9%	7%
<b>Virginia</b>	7%	36%
<b>North Carolina (ocean only)</b>	--	72%
<b>Coastwide</b>	29%	15%

**Table 3. Summary of indices used in the striped bass stock assessment model.**

Index Name	Index Metric	Design	Time of		
			Year	Years	Age
MRIP Total Catch Rate Index	Total catch per unit effort	Stratified random	Mar-Dec	1982-2021	1+
Connecticut Long Island Sound Trawl Survey (CTLISTS)	Mean number per tow	Stratified random	Apr-Jun	1984-2021	1+
New York Ocean Haul Seine (NYOHS)	Geometric mean per haul	Fixed station	Sep-Oct	1987-2006	1+
New York Young-of-the-Year (NYYOY)	Geometric mean per haul	Fixed station	Jul-Nov	1985-2021	YOY
New York Western Long Island Beach Seine Survey (NY Age-1)	Geometric mean per haul	Fixed station	May-Aug	1984-2021	1
New Jersey Bottom Trawl Survey (NJTRL)	Stratified mean per tow	Stratified random	April	1990-2018	1+
New Jersey Young-of-the-Year Survey (NJYOY)	Geometric mean per haul	Fixed station	Aug-Oct	1982-2021	YOY
Delaware Spawning Stock Electrofishing Survey (DESSN)	Geometric mean per tow	Fixed station	Apr-Jun	1996-2021	1+
Delaware 30' Bottom Trawl Survey (DE30)	Geometric mean per tow	Fixed station	Nov-Dec	1990-2021	1+
Maryland Spawning Stock Survey (MDSSN)	Selectivity-corrected CPUE	Stratified random	Mar-May	1985-2021	1+
Maryland Young-of-the-Year and Yearlings Surveys (MDYOY and MD Age-1)	Geometric mean per haul	Fixed station	Jul-Sep	1954-2021	0-1
Virginia Young-of-the-Year Survey (VAYOY)	Geometric mean per haul	Fixed station	Jul-Sep	1980-2021	YOY
Chesapeake Bay Multispecies Monitoring and Assessment Program (ChesMMAP)	Stratified mean per tow	Stratified random	Mar-Nov	2002-2018	1+



**Table 4. Status of age-1+ striped bass surveys from 2018-2021. Empty cells indicate the survey occurred without interruption.**

Year	CT LISTS	NJ TRL	DE SSN	DE 30'	MD SSN	ChesMMAP
2018						
2019		Did not occur				Unavailable
2020	Did not occur	Did not occur	Did not occur			Unavailable
2021		Did not occur			Delayed	Unavailable

**Table 5. Status of striped bass recruitment surveys from 2018-2021. Empty cells indicate the survey occurred without interruption.**

Year	NY JAI	NY Age-1	NJ JAI	MD JAI	MD Age-1	VA JAI
2018						
2019						
2020		Interrupted	Did not occur			
2021						

**Table 6. Model structure summary for the 2021 striped bass update.**

	Value(s)
<b>Years in Model</b>	1982-2021
<b>Size/Age Plus Group</b>	15+
<b>Fleets</b>	2 (Bay and Ocean)
<b>Selectivity blocks</b>	Bay fleet: 1982-1984, 1985-1989, 1990-1995, 1996-2019, 2020-2021 Ocean fleet: 1982-1984, 1985-1989, 1990-1996, 1997-2019, 2020-2021

**Table 7. Striped bass life history information used in the 2021 stock assessment update.**

<b>Age</b>	<b>Proportion Mature</b>	<b>Proportion Female</b>	<b>Natural Mortality</b>
<b>1</b>	0	0.53	1.13
<b>2</b>	0	0.56	0.68
<b>3</b>	0	0.56	0.45
<b>4</b>	0.09	0.52	0.33
<b>5</b>	0.32	0.57	0.25
<b>6</b>	0.45	0.65	0.19
<b>7</b>	0.84	0.73	0.15
<b>8</b>	0.89	0.81	0.15
<b>9</b>	1	0.88	0.15
<b>10</b>	1	0.92	0.15
<b>11</b>	1	0.95	0.15
<b>12</b>	1	0.97	0.15
<b>13</b>	1	1	0.15
<b>14</b>	1	1	0.15
<b>15+</b>	1	1	0.15

**Table 8. Population estimates from the 2021 striped bass assessment update.**

<b>Year</b>	<b>Full <i>F</i></b>	<b>Recruitment (millions of age-1 fish)</b>	<b>Female SSB (mt)</b>	<b>Total Abundance (millions of fish)</b>	<b>Age 8+ Abundance (millions of fish)</b>
1982	0.17	36.2	18,498	54.5	1.7
1983	0.14	70.1	15,614	92.4	1.5
1984	0.07	60.5	15,783	95.8	1.3
1985	0.19	66.8	16,452	106.2	1.5
1986	0.05	64.5	14,838	109.0	1.7
1987	0.03	71.2	18,247	118.9	2.0
1988	0.04	92.5	24,125	145.2	2.5
1989	0.05	104.6	36,060	167.5	3.3
1990	0.07	128.3	42,017	201.1	5.3
1991	0.09	100.6	49,377	186.7	6.5
1992	0.11	106.0	62,663	190.7	7.5
1993	0.09	131.1	70,390	217.9	8.0
1994	0.12	285.6	79,213	382.5	8.6
1995	0.21	184.3	85,457	336.1	9.6
1996	0.27	232.1	95,380	378.2	9.9
1997	0.21	261.2	90,227	422.4	10.2
1998	0.22	147.1	83,863	325.8	9.7
1999	0.21	152.1	83,024	304.0	9.3
2000	0.21	121.4	95,101	263.3	9.7
2001	0.20	192.2	99,421	318.3	13.6
2002	0.22	228.7	111,329	369.2	14.1
2003	0.24	118.3	113,506	276.1	15.3
2004	0.26	323.3	109,337	453.8	16.6
2005	0.26	157.0	108,416	340.1	14.5
2006	0.30	138.7	102,105	293.5	13.1
2007	0.23	81.2	99,830	216.9	10.9
2008	0.24	131.8	106,075	240.7	11.6
2009	0.23	70.6	104,599	181.2	12.8
2010	0.27	92.3	104,749	182.0	11.4
2011	0.28	118.3	97,556	203.0	14.5
2012	0.28	208.6	95,936	297.5	12.8
2013	0.39	63.6	84,750	182.7	11.2
2014	0.31	76.9	73,346	162.6	8.1
2015	0.27	152.4	63,415	228.0	7.5
2016	0.31	238.7	64,227	333.0	6.2
2017	0.35	101.7	57,106	231.5	5.6
2018	0.26	130.7	55,120	234.8	5.4
2019	0.23	159.6	56,634	263.7	7.4
2020	0.14	109.5	59,980	223.1	6.4
2021	0.14	116.0	64,805	218.9	6.6


**Table 9. Updated biological reference points and 2021 estimates for *F* and female SSB compared with the estimates from the 2018 benchmark.**

Metric	2018		2021		2021 Value
	2018 Target	Threshold	2021 Target	Threshold	
<b>Fishing Mortality</b>	0.20	0.24	0.17	0.20	0.14
<b>Female SSB</b>	114,295 mt (252 million lbs)	91,436 mt (202 million lbs)	106,820 mt (235 million lbs)	85,457 mt (188 million lbs)	64,805 mt (143 million lbs)


**Table 10. Probability of SSB being at or above the SSB threshold or target under different constant *F* scenarios. Shaded row indicates 2029, the rebuilding deadline.**

Year	Probability SSB ≥ SSB threshold under current <i>F</i>	Probability SSB ≥ SSB target under current <i>F</i>	Probability SSB ≥ SSB threshold under <i>F</i> target	Probability SSB ≥ SSB target under <i>F</i> target	Probability SSB ≥ SSB threshold under <i>F</i> threshold	Probability SSB ≥ SSB target under <i>F</i> threshold
2021	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2022	34.4%	0.4%	34.5%	0.4%	34.5%	0.4%
2023	70.2%	14.9%	61.9%	13.1%	53.2%	11.6%
2024	86.0%	39.0%	74.1%	29.2%	61.8%	23.2%
2025	91.8%	56.1%	79.3%	40.3%	64.3%	28.6%
2026	94.1%	65.7%	81.4%	45.5%	63.4%	30.3%
2027	95.7%	72.7%	82.8%	49.9%	63.4%	31.9%
2028	96.4%	76.6%	82.8%	52.0%	61.7%	31.6%
2029	96.7%	78.6%	82.4%	52.5%	59.4%	30.5%
2030	97.0%	80.6%	82.8%	53.7%	58.6%	30.5%


## FIGURES




**Figure 1. Total striped bass removals by fleet.**



**Figure 2. Total striped bass removal by sector.**



**Figure 3. Indices of age-1+ abundance for striped bass, 1982-2021.**



**Figure 4. Striped bass juvenile abundance indices, including the composite Chesapeake Bay index (MD-VA), 1954-2021.**






Figure 5. Age-1 recruitment indices for striped bass, 1954-2021.







Figure 6. Selectivity patterns for the Bay fleet (top) and the Ocean fleet (bottom).




**Figure 7. Fully recruited fishing mortality for the Bay and Ocean fleets plotted with the total fully recruited  $F$ .**




**Figure 8. Estimates of striped bass recruitment plotted with the time series mean.**




**Figure 9. Total abundance (top) and age-8+ abundance of striped bass over time.**



**Figure 10. Female spawning stock biomass (top) and exploitable biomass of striped bass over time.**




**Figure 11. Retrospective plots of seven-year peels for fishing mortality (top), female spawning stock biomass (middle), and recruitment (bottom).**




**Figure 12. Comparison of fishing mortality (top), female SSB (middle), and recruitment (bottom) estimates from the update assessment and an assessment in which the 2020 NY Age 1 and 2021 MDSSN index values were set as missing. Absolute values are on the left and relative percent difference is on the right.**






**Figure 13. Comparison of fully-recruited fishing mortality (top) and female SSB (bottom) from the update assessment base model and sensitivity runs with a new 2020-2021 selectivity block for the Ocean region only and no new selectivity blocks.**



**Figure 14. Comparison of estimates of female spawning stock biomass (top) and total fishing mortality (bottom) from the 2018 benchmark assessment and current assessment update.**



**Figure 15. 2020-2021 average selectivity pattern used in the projections to determine fishing mortalities associated with the SSB threshold and targets compared to the overall selectivity in each individual year.**






Figure 16. Plot comparing the 2021 retrospective-adjusted  $F$  and female SSB values with the unadjusted  $F$  and SSB estimates and their associated 90% confidence intervals.



**Figure 17. Female SSB (top) and total F estimates (bottom) plotted with their respective targets and thresholds. Shaded area indicates 95% confidence intervals of the estimates.**



**Figure 18.** Projections of female spawning stock biomass through 2030 under current  $F$  (top), target  $F$  (middle), and threshold  $F$  (bottom). Absolute values are on the left and the probability of female SSB being above the target and threshold values is on the right.