

# A. STOCK ASSESSMENT FOR BLACK SEA BASS FOR 2016

## TERMS OF REFERENCE

TOR 1. Summarize the conclusions of the February 2016 SSC peer review regarding the potential for spatial partitioning of the black sea bass stock. (The consequences for the stock assessment will be addressed in TOR-6.)

TOR 2. Estimate catch from all sources including landings and discards. Characterize the uncertainty in these sources of data. Evaluate available information on discard mortality and, if appropriate, update mortality rates applied to discard components of the catch. Describe the spatial and temporal distribution of fishing effort.

TOR 3. Present the survey data being used in the assessment (e.g., indices of abundance, recruitment, state surveys, age-length data, etc.). Investigate the utility of fishery dependent indices as a measure of relative abundance. Characterize the uncertainty and any bias in these sources of data.

TOR 4. Consider the consequences of environmental factors on the estimates of abundance or relative indices derived from surveys.

TOR 5. Investigate implications of hermaphroditic life history on stock assessment model. If possible, incorporate parameters to account for hermaphroditism.

TOR 6. Estimate annual fishing mortality, recruitment and stock biomass (both total and spawning stock), using measures that are appropriate to the assessment model, for the time series (integrating results from TORs-1,-4, & -5 as appropriate), and estimate their uncertainty. Include a historical retrospective analysis and past projection performance evaluation to allow a comparison with most recent assessment results.

TOR 7. Estimate biological reference points (BRPs; point estimates or proxies for BMSY, BTHRESHOLD, FMSY, and MSY), including defining BRPs for spatially explicit areas if appropriate, and provide estimates of their uncertainty. If analytic model-based estimates are unavailable, consider recommending alternative measurable proxies for BRPs. Comment on the appropriateness of existing BRPs and the “new” (i.e., updated, redefined, or alternative) BRPs.

TOR 8. Evaluate overall stock status with respect to a new model or new models that considered spatial units developed for this peer review.

TOR 9. Develop approaches and apply them to conduct stock projections.

- a. Provide numerical annual projections (3-5 years) and the statistical distribution (e.g., probability density function) of the OFL (overfishing level) that fully incorporates observation, process and model uncertainty (see Appendix to the SAW TORs). Each projection should estimate and report annual probabilities of exceeding threshold BRPs for F, and probabilities of falling below threshold BRPs for biomass. Use a sensitivity

analysis approach in which a range of assumptions about the most important uncertainties in the assessment are considered (e.g., terminal year abundance, variability in recruitment, and definition of BRPs for black sea bass).

b. Comment on which projections seem most realistic. Consider major uncertainties in the assessment as well as the sensitivity of the projections to various assumptions.

c. Describe this stock's vulnerability (see "Appendix to the SAW TORs") to becoming overfished, and how this could affect the choice of ABC.

TOR 10. Review, evaluate and report on the status of the SARC and Working Group research recommendations listed in recent SARC reviewed assessments and review panel reports. Identify new research recommendations.

### **Appendix to the SAW Assessment TORs: Clarification of Terms used in the SAW/SARC Terms of Reference**

#### **Guidance to SAW WG about "Number of Models to include in the Assessment Report":**

In general, for any TOR in which one or more models are explored by the WG, give a detailed presentation of the "best" model, including inputs, outputs, diagnostics of model adequacy, and sensitivity analyses that evaluate robustness of model results to the assumptions. In less detail, describe other models that were evaluated by the WG and explain their strengths, weaknesses and results in relation to the "best" model. If selection of a "best" model is not possible, present alternative models in detail, and summarize the relative utility each model, including a comparison of results. It should be highlighted whether any models represent a minority opinion.

#### **On "Acceptable Biological Catch" (DOC Nat. Stand. Guidel. Fed. Reg., v. 74, no. 11, 1-16-2009):**

*Acceptable biological catch (ABC)* is a level of a stock or stock complex's annual catch that accounts for the scientific uncertainty in the estimate of Overfishing Limit (OFL) and any other scientific uncertainty..." (p. 3208) [In other words,  $OFL \geq ABC$ .]

*ABC for overfished stocks.* For overfished stocks and stock complexes, a rebuilding ABC must be set to reflect the annual catch that is consistent with the schedule of fishing mortality rates in the rebuilding plan. (p. 3209)

NMFS expects that in most cases ABC will be reduced from OFL to reduce the probability that overfishing might occur in a year. (p. 3180)

ABC refers to a level of "catch" that is "acceptable" given the "biological" characteristics of the stock or stock complex. As such, Optimal Yield (OY) does not equate with ABC. The specification of OY is required to consider a variety of factors, including social and economic factors, and the protection of marine ecosystems, which are not part of the ABC concept. (p. 3189)

**On “Vulnerability” (DOC Natl. Stand. Guidelines. Fed. Reg., v. 74, no. 11, 1-16-2009):**

“*Vulnerability*. A stock’s vulnerability is a combination of its productivity, which depends upon its life history characteristics, and its susceptibility to the fishery. Productivity refers to the capacity of the stock to produce Maximum Sustainable Yield (MSY) and to recover if the population is depleted, and susceptibility is the potential for the stock to be impacted by the fishery, which includes direct captures, as well as indirect impacts to the fishery (e.g., loss of habitat quality).” (p. 3205)

**Participation among members of a Stock Assessment Working Group:**

Anyone participating in SAW meetings that will be running or presenting results from an assessment model is expected to supply the source code, a compiled executable, an input file with the proposed configuration, and a detailed model description in advance of the model meeting. Source code for NOAA Toolbox programs is available on request. These measures allow transparency and a fair evaluation of differences that emerge between models.

## **Executive Summary**

**Term of reference 1. Summarize the conclusions of the February 2016 SSC peer review regarding the potential for spatial partitioning of the black sea bass stock. (The consequences for the stock assessment will be addressed in TOR-6.)**

The Black Sea Bass Working Group (WG) presented information to a review panel of the Mid-Atlantic Fishery Management Council Science and Statistical Committee (MAFMC SSC) regarding spatial patterns in the black sea bass northern stock. The WG concluded that recognizing two distinct areas separated by Hudson Canyon would provide the best spatial partitioning for the stock assessment. The WG emphasized that the two areas do not constitute discrete stock areas. The review panel agreed with this spatial boundary as a starting point for evaluation in the assessment.

**Term of reference 2. Estimate catch from all sources including landings and discards. Characterize the uncertainty in these sources of data. Evaluate available information on discard mortality and, if appropriate, update mortality rates applied to discard components of the catch. Describe the spatial and temporal distribution of fishing effort.**

The principal gears used in commercial fishing for black sea bass are fish pots, otter trawls and hand-lines. Commercial landings peaked in 1952 at 9,900 mt then declined substantially during the 1960s until commercial landings during the late 1980s and 1990s averaged 1,300 mt. Commercial fishery quotas were implemented in 1998 but commercial landings remained stable between 1,300 mt and 1,600 mt until 2007. After quota restrictions, commercial landings declined to 523 and 751 mt in 2009 and 2010, respectively. Commercial landings increased to 1,088 mt in 2013 and have since remained above 1,000 mt (1,113 mt in 2015). The recreational rod-and-reel fishery for black sea bass harvests a significant proportion of the total catch. After peaking in 1986, recreational landings averaged 1,700 mt annually until 1997. Recreational fishery harvest limits were implemented in 1998 and recreational landings have since ranged between 500 mt and 2,000 mt. Recreational landings in 2015 were 1,864 mt. Commercial fishery discards, although poorly estimated, appear to be a minor part of the total fishery removals from the stock. Commercial discards were generally less than 200 mt per year, but increased to 416 mt and 335 mt in 2014 and 2015, respectively. Recreational discard losses

assuming 15% hook and release mortality are similar, generally less than 500 mt per year. Estimated mortality from recreational discards was 371 mt in 2015. The distributions of the commercial and recreational fisheries were described and maps were produced based on observed trips and vessel trip reports.

**Term of reference 3. Present the survey data being used in the assessment (e.g., indices of abundance, recruitment, state surveys, age-length data, etc.). Investigate the utility of fishery dependent indices as a measure of relative abundance. Characterize the uncertainty and any bias in these sources of data.**

The 2016 assessment model for black sea bass applied spatially explicit age-based statistical catch at age models using catch and age information since 1989. The fishery catch in two areas (north and south of Hudson Canyon) is modeled as two fleets (trawl and non-trawl) with area specific indices of stock abundance from NEFSC winter and spring surveys, the NEAMAP spring survey, recreational catch per angler, as well as state survey indices from VA, MD, DE, NJ in the south and NY, CT, RI and MA in the northern area. Several state survey index series were modeled to standardize factors that influence the indices. The standardized values were used as relative indices of abundance. Indices in the north have increased over the past decade and show a very strong 2011 year class. Indices in the southern area indicated a strong 1999 cohort, however the 2011 cohort is only average in size for the southern area. Recreational catch per angler trip (CPA) was developed using effort targeting a suite of species commonly caught with sea bass and was used as a relative index of abundance.

**Term of reference 4. Consider the consequences of environmental factors on the estimates of abundance or relative indices derived from surveys.**

The significant factors in the state surveys were depth, salinity and temperature. In response to issues raised in previous black sea bass assessments, a study was also conducted to evaluate the influence of winter oceanographic conditions on distribution and survival. The study concluded that warm saline conditions improved juvenile survival and the location of the shelf-slope front dictates the distribution of adults in the winter offshore habitat.

**Term of reference 5. Investigate implications of hermaphroditic life history on stock assessment model. If possible, incorporate parameters to account for hermaphroditism.**

In response to previous research recommendations the implication of black sea bass life history on exploitation, was evaluated in a simulation study. The study concluded that black sea bass in the northern unit stock was more robust to exploitation than typical hermaphroditic species depending on spawning contributions from secondary males. Given these findings and the findings of previously published studies, black sea bass spawning stock biomass (SSB) was defined as male and female mature biomass. Since growth between males and female black sea bass is similar, the WG concluded that use of male and female SSB accounted for the unique life history of black sea bass. Additionally, an operating model conditioned on the available black sea bass data including sex structure and hermaphroditism was used to simulate data for performance testing of alternative estimation models. The pooled-sex modeling approach developed by the WG performed relatively well for determining stock status.

**Term of reference 6. Estimate annual fishing mortality, recruitment and stock biomass (both total and spawning stock), using measures that are appropriate to the assessment model, for the time series (integrating results from TORs 1,4, & 5 as appropriate), and estimate their uncertainty. Include a historical retrospective analysis and past projection performance evaluation to allow a comparison with most recent assessment results.**

Fishing mortality (F) from the combined north and south areas varied between  $F = 1.34$  (1993) and  $F = 0.24$  (2015). The mortality averaged greater than 1.0 prior to implementation of management regulations in 1997. Fishing mortality remained above 0.50 between 1998 and 2008 then declined steadily to 0.24 in 2015 (value not adjusted for retrospective bias). Both the north and south area models had a pattern of retrospective bias. However the direction of patterns was different between areas, with apparently over-estimated fishing mortality (F) in the north and apparently under-estimated F in the south. A combination of retrospective adjusted Fs from each area produced a 2015  $F = 0.27$ . Spawning stock biomass (SSB) increased from about 2,485 mt in 1991 to about 8,500 mt in 2002, then decreased to about 4,072 mt by 2007. With improved recruitment and declining fishing mortality rates, SSB has steadily increased since 2007. Total spawning biomass peaked in 2014 at 17,158 mt then declined in 2015 to 16,552 mt (value not adjusted for retrospective bias) as the 2011 cohort abundance declined. Total biomass

increased from 6,558 mt in 1989 to 16,205 mt in 2002, but decreased to 9,777 mt by 2006. However, beginning in 2007 biomass began increasing, peaking at 27,125 mt in 2014. Total biomass was slightly lower in 2015 at 24,143 mt (retrospective bias adjusted 2015 biomass equaled 32,010 mt) while the SSB retrospective adjusted value equaled 22,176 mt. Recruitment at age 1 averaged 24.318 million fish from 1989 to 2015, with peaks in 1991 (29.149 million, 1990 cohort), 1994 (27.188 million, 1994 cohort), 2000 (37.256 million, 1999 cohort) and most importantly in 2012 (68.932 million) from the 2011 cohort. The distribution of the 2011 cohort was more prevalent in the northern area of the stock. The 2014 cohort in 2015 was estimated as 24.917 million fish with a retrospective adjusted recruitment equal to 18.002 million fish. Model results in both areas show a generally decreasing trend in fishing mortality but different historical trends in recruitment and spawning stock biomass. Recent increased biomass occurred in both areas and relatively high biomass in the last year. Several alternative modeling approaches produced similar results, suggesting that the results are robust to a range of data and model decisions.

**Term of reference 7. Estimate biological reference points (BRPs; point estimates or proxies for BMSY, BTHRESHOLD, FMSY, and MSY), including defining BRPs for spatially explicit areas if appropriate, and provide estimates of their uncertainty. If analytic model-based estimates are unavailable, consider recommending alternative measurable proxies for BRPs. Comment on the appropriateness of existing BRPs and the “new” (i.e., updated, redefined, or alternative) BRPs.**

The 2008 Northeast Data Poor Stocks Working Group (NEDPSWG) Review Panel (NEFSC 2009a) recommended  $F_{40\%}$  be used as a proxy for  $F_{MSY}$  and total spawning stock biomass at  $F_{40\%}$  ( $SSB_{40\%}$ ) be used as the proxy for the stock biomass target reference point. Considering the weak stock-recruitment relationship in the range of estimated stock sizes and biological reference points for comparable species,  $F_{40\%}$  was maintained as the proxy for black sea bass  $MSY$  related reference points in the 2016 assessment. The average of the two area specific  $F_{40\%}$  values produced a  $F_{MSY}$  proxy of 0.36. Based on a long-term projection at  $F_{MSY}$  proxy, the associated  $SSB_{MSY}$  proxy equaled 9,667 mt and  $B_{MSY}$  proxy equaled 17,256 mt.

**Term of reference 8. Evaluate overall stock status with respect to a new model or new models that considered spatial units developed for this peer review.**

Stock status was evaluated using the combined results of the two area models (2015 retrospective adjusted average fishing mortality  $F=0.27$ ) compared to the area-averaged  $F_{MSY\ proxy}$  (0.36).

Results indicate that overfishing is not occurring on the stock. The 2015 combined retrospective adjusted total biomass was (32,010 mt) was much greater than the  $B_{MSY\ proxy}$  (17,256 mt) and biomass threshold of 8,628 mt. In addition, the combined retrospective adjusted SSB of 22,176 mt was well above the  $SSB_{MSY\ proxy}$  of 9,667 mt and SSB proxy threshold of 4,834 mt.

Therefore the stock is not considered over-fished.

**Term of reference 9. Develop approaches and apply them to conduct stock projections.**

**a. Provide numerical annual projections (3-5 years) and the statistical distribution (e.g., probability density function) of the OFL (overfishing level) that fully incorporates observation, process and model uncertainty (see Appendix to the SAW TORs). Each projection should estimate and report annual probabilities of exceeding threshold BRPs for F, and probabilities of falling below threshold BRPs for biomass. Use a sensitivity analysis approach in which a range of assumptions about the most important uncertainties in the assessment are considered (e.g., terminal year abundance, variability in recruitment, and definition of BRPs for black sea bass).**

**b. Comment on which projections seem most realistic. Consider major uncertainties in the assessment as well as the sensitivity of the projections to various assumptions.**

**c. Describe this stock's vulnerability (see "Appendix to the SAW TORs") to becoming overfished, and how this could affect the choice of ABC.**

Projections were made for two fishing mortality scenarios. One assumes that fishing mortality from 2017 to 2019 equaled the  $F_{MSY\ proxy}$ , and a second scenario assumes that  $F$  equals to the 2015 terminal year status quo value (North  $F_{SQ}=0.14$  and South  $F_{SQ}=0.39$ ). Future annual recruitments were drawn from empirical values from 2000 to 2015 and catch in 2016 was assumed equal to the allowable biological catch (3,024 mt). Starting stock size was based on area-specific estimates of 2016 abundance at age with retrospective adjustments and recent selectivity estimates.  $F_{MSY}$  projections suggest that spawning biomass would decline to 11,849 mt by 2019 and total biomass would equal 20,788 mt by 2019. Total catch under that scenario



would increase to 5,467 mt in 2017 then decline to 3,901 mt by 2019.  $F_{SQ}$  projections suggest that combined area spawning biomass would decline to 15,349 mt by 2019 and total biomass would equal 24,704 mt by 2019. Total catch under that scenario would decrease to 3,024 mt in 2017 then decline to 2,764 mt by 2019.

The projections were developed for a single stock using the combined results of the north and south areas. It was clear from the results that the area specific conditions vary and therefore the vulnerability of each area to exploitation at a common  $F$  may vary.

**Term of reference 10. Review, evaluate and report on the status of the SARC and Working Group research recommendations listed in recent SARC reviewed assessments and review panel reports. Identify new research recommendations.**

The WG reviewed previous research recommendations and concluded they had been considered and when possible completed. New recommendations for future consideration include:

expand on previous genetic studies with smaller spatial increments in sampling; consider the impact of climate change on black sea bass, particularly in the Gulf of Maine; evaluate population sex change and sex ratio, particularly comparing dynamics among communities; study black sea bass catchability in a variety of survey gear types; investigate and document social and spawning dynamics of black sea bass; increase work to understand habitat use in sea bass and seasonal changes; evaluate use of samples collected by industry study fleets.

**Black Sea Bass Working Group :**

The SARC62 Black Sea Bass Working Group conducted two data meetings (June 29-July 2 2015, Warwick RI; June 27-29, 2016, Providence RI), a spatial structure workshop (December 16 2015, Warwick RI), one model meeting (September 6-9 2016, Woods Hole MA), and several video conference calls in the development of this assessment. The formal working group consisted of:

John Maniscalco – New York Dept. of Environmental Conservation - chair

Gary Shepherd – Northeast Fisheries Science Center –assessment lead

Mike Bednarski – Massachusetts Division of Marine Fisheries (currently Virginia Dept. Game and Inland Fisheries)

Jeffrey Brust – New Jersey Department of Environmental Protection

Steve Cadrin – UMass Dartmouth

Steve Doctor – Maryland Dept. of Natural Resources - Fisheries Service

Gavin Fay – UMass Dartmouth

Robert Leaf – University of Southern Mississippi

Chris Legault – Northeast Fisheries Science Center

Jason McNamee – Rhode Island DEM- Division of Fish and Wildlife, Marine Fisheries

Other participants included Kirby Rootes-Murdy (ASMFC), Kiley Dancy (MAFMC), Kiersten Curti (NEFSC), Olaf Jensen (Rutgers Univ.), Joe Myers (ACCSP), Greg Wojcik (CT), Peter Clarke (NJDEP), Tom Wadsworth (NCDMF), Greg DeCelles (UMass Dartmouth), Mark Hoffman (AP), Mark Terceiro (NEFSC), Rich Seagraves (MAFMC), Moira Kelly (GARFO), Rich Wong (DEDFW), Alicia Miller (NEFSC), Marissa McMahan (Northeastern Univ.), Eric Powell (SCEMFIS), Alexei Sharov (MDDNR), Mike Palmer (NEFSC), Paul Nitschke (NEFSC), Chuck Adams (NEFSC), Tim Miller (NEFSC), Jeff Kipp (ASMFC), Jessica Blaylock (NEFSC).

## INTRODUCTION

### *Life History*

Black sea bass (*Centropristis striata*) are distributed from the Gulf of Maine to the Gulf of Mexico, but fish north of Cape Hatteras, NC are considered part of a single unit stock (Figure A1). Over the past decade, the distribution of sea bass has expanded into the Gulf of Maine (Bell et al. 2014) as far as eastern coastal Maine (M. McMahan, Northeastern Univ, pers. comm.). Within the stock area, distribution changes on a seasonal basis and the extent of the seasonal change varies by location. In the northern end of the range (New York to Massachusetts), black sea bass move offshore crossing the continental shelf, then south along the edge of the shelf (Moser and Shepherd 2009). By late winter, northern fish may travel as far south as Virginia, but most return to the northern inshore areas by May. Black sea bass originating inshore along the Mid-Atlantic coast (New Jersey to Maryland) head offshore to the shelf edge during late autumn, travelling in a southeasterly direction. They return inshore in spring to the general area from which they originated. Black sea bass in the southern extent of the stock (Virginia and

North Carolina) move offshore in late autumn/early winter. Given the proximity of the shelf edge, they transit a relatively short distance, due east, to reach over-wintering areas.

Black sea bass are protogynous hermaphrodites and can be categorized as temperate reef fishes (Steimle et al. 1999, Drohan et al. 2007). Transition from female to male generally occurs between the ages of two and five years (Lavenda 1949, Mercer 1978). Based on sex ratio at length from NMFS surveys, males constitute approximately 25% of the population by 15 cm, with increasing proportions of males with size (Figure A2). Following transition from female to male, sea bass can follow one of two behavioral pathways; either becoming a dominant male, characterized by a larger size and a bright blue nuchal hump during spawning season, or subordinate males (secondary males) which have few distinguishing features from females. The initiation of sexual transition appears to be based on visual rather than chemical cues (Dr. David Berlinsky, UNH, Personal communication). In studies of protogyny, among several coral reef fish species, transition of the largest female to male may occur quickly if the dominant male is removed from the reef, however, similar studies have not been published for black sea bass.

Spawning in the Middle Atlantic peaks during spring (May and June) when the fish reside in coastal waters (Drohan et al. 2007). The social structure of the spawning aggregations is poorly known although some observations suggest that large dominant males gather a harem of females and aggressively defend territory during spawning season (Nelson et al. 2003). The bright coloration of males during spawning season suggests that visual cues may be important in structuring of the social hierarchy. Black sea bass populations may also have secondary males present which are capable of spawning but do not present the same morphological features as dominant males (NEFSC 2012). Recent analysis of morphological features in mature female, primary and secondary male black sea bass showed that mature secondary males may be indistinguishable from mature females (Keigwin et al. 2016).

Age estimation for black sea bass has been routinely conducted at the NEFSC and other institutions (e.g. Virginia Institute of Marine Science (VIMS)) using both scales and otoliths. Validation of the ages and comparison between age readers is presented in Appendix A1. The study concluded that age determinations were valid using either structure, but older ages were easier to determine from otoliths. Some discrepancy remains between readers determining age 0 vs age 1 in fall samples. Consequently and for other reasons described below, fall indices were

not used in the stock assessment analysis. Any fall catch data age 0 or 1 were based on NEFSC age data.

Black sea bass attain a maximum size around 60 cm and 4 kg. Published growth curves from Lavenda (1949) suggested a maximum age for females of 8 and age 12 for males. However he noted the presence of large males (>45 cm) in deeper water that may have been older. Von Bertalanffy growth curves from NEFSC age and growth data produce estimates of  $L_{\infty}$  equal to 58.9 cm with  $K$  and  $t_0$  of 0.22 and 0.207, respectively (Figure A3). Length at 50% maturity for both sexes combined occurs at 21.0 cm (age2), based on NEFSC maturity data from spring and winter surveys since 1984 (Figure A4). Black sea bass reach full maturity by 35 cm (age 5) (Figure A5).

### *Habitat*

Black sea bass are commonly associated with live-bottom and reef habitats although use a variety of habitats throughout the year. Habitat use by life stage and season was summarized by Steimle et al. (1999) and provided in Table A1. Fabrizio et al. (2013,2014) examined habitat use and home range with acoustic tags within a defined area of complex habitat. The results showed that black sea bass were relatively mobile, moving among habitats from structure (e.g. rocks, wrecks, reefs) to coarse bottom until leaving the area during the annual migration. A NEFSC outreach project prior to this assessment requested information from stakeholders on habitats where black sea bass were found (Appendix A2). Among the 75 respondents, 50 fishermen (including those using net, hook and line and other gear) noted that black sea bass were common on open bottom (not structured).

Past assessment reviewers expressed concern that indices based on trawl gear used inshore cannot be representative of abundance since such gear cannot tow in structured habitat commonly used by black sea bass. However, the NEFSC spring and winter surveys in offshore strata are conducted in the same area as the commercial winter otter trawl fishery, which is towable bottom. An analysis was done in SARC 53 to examine potential biases in the offshore spring survey indices due to avoidance of structure. The frequency of gear damage during tows

was evaluated relative to black sea bass indices to identify tows in structured habitat (Nieland and Shepherd 2011). The analysis concluded that there is no evidence of a bias in black sea bass catches in the offshore strata resulting from structured habitat.

To further address these concerns over the use of inshore trawl surveys to track relative abundance and size/age composition of black sea bass, an in-depth analysis of available data was conducted to compare catch characteristics of black sea bass by different gear types and in different habitats (Appendix A3). Results indicate that both catch rates and size distribution of black sea bass are similar in structured and non-structured habitat. These results are supported by scientific literature, as well as anecdotal information from the NEFSC outreach program, that indicate that black sea bass are not dependent on areas of high relief habitat and commonly occur in areas of open (trawlable) bottom. Additionally, the analysis indicated similarities between the segment of the population sampled by trawl gear and more specialized gear types like fish pots. In fact, trawl gear seemed to sample both small and large fish better than fish pots, again implying that trawl gear is an appropriate sampling gear for black sea bass. Based on the available information and analyzes, it was determined that trawl surveys operating in open bottom areas are sampling the same population of black sea bass as would be found in structured habitat, and as a result, inshore trawl surveys are appropriate for characterizing size structure and relative abundance of black sea bass.

#### *Food habits*

Black sea bass are visual predators and have a generalized carnivorous diet of predominantly motile, benthic species including amphipods, decapods, and fishes (Sedberry 1988). Items consumed by black sea bass depend on life stage and size. Smaller black sea bass consume mostly small crustaceans; including amphipods, isopods, small crabs and shrimp; but also feed on polychaetes. Larger black sea bass feed more on fish and decapod crustaceans; including rock crabs and juvenile American lobster. Other diet items include molluscs (cephalopods) and ascidians (Steimle et al. 1999).

Stomach content data was collected from fish caught during both the spring and fall NEFSC bottom trawl surveys from 1973-1990. Fish remains found in the stomachs of 802 trawl caught black sea bass included Alepocephalids, Atlantic herring, anchovy, sea horses, northern pipefish, lanternfish, sand lance, short-horned sculpin, cusk eel, scup, windowpane flounder, and rock gunnel. The invertebrate content of black sea bass stomachs was very diverse and included

hydrozoans, anthozoans, cerianthids, bryozoans, ctenophores, annelids (including polychaetes), amphipods, isopods, copepods, ostracods, chaetognatha, lancelets, sea spiders, crabs (including cancer, swimming, box, mud, mole, spider, hermit), shrimp, hooded shrimp, krill, mysids, squat lobster, barnacles, molluscs (including bivalves, snails, sea slugs, limpets, pteropods, cephalopods - squid), echinoderms (including brittle star and sea cucumbers) and tunicates (Rountree 1999).

Sampling in the NEFSC Bottom Trawl Survey from 1973-2008 also found black sea bass as prey items in the stomachs of various predators. In order of frequency of occurrence, these included demersal sharks, summer flounder, fourspot flounder, black sea bass, monkfish, spiny dogfish, skate, windowpane, little skate, bluefish, smooth dogfish, red hake, striped bass, silver hake, Atlantic cod and winter skate. Other food web research conducted in Chesapeake Bay found young of the year black sea bass in the stomachs of Age 1 weakfish (Hartman and Brandt 1995).

Sampling of stomach contents from fish caught in the Hudson Raritan Estuary by otter trawl also provides data on black sea bass as both predator and prey. Most of the black sea bass sampled were young of year or juveniles. Prey species were dominated by crustaceans, particularly shrimp, mysids and juvenile rock crab but also included copepods, amphipods, isopods and other juvenile decapods. Several stomachs contained fish species including cunner, goby, menhaden and anchovy. Juvenile black sea bass were found as prey items in the stomachs of summer flounder, striped searobins, northern searobin, clearnose skate, bluefish, and grubby sculpin (Steimle et al. 2000).

The Northeast Area Monitoring and Assessment Program Near Shore Trawl Survey (NEAMAP) samples from Cape Cod, MA south to Cape Hatteras, NC and targets both juvenile and adult fishes. The survey sampled diet composition of 1,245 black sea bass caught by the survey from 2007 through 2013. Gut contents analysis of these black sea bass found 55% crustaceans, 26.7% fishes, 10.8% mollusks, 4.2% miscellaneous, and 2.7% worms by weight. Crustaceans were the largest portion of the diet by weight and rock crabs led the crustaceans in abundance and weight. Bay anchovy was the most frequent of the fishes and the other important food items were squid, mollusks, gastropods, and worms. Sea bass examined were predominately age zero to age four fish.

The Chesapeake Bay Multispecies Monitoring and Assessment Program (ChesMMAP) began in 2002 and uses a large-mesh bottom trawl to sample juvenile-to-adult fishes from the head of Chesapeake Bay at Poole's Island, MD to the mouth of the Bay just outside the Chesapeake Bay Bridge Tunnel. The ChesMMAP survey sampled the diet composition of 233 black sea bass from 2002 to 2013. Black sea bass stomach contents were composed of 70.5% crustaceans, 10.5% fishes, 10.2% miscellaneous, 4.7% worms, and 4.3% molluscs by weight. Crustaceans were dominated by mysids and amphipods. Sea bass examined were predominantly age one and two fish.

The ChesMMAP survey is conducted in the Chesapeake Bay versus the NEAMAP survey which is conducted in near shore waters. Comparison of the two surveys demonstrates the reliance of the smaller fish found in the Chesapeake Bay on mysids, mud crabs, amphipods, and sand shrimp while the larger fish caught in the NEAMAP survey rely more on rock crabs and fishes. Sand shrimp, amphipods, and bay anchovies were popular food items in both surveys. Stomach contents analyses presented above are all from trawl caught fish but the results agree with those reported by Lindquist et al. (1994) and Steimle and Figley (1996) which collected fish using hook and line, spear and traps from Onslow Bay, NC and the Atlantic Ocean, NJ, respectively.

As part of an NEFSC outreach project of stakeholders as part of the SAW 62 assessment process, fishermen were asked if they had witnessed adult black sea bass preying on young of the year sea bass. They were also asked for their observations regarding predation on adult sea bass. Among 94 respondents only 6 had ever seen evidence of cannibalism. Predation on adults was observed primarily from adult bluefish, striped bass, sharks and dogfish and cormorants.

All of the stomach content analysis done on black sea bass indicates a generalist carnivore that feeds primarily on benthic crustaceans but will opportunistically feed on any prey encountered including small fish. This includes a limited amount of cannibalism. Also common in the datasets noted above is the ontogenetic shift from small crustaceans to larger decapods including juvenile crab and lobster.

### *Natural Mortality*

Instantaneous natural mortality (M) has been explored in several previous assessments and a value of 0.4 has been used in most the recent assessment. A maximum age of 12 (oldest age in

NEFSC data) was assumed in re-examining  $M$  using available published methods. A constant  $M$  was chosen for use in the assessment and the average value among 20 estimates equaled 0.4, with a range from 0.1 to 0.9 (Figure A6, Table A2).

### *Fisheries*

In the Northwest Atlantic, black sea bass support commercial and recreational fisheries. In 1939 and 1940, 46-48% of the commercial landings were in New England, primarily in Massachusetts. After 1940, the center of the fishery shifted south to New York, New Jersey and Virginia. Landings peaked in 1952 at 9,883 mt (Figure A7) with the bulk of the commercial landings from otter trawls, then declined steadily to a low point of 566 mt in 1971. Historically, trawl fisheries for sea bass focused on the over-wintering areas near the shelf edge. Inshore pot fisheries, which were primarily off New Jersey, showed a similar downward trend in landings between the peak in 1952 and the late 1960s. The large increase in landings during the 1950's appears to be the result of increased landings from otter trawlers, particularly from New York, New Jersey and Virginia. During the same period, a large increase in fish pot effort, and subsequent landings, occurred in New Jersey. In recent years, fish pots and otter trawls account for the majority of commercial landings with increasing contributions from hand-line fisheries (Figure A8). Inshore commercial fisheries are prosecuted primarily with fish pots (baited and unbaited) and hand-lines. Recreational fisheries, mostly boat based, generally occur during the period that sea bass are inshore. Once fish move offshore in the winter, they are caught in a trawl fishery targeting summer flounder, scup and *Doryteuthis* squid (Shepherd and Terceiro 1994). Hand-line and pot fisheries in the southern areas may still operate during this offshore period. A small sector of the NJ charter fleet targets sea bass offshore during the winter.

### *Stock assessment history*

Black sea bass stock assessments have been reviewed in the SARC/SAW process (SAWs 1, 9, 11, 20, 25, 27, 39, 43, Data Poor Workshop and 53) beginning with an index based assessment in 1991 (summary in Table A3). In 1995 a VPA model was approved and the results generally showed fishing mortalities exceeding 1.0 (estimated using an  $M=0.2$ ). The VPA was reviewed again in 1997 when it was considered indicative of general trends but too uncertain to determine stock status. In 1998, another review was conducted and both VPA and production models were rejected as either too uncertain or inappropriate for use with a hermaphroditic species. A



suggestion was made to use an alternative method such as a tag/recapture approach. The NEFSC survey remained the main source of information regarding relative abundance and stock status. A tagging program (Moser and Shepherd 2009) was initiated in 2002 and the first year results were presented for peer review in 2004. The review panel concluded that a simple tag recovery model using the proportion recovered in the first year at large, as well as an analysis of survey indices, produced acceptable results to determine exploitation rate and stock status. The release of tags continued through 2004 and results of tag models as well as survey indices were presented for SARC review in 2006. Their findings were that the tag model did not meet the necessary assumptions and the variability in the survey indices created uncertainty which prevented determination of stock status. The panel did not recommend any alternative reference points, but they recommended continued work on length based analytical models. The black sea bass assessment was reviewed by the Northeast Data-Poor Stocks Working Group (NDPSWG) in December 2008. The review panel considered a statistical catch-at-length model (SCALE) and a variety of natural mortality options. They concluded that the length-based model was suitable for evaluating stock status and recommended a constant natural mortality option of 0.4. Although the stock was considered not overfished or experiencing overfishing, the uncertainty in the results prompted the reviewers to recommend caution in applying the results for management.

A stock assessment using a statistical catch at age model (ASAP, Legault and Restrepo 1998) was most recently reviewed by SAW53 (December 2011) but was not accepted for use in management. The review panel concluded that spatial structure within the stock and the species strong association with complex habitats when inshore compromised the ability of a single age-structured model to adequately capture the dynamics. They also concluded that new data would be required to produce an assessment useful for management.

### *Management Overview*

The Atlantic States Marine Fisheries Commission (ASMFC or Commission) and Mid-Atlantic Fishery Management Council (MAFMC or Council) cooperate to develop fishery regulations for black sea bass from Maine through Cape Hatteras, North Carolina (for details see Appendix A4). The Council and Commission work in conjunction with the National Marine Fisheries Service (NMFS), which serves as the federal implementation and enforcement entity. The Fishery

Management Plan (FMP) defines the management unit as black sea bass (*Centropristis striata*) in US waters in the western Atlantic Ocean from Cape Hatteras, North Carolina to the US-Canadian border (MAFMC 1996).

Commercial and recreational black sea bass fisheries are managed using annual catch limits, commercial quotas, recreational harvest limits, minimum fish sizes, gear regulations, permit requirements, and other provisions as prescribed by the FMP. Based on the allocation percentages in the FMP, 49% of the total allowable landings of black sea bass is allocated to the commercial fishery as a commercial quota and 51% is allocated to the recreational fishery as a recreational harvest limit. These measures are described in more detail in subsequent sections of this document.

The Council and the Commission's Summer Flounder, Scup, and Black Sea Bass Management Board (the Board) developed a cooperative management process for black sea bass including complementary FMPs, regular joint meetings, and joint decision-making on most aspects of management. This cooperative management endeavor was developed because a significant portion of the catch is taken from both state waters (0-3 miles offshore) and federal waters (3-200 miles offshore, also known as the Exclusive Economic Zone or EEZ). Primary responsibility for management in federal waters falls to the Council, while the Commission has primary jurisdiction over the fisheries in state waters. However, many measures, including annual catch limits, are jointly established on a coastwide basis and apply in both federal and state waters.

The Commission operates under the authority of the Atlantic Coastal Fisheries Cooperative Management Act (ACFCMA) of 1993. Commission member states include all Atlantic coast states, and for black sea bass, states that have declared an interest in the fishery include Maine, New Hampshire, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Delaware, Maryland, Virginia, and North Carolina. All states that are included in a Commission fishery management plan must implement required conservation provisions of the FMP, or the Secretary of Commerce may impose a moratorium for fishing in the noncompliant state's waters.

The Council operates under the authority of the Magnuson-Stevens Fishery Conservation and Management Act (MSA), passed in 1976 and amended in 1996 and 2007. The MSA extended U.S. jurisdiction to 200 nautical miles and established eight regional fishery management councils with representation from the coastal states and fishery stakeholders. The Mid-Atlantic

Council's representation includes 21 voting members and four non-voting members. Seven of the voting members represent the constituent states' fish and wildlife agencies from the states of New York through North Carolina. Thirteen members are private citizens who are knowledgeable about recreational fishing, commercial fishing, or marine conservation. The four non-voting members represent the Atlantic States Marine Fisheries Commission, the U.S. Fish and Wildlife Service, the U.S. Department of State, and the U.S. Coast Guard.

As mandated under the MSA, the Council's Scientific and Statistical Committee (SSC) gives ongoing scientific advice for annual catch limits and other issues as needed. Other technical issues and analyses are addressed through the joint Summer Flounder, Scup, and Black Sea Bass Monitoring Committee and the Commission's Black Sea Bass Technical Committee. The Monitoring Committee is a joint committee of the Council and the Commission made up of staff representatives of the Mid-Atlantic and South Atlantic Fishery Management Councils, the Atlantic States Marine Fisheries Commission, the Greater Atlantic Regional Office, and the Northeast Fisheries Science Center. The Monitoring and Technical Committees annually review the performance of management measures and recommend any adjustments necessary to account for the current stock status, catch limits, and issues facing the commercial and recreational fisheries.

The Council and Commission each have Summer Flounder, Scup, and Black Sea Bass Advisory Panels, comprised of commercial, recreational, and environmental stakeholders with an interest in these fisheries. The Advisory Panels provide ongoing advice to the Council and Commission regarding on-the-water trends and issues in the fisheries.

In an annual process the Council and Commission review catch and landings limits and a specific set of commercial and recreational measures that can be modified annually. The Council and Board meet jointly each year to consider the annual catch limits for the upcoming fishing year or years, along with the commercial and recreational size limits, seasons, gear requirements, and possession limits. Multi-year specifications may be set for black sea bass for up to three years at a time. The Council and Board consider the recommendations of the SSC, the Monitoring Committee, Advisory Panel members, and public comments before recommending commercial quotas, recreational harvest limits, and other annual management measures for all three species.

More substantial modifications to the FMP or regulations are undertaken through FMP amendments, addenda, and framework actions. FMP “amendments” include major changes to the management regime that require extensive analysis, and are typically developed jointly by both the Council and the Commission. For less complicated actions, the Commission may also make changes through an “addendum” process, while the Council can make minor changes to the FMP through a “framework adjustment” process, both of which typically take less time and resources than an amendment requires.

The Council submits any recommended changes in regulations or FMP elements to the NMFS Greater Atlantic Regional Administrator to consider for implementation. The Regional Administrator reviews the recommendations in this document and may revise them, if necessary, to achieve FMP objectives and to meet statutory requirements. The Commission’s actions do not require review by NMFS, and member states are responsible for implementing the Board’s decisions.

**TOR 1.** Summarize the conclusions of the February 2016 SSC peer review regarding the potential for spatial partitioning of the black sea bass stock. (The consequences for the stock assessment will be addressed in TOR-6.)

The review panel for the SAW/SARC 53 black sea bass assessment noted “*concerns over the potential for spatial structure and incomplete mixing within the stock area that compromised the ability of the forward projecting catch at age model to index abundance and fishing mortality reliably based on the data available.*” In response to this need to introduce a spatial element into the sea bass assessment, the WG developed a recommendation to partition the stock north of Cape Hatteras, NC into two sub-units divided by the Hudson Canyon for development of the assessment model. The basis for this recommendation was presented to a four member ad-hoc review panel of the Mid-Atlantic Fishery Management Council Science and Statistical Committee (MAFMC SSC) on February 16, 2016. The panel presented its report on March 3, 2016 to the full SSC committee and concluded that the recommended separation was reasonable and appropriate to use as a starting point for developing a black sea bass stock assessment model. The full report to the ad-hoc review panel and their response to the SSC are provided in Appendix A5

The implication of a split area assessment was evaluated using a simulated black sea bass population. Empirical sea bass data was used as the basis to develop a migratory sea bass population within a Stock Synthesis (SS) model framework. The pseudo-datasets were used within an SS model framework to compare results with area partitioning, without area partitioning and with both movement and area partitioning. The details of the simulations are provided in Appendix A6. The conclusion of the simulated assessment was that the two area model was more effective in capturing the dynamics of the overall population than a one-area assessment, and more complex models that account for movement among sub-units did not perform significantly better than the two-area estimation model approach.

**TOR 3.** Present the survey data being used in the assessment (e.g., indices of abundance, recruitment, state surveys, age-length data, etc.). Investigate the utility of commercial or recreational LPUE as a measure of relative abundance. Characterize the uncertainty and any bias in these sources of data.

Survey data considered for the assessment included NMFS winter and spring surveys and state survey data from MA, RI, CT, NY, NJ, DE, MD, VA and NEAMAP (Northeast Area Monitoring and Assessment Program) (Table A4) (see also Maps in Appendix A7 ). A recently published study (see TOR 4) examined the influence of winter oceanographic conditions on black sea bass abundance. The study concluded that overwintering conditions are a critical determinant of juvenile survival and abundance at age 1. Autumn survey indices are generally dominated by age 0 fish but may not be indicative of subsequent cohort strength. Consequently, the WG made a decision to limit survey abundance indices to winter or spring surveys when possible.

#### *State Surveys*

The Virginia Institute of Marine Science (VIMS) conducts a monthly trawl survey targeting juvenile fish within Virginia tributaries of the Chesapeake Bay and provided a random stratified index of black sea bass abundance (Figure A9). The index is for black sea bass sampled in May, June, and July since 1989 and contains fish that are less than 110, 150, and 175 mm total length, respectively. All are age-1 fish, assuming a Jan 1 birthdate. Thus, the mean number per tow index for 2015 represents the 2014 year class (spawned in 2014). The results shows variable recruitment trends with above average year classes in 1989 (the largest in the time series at 2.36

fish per tow), 2001, 2007 and 2010. The 2010 index (1.11 fish/tow) was above the series average of 0.67 fish/tow. Recruitment since 2013 has been below the series average (0.674 fish per tow) with a 2015 index of 0.290 fish per tow. The VA index was applied to the southern area.

The Maryland Dept. of Natural Resources conducts surveys from April through October in coastal bays using a 16ft trawl. Twenty sites have been sampled monthly since 1989. Black sea bass collected in the survey are all less than 21 cm and age 1 or less. The index (log mean) has shown a downward trend since a 2008 index of (1.42 fish per tow) (Figure A10). The most recent index in 2015 equaled 0.27 fish per tow, below the series average of 0.64 fish per tow. The MD index was applied to the southern area.

The Delaware Division of Fish and Wildlife operates a monthly (April to June) fixed station trawl survey in the Delaware portion of the Delaware Bay using a 16' trawl. The index serves as a spring index of age one abundance for the southern area (Figure A11) with a time series average of 0.26 fish per tow. The peak index occurred in 2001 (2000 cohort) at 1.82 fish per tow. There was evidence of the 2011 cohort in 2012 which had an above average index of 0.565 fish per tow. The 2015 index of 0.22 was below the time series average index.

The New Jersey Department of Environmental Protection conducts a depth and area stratified random trawl survey in state waters during January, April, June, August, and October. Data from June cruises in 1989 to 2016 were used as an index of abundance (Figure A12). The index was developed using a negative binomial GLM. Available covariates included tow depth, bottom temperature (BTmp), bottom salinity (BotS), and bottom DO, but only covariates that explained at least 5% of total deviance were retained. The final model was specified as  $Catch \sim Year + Depth + BotS$ . The resulting index was decomposed by annual length frequencies, and indices at age were developed using the southern region ALK. The composite index peaked in 2009 at 13.73 fish per tow. The 2015 total index (4.52 fish/tow) was above the series average (3.29 fish/tow). A 2016 spring index was 1.71 fish per tow, well below the series average. The NJ index was applied to the southern area.

New York Department of Environmental Conservation operates a small mesh trawl survey which targets age 0 and juvenile finfish species in the Peconic Estuary since 1987. Tows are conducted weekly from May through October at 16 randomly chosen stations out of a possible 77. Tows from eastern survey stations that occurred in May, June and July from 1990 to 2015 were used to

generate an age 1 spring index. Black sea bass  $\leq 15$ cm TL in the spring were considered Age 1 (Figure A13). The index was developed using a 0-inflated negative binomial GLM with depth and salinity as significant environmental covariates that explained a combined 7% of total deviance. Bottom temperature was not a useful covariate. While the time series average is 0.11 fish/tow, significant peaks occurred in 2002 (0.34 fish/tow), 2012 (1.11 fish/tow) and 2016 (0.58 fish/tow).

The Connecticut Department of Energy and Environmental Protection Long Island Sound Trawl Survey (LISTS) is conducted from longitude 72° 03' (New London, Connecticut) to longitude 73° 39' (Greenwich, Connecticut). The sampling area includes Connecticut and New York waters from 5 to 46 m in depth and is conducted over mud, sand and transitional (mud/sand) sediment types. Prior to each tow, temperature (°C) and salinity (ppt) are measured at 1 m below the surface and 0.5 m above the bottom using a YSI model 30 S-C-T meter. Water is collected at depth with a five-liter Niskin bottle, and temperature and salinity are measured within the bottle immediately upon retrieval.

Sampling is divided into spring (April-June) and fall (Sept-Oct) periods, with 40 stratified-random sites sampled monthly for a total of 200 sites annually. For this assessment, only the spring leg of the survey was used which is believed to represent age 1 and older black sea bass. The sampling gear employed is a 14 m otter trawl with a 51 mm cod-end. The sampling area is divided into 1.85 x 3.7 km (1 x 2 nautical miles) sites, with each site assigned to one of 12 strata defined by depth interval (0 - 9.0 m, 9.1 - 18.2 m, 18.3 - 27.3 m or, 27.4+ m) and bottom type (mud, sand, or transitional as defined by Reid et al. 1979). The number of sites sampled in each stratum was determined by dividing the total stratum area by 68 km<sup>2</sup> (20 square nautical miles), with a minimum of two sites sampled per stratum. Discrete stratum areas smaller than a sample site are not sampled. The survey's otter trawl is towed from the 15.2 m aluminum R/V John Dempsey for 30 minutes at approximately 3.5 knots, depending on the tide. Black sea bass at each station are counted and total lengths measured to the nearest centimeter.

This survey was not designed to target black sea bass. In order to generate a black sea bass index of abundance, a statistical model-based standardization of the survey data was conducted to account for factors that affect black sea bass catchability. Following the approach described in the ASMFC's standardization guidelines (SEDAR 2015), an index of age 1 and older black sea

bass was created using a negative binomial generalized linear model (GLM) with a log link and bootstrapped estimates of uncertainty. Zero inflated negative binomial models were also tested and compared using a Vuong non-nested hypothesis test (Vuong 1996).

A full model that predicted catch as a linear function of year (categorical data), month (categorical data), depth (continuous data), and stratum (categorical data) was compared with nested sub-models using AIC. Temperature and salinity were not able to be used for this analysis because the recording of those covariates began late in the time series. The model that included year and month was selected because it produced the lowest AIC coupled with the lowest variance inflation. The index exhibited a marked increase from low catches early in the time-series with the increase beginning in the early 2000s to present with inter-annual variability. Diagnostics identified slight over-prediction of average annual catch per tow at the end of the time series.

The indices show low abundance until an increasing trend beginning in 2012 (Figure A14). Large age 1 indices occurred in 2002, 2012 and 2014 with the recent cohorts contributing to the sharp increase in abundance. Prior to 2012 the average index was 0.5 fish per tow compared to the 2014 and 2015 indices of 14.3 (CV=0.21) and 14.2 (CV=0.21) fish per tow, respectively.

The Rhode Island Department of Environmental Management research trawl survey is conducted with a  $\frac{3}{4}$  high-rise heavy-duty bottom trawl towed for 20 minutes at 2.5 knots. Sampled areas include Narragansett Bay and Rhode and Block Island Sounds. Data include a mixture of fixed and random sampling stations, depending on the component of the survey being examined (seasonal = random stratified, monthly = fixed station). Data collection has been generally consistent from 1979 to the present, with the exception of tow numbers early in the time series, and a change to the survey doors in 2012. Investigations between survey doors during a calibration experiment indicated that there are not major differences in catchability of black sea bass between the two door designs. Data elements collected include numbers, weights, and lengths caught by species as well as a suite of environmental information including bottom and sea surface water temperature, depth, and sea conditions.

The survey has two seasonal components, a spring (April/May) and a fall (September/October) survey that has been conducted annually since 1979. Sampling is conducted during daylight hours only. For assessment purposes, only the spring survey is used, which involves



approximately 42 tows per year. The spring seasonal survey employs a stratified-random sampling design. The sampling area is divided into strata defined by depth. For each seasonal sampling cruise, sites are selected randomly from within each stratum.

All black sea bass collected are weighed (kg) and measured for total length (cm). The number of individuals measured from each tow varies by species, and also depends on the size of the catch and range of lengths. If a species is subsampled, the length frequency of the catch is determined by multiplying the proportion of measured individuals in each centimeter interval by the total number of individuals caught.

In order to generate a black sea bass index of abundance for the stock assessment, statistical, model-based standardization of the survey data was conducted to account for factors that affect black sea bass catchability. Only ages 1 and above are represented in the spring component of the RI seasonal trawl survey. Following the approach described in the ASMFC's standardization guidelines, an index of age 1 and above black sea bass was created using a negative binomial generalized linear model with a log link and bootstrapped estimates of uncertainty. Proportion of positive tows for black sea bass averaged approximately 11% across the time series, with that average increasing to 19% since 2000.

A full model that predicted catch as a linear function of year (categorical data), bottom temperature (continuous data), month (categorical data), station (categorical data), and depth (continuous data) was compared with nested sub-models using AIC. The model that included year, bottom temperature, and depth was selected because it produced the lowest AIC coupled with the lowest variance inflation. The index exhibited a marked increase from low catches early in the time series with the increase beginning in the mid-1990s to present. There is some degree of small decline at the end of the time series. Diagnostics identified slight under-prediction of average annual catch per tow

The indices have been highly variable over time, although the spring index includes several above-average years since 1999 (Figure A15). The 2015 overall index (1.025 fish/tow) was well above the series average (0.180 fish/tow, std error = 0.073). The Department also conducts a coastal pond seine survey and although the mean catches per tow are small, it does show an increasing trend, peaking in 2014 at 2.04 fish per tow.

The Massachusetts Division of Marine Fisheries has conducted trawl surveys each spring and fall since 1978. The spring survey occurs in May and the fall survey occurs in September. The survey employs a stratified random design, with stratification based on five bio-geographic regions and six depth zones. The six depth zones include 0-9.1 m, 9.1-18.2 m, 18.2-27.4 m, 27.4-36.6 m, 36.6-54.9 m, and >54.9m. Surveys from 1978-1981 were conducted aboard the F/V Frances Elizabeth. Surveys since 1982 have employed the R/V Gloria Michelle. Trawl design and trawl doors have been consistent for the duration of the survey. The net is a ¾ size North Atlantic type two-seam otter trawl (11.9 m headrope/15.5 m footrope) rigged with an 8.9 cm rubber disc sweep and a 0.6 cm knotless codend liner (1.3 cm stretched mesh). The trawl is spread behind 1.8 m x 1.0 m 147 kg wooden trawl doors, 19 m 1.0 cm chain bottom legs and 18.3 m 1.0 cm wire top legs.

Black sea bass are encountered in both the spring and the fall survey, and are captured in all regions. All sizes of BSB are captured by the survey. Black sea bass are most common in Regions 1-3, which include waters from the Massachusetts-Rhode Island border, the waters of Nantucket and Vineyard Sound, and the eastern shore of Cape Cod.

Only the spring survey was standardized for the assessment. The survey was standardized using negative binomial regression. No offset was included because tow time was standardized throughout the entire time series. Potential predictors included temperature and stratum. Models were evaluated based on AIC. Results indicated that the model with the lowest AIC was one that included an effect of temperature and stratum. Diagnostic criteria, such as qq and residual plots, indicated acceptable model fit.

The MADMF spring index declined during the 1990s, rose briefly in 2000, then again in 2008 (Figure A16). The spring 2012 mean number per tow (3.882 fish/tow and CV=0.309) was the beginning of a period of indices above the series average (1.88 fish/tow). The index spiked in 2014 (10.458 fish/tow with CV=0.29) due to the incoming 2011 year class. The index has remained above average through 2016 (4.301 fish/tow with CV=0.30).

The Northeast Area Monitoring and Assessment Program (NEAMAP) is a stratified random trawl survey conducted in coastal waters between Rhode Island and Virginia. The spring series began in 2008 when the NEFSC dropped sampling of those strata. The overall ln re-transformed mean number per tow increased slowly since 2010 (0.19 fish per tow) to a 2015 mean of 0.30

fish per tow. The spring time series was further divided into two separate north and south series. The south series included strata off the coast of NJ and south, while the north was NY and RI coastal waters. The average catch per tow in the north (1.67 fish per tow with a CV =0.64) (Figure A17) was significantly larger than the south (0.14 fish per tow with CV=0.22) (Figure A18). The northern series increased since 2008, peaking in 2014 at 3.47 fish per tow. The 2015 mean number per tow decreased to 1.93 fish per tow. The southern index had an initial downward trend but has remained generally stable with the 2015 index approximately equal to the time series average. Catches in the southern strata were small and the final assessment model used only the age 1 indices for the southern area.

#### *NMFS surveys*

The NEFSC winter bottom trawl survey was conducted between Virginia and Georges Bank with stratified random tows from 1992 until 2007. Survey strata used for an index of abundance were offshore strata in areas comparable to the location of the winter trawl fishery. The trawl gear was a flatfish net with a cookie sweep. The overall series peaked in 2003 at 17.70 fish per tow, which followed from the 2002 index of 10.84 fish per tow. In 2007 when the survey was terminated, the survey index equaled 4.36 fish per tow.

For the assessment area models, the survey area was split into north and southern strata groups relative to Hudson Canyon. The northern stratified mean number per tow reached a peak in 2003 of 4.64 fish/tow before declining to average values by 2007 of 1.18 fish per tow (Figures A19). The comparable survey indices for the southern area were considerably larger, peaking at 31.32 fish per tow in 2003 (Figure A20). The index in the final year of the survey was 7.67 fish per tow.

Indices were converted to swept area abundance estimates for use in the assessment model. The area swept per tow between the wings during the winter survey equaled 0.0131 NM<sup>2</sup>. The winter survey indices were not included in the final northern model.

The NEFSC spring bottom trawl survey has been conducted between Nova Scotia and North Carolina since 1968. The indices (stratified mean number per tow) for black sea bass were developed using offshore strata containing at least one positive tow in the time series. Previous assessments using the NMFS data considered a log transformation of catch per tow to reduce the influence of high catches. The survey is designed to account for variation and the

transformation can violate the underlying assumption of the designed survey (T. Miller, NEFSC, pers. comm.), therefore indices were calculated as the arithmetic mean number or mean weight per tow. Sampling was conducted aboard the *FRV Albatross IV* using a Yankee 35 haddock net with a roller sweep. In 2009 the FSV *Henry B. Bigelow* replaced the *FRV Albatross IV* in conducting the survey. The *Bigelow* gear consists of 4-seam, 3 bridle net with a rock-hopper sweep and is towed for 20 minutes rather than 30 minutes with *Albatross*. Although calibration factors for the two vessels are available (calibration factor=3.41), the *Bigelow* time series was considered as a separate index series of black sea bass abundance.

The NEFSC spring mean number per tow in the *Albatross* series followed a pattern of an increasing index during the late 1970s, followed by a decline during the 1980s and 1990s (Figure A21). A second increase in the index occurred beginning in 1998, peaking in 2003 at 9.51 fish per tow, followed by a decline to 0.61 by 2008. The uncalibrated *Bigelow* series began in 2009 with an index of 2.44 fish per tow and peaked in 2013 with 11.42 fish per tow. The 2015 and 2016 indices were 3.12 and 4.97 fish per tow, respectively.

The area split for the NEFSC spring series highlights a general northward spatial distribution shift. The *Albatross* series north of Hudson Canyon peaked in 1986 with 2.30 fish per tow, however the time series average was only 0.21 fish per tow (Figure A22). The numbers per tow increased significantly during the years of the *Bigelow* series, peaking at 11.02 fish per tow in 2014. The 2015 and 2016 indices were 1.88 and 8.20 fish per tow, respectively.

The indices in the southern area reflected the multiple peaks seen in the overall index (Figure A23). The *Albatross* index series peaked in 1977 at 18.25 fish per tow and again in 2003 at 20.60 fish per tow. The *Bigelow* series since 2009 had an index of 22.56 in 2013 but declined to 4.64 and 1.01 fish per tow in 2015 and 2016, respectively.

For the purposes of the area exchange model, the north and south indices were converted to swept area abundance estimates. The area swept per tow between the wings during the spring *Albatross* series equaled 0.0112 NM<sup>2</sup> while the *Bigelow* area swept per tow was 0.007 NM<sup>2</sup>

### *Recreational CPUE*

In addition to fishery independent survey indices of abundance, the Working Group also developed a fishery dependent index of abundance from the Marine Recreational Fisheries

Statistics Survey (MRFSS<sup>1</sup>). For this exercise, catch was estimated using the raw (unexpanded) Type 2 and Type 3 MRFSS data files. Estimates of potential effort for a given species can be difficult using the MRFSS data due to the self-reported nature of the data. In an attempt to account for potential effort (rather than just positive trips or self-reported directed trips), effort estimates were based on effort for a species guild. Species to include in a guild were identified using the Jaccard index of similarity (Jaccard 1912). Estimation of the similarity coefficient can be summarized as follows.

- 1) Determine the number of trips (MRFSS intercepts) that caught the target species
- 2) Determine which non-target species were caught on trips when the target species was caught.
- 3) Determine the number of trips (MRFSS intercepts) that caught a given non-target species.
- 4) Divide the number of trips that caught both the target and non-target species by the number of trips that caught either the target species or the non-target species.

Mathematically, this can be expressed as

$$J = \frac{N_{11}}{N_{10} + N_{01} + N_{11}}$$

where N is the number of trips and the subscripts of 0 and 1 are binary for observation of the target and non-target species. High values of  $J$  suggest high correlation between the target and non-target species (*e.g.* habitat utilization), so observation of the non-target species implies presence of the target species even if it is not observed.

For the current analysis, species associations were evaluated at the regional level using all years combined. Guilds were composed of the target species and any species with a similarity coefficient greater than 5% ( $J \geq 0.05$ ). Any trip that caught any one of the guild species was considered a potential black sea bass trip. Regional species guilds and time series of effort (# of intercepted trips with guild species) are shown in Tables A5-A6, respectively.

For each potential black sea bass trip identified through the guild analysis, trip level CPUE was estimated as the black sea bass catch divided by the number of anglers contributing to the catch (catch per angler or CPA). Because observed (Type A) and unobserved (Type B1 and B2) catch

---

<sup>1</sup> Although the MRFSS was officially replaced by the MRIP in 2012, MRFSS-based raw data files are available through 2015, allowing a continuous time series of MRFSS data for this analysis.

are handled separately by MRFSS and do not necessarily have the same number of anglers associated with the two types of catch on a given trip, it was necessary to develop separate CPA estimates for observed and unobserved fish and sum them ( $CPA = CPA\_A + CPA\_B$ ).

Admittedly, this is not ideal, but should not have an overall large effect on the results.

Trip specific CPA was then modeled using R software with a delta-lognormal GLM. Full models included covariates of year, area, wave, state, mode, and hours fished. Final models included only covariates that explained at least 5% of total deviance (Table A7).

Effort in the northern region generally increased during the 1980s, rising from less than 1000 intercepted trips in 1981 to over 4000 intercepts by 1990 (Table A6). Effort subsequently leveled off, and varied without trend between 2000 and 4000 intercepts annually for the years 1990 to 2010 before showing a general increase in recent years. CPA in the northern region remained generally stable below 0.25 fish per trip between 1989 and 1998 (Figure A24). Catch rates increased to around 0.5 for a brief period, before dropping back to previous levels in 2004 and 2005. Over the last decade, recreational catch rates of black sea bass in the northern region have shown a dramatic increase, rising from 0.23 fish per trip in 2005 to 1.7 fish per trip in 2015.

From the early 1980s to early 2000s recreational black sea bass effort in the southern region increased more than two-fold, rising from around 3000 intercepted trips per year to a peak of over 9000 intercepts in 2001 (Table A6). Since that time, effort has gradually declined, dropping to approximately 6400 intercepts in 2015. CPA in the southern region follows a similar pattern as the associated effort. Catch rates increased from around 1.0 fish per trip in early years to over 3.0 fish per trip by the early 2000s (Figure A25). CPA subsequently dropped by approximately 35% by 2004, and has varied without trend around 2.0 fish per trip since that time. Recreational black sea bass CPA in the southern region was estimated at 1.74 fish per trip in 2015.

The overall stock-wide CPA peaked in the late 1990s-early 2000s, which followed the southern CPA pattern (Figure A26). Since 2005 there has been a steady increase in catch rate with a sharp spike in 2012.

**TOR 2.** Estimate catch from all sources including landings and discards. Characterize the uncertainty in these sources of data. Evaluate available information on discard mortality and, if

appropriate, update mortality rates applied to discard components of the catch. Describe the spatial and temporal distribution of fishing effort.

### *Spatial and Temporal Distribution*

Spatial patterns in the commercial fisheries were examined and distribution maps presented in Appendix A7. The trawl fishery north of Hudson Canyon generally catches black sea bass across the shelf throughout the year. There is a trawl fishery within the New York Bight that targets scup and summer flounder as well as sea bass. As sea bass migrate out to the shelf edge with declining water temperatures, the trawl fleet follows and operates south of Hudson Canyon along the shelf edge. Pot fisheries generally operate further inshore than the trawl fleet as does the hand-line fleet. However, there is a hand-line fishery in the southern end of the stock that operates throughout the winter closer to the shelf edge.

The recreational fishery occurs from late April into October/November while black sea bass are distributed in coastal waters. There are party and charter boats that occasionally target black sea bass in January and February near the Hudson Canyon. The distribution of trips is presented across years due to confidentiality with the limited number of vessels involved. The wave 1 (January- February) fishery has been closed in recent years.

A tagging study in the mid-2000s demonstrated the migratory pathways of black sea bass originating in different areas (Moser and Shepherd 2009). Fish originating north of Hudson Canyon follow the shelf break south and are captured in the winter trawl fishery. Fish originating to the south of Hudson Canyon remain in the south. The pot and hand-line fisheries operate inshore (with the exception of the southern hand-line fishery which operates offshore adjacent to the point of origin). Consequently, mixing among the two defined spatial units is most likely to occur south of Hudson Canyon with fish from both areas present in the southern trawl catch. In order to account for the mixing, the fisheries were divided into trawl and non-trawl components with the winter trawl fishery acting as a proxy for the mixed component of the stock.

### *Commercial fishery*

The commercial fishery on the northern black sea bass stock (Maine to Cape Hatteras, NC) is prosecuted primarily with fish pots, otter trawls and hand-lines (Figure A8). Fish pots and hand-

lines are generally fished in inshore waters when targeting black sea bass (with the exception of some lobster and sea bass pots in NY). Trawls are generally offshore in the winter months in conjunction with summer flounder and scup fisheries (Shepherd and Terceiro 1994). Fish pots have accounted for 44% of landings since 1989, followed by otter trawls at 41% and hand-lines at 9%. Other gears account for 7%. The majority of the landings occur in January through June, peaking in May.

Trends in landings were relatively stable at around 1,300 mt until 2007 (Table A8, Figures A7, Figure A27). State and Federal management plans were implemented in 1998 which included minimum size restrictions and commercial quotas. In 2008, additional quota regulations were enacted which decreased landings to an average of 739 mt between 2008 and 2012. Since 2013 commercial landings have increased to 1,113 mt in 2015. The commercial sea bass fishery is prosecuted in all states between Massachusetts and North Carolina however Massachusetts, New Jersey, Maryland and Virginia account for over 70% of total commercial landings (Table A8). The majority of landings since 1989 (54.6%) are from areas 611-622 with most trawl landings coming from areas 616 (17%), 622 (26%) and 626 (17%). Total landings by NMFS statistical areas are presented in Table A9. Black sea bass landings have been dominated by medium, large and jumbo sizes since 2000 (Figure A28, Tables A10-A17). Beginning in 2011, large and jumbos have accounted for 60% or greater of the landed fish. In 2015 sea bass from medium, large and jumbos market categories represented 23%, 37% and 29% of overall landings, respectively.

The time series of fishery data in the assessment was limited to 1989 to 2015. The NEFSC fishery observer program began in 1989. Therefore estimates of discards prior to 1989 would have to be determined via ratio estimates to other fisheries. The consensus of the WG was to limit the data series to years with empirical estimates of commercial discards. Commercial data was divided into half year time blocks for gear categories of trawl and non-trawl. The north-south criteria adopted in TOR1 was applied in the expansion of the catch at length and catch-at-age. Spatial coverage of length sampling was insufficient to allow region specific samples to be applied to the landings by market category. The WG concluded that regional variation in landing by market category would adequately capture any size differences between north and south and length distribution within market category was narrow enough to minimize any regional effect.



Average size by market category was also compared across years (Figure A29) and areas (Figure A30) to evaluate the inter-annual stability of size composition within market grades. Length distributions within market categories were relatively stable. Therefore size composition within a market category was applied between adjacent years when needed, i.e. a medium was considered a medium regardless of time or space. In summary, length distributions within market categories (small, medium, large, jumbo and unclassified) by half year period (Jan-Jun, Jul-Dec) were applied to landings by half year, region (north and south), market category and gear category (trawl vs non-trawl) for 1989-2015, using half-year region specific length-weight equations.

Length measurements (cm) of sea bass in the commercial landings are sampled by NMFS in ports from Maine to North Carolina. Samples are collected from boxes of fish available from dealers and sorted by market category. Market categories are extra small, small, medium, large, jumbo and unclassified. Length frequencies by market category and half year were expanded to total catch beginning with 1989. NMFS samples were supplemented with similar information collected by the state of North Carolina between 1989 and 1998. The NC lengths measurements were combined with NMFS data by market category and half year. Landings from the extra small category since 1989 were minimal and the size distributions were comparable to smalls. Therefore extra smalls were combined with smalls. Sample sizes and total number of fish measured from NMFS and NC data are provided in Tables A18. Sampling intensity is expressed as metric tons (mt) of landings per 100 fish measured (Tables A19-A20). Since 1989 the overall commercial sampling has averaged 21 mt per 100 lengths, with the best sampling years between 2008 and 2012 when sampling was on the order of 7 mt per 100 lengths (Figure A31).

Length weight information to convert landed weight to number was available from NMFS spring and autumn survey data since 1992. The equations applied to all length samples by season and region were:

$$\begin{aligned} \text{North Spring:} & \quad 1.0157e-5 * \text{length(cm)}^{3.0769} \\ \text{North Autumn:} & \quad 1.2649e-5 * \text{length(cm)}^{3.0173} \\ \text{South Spring:} & \quad 1.1448e-5 * \text{length(cm)}^{3.0435} \\ \text{South Autumn:} & \quad 1.9053e-5 * \text{length(cm)}^{2.8937} \end{aligned}$$

In the expansion process, missing cells (Figure A32) were replaced with lengths from the same market category and the closest year or years containing measurements. The most frequent substitutions occur in the second half of the year within the unclassified category between 1993

and 2002 (Figure A33). In addition, jumbos in the second half year prior to 2002 were also poorly sampled. However, the lack of samples also reflects limited availability due to minimal landings. Since 2003 the only missing cells were unclassified in the second half of 2010 and 2011, which represented 2% and 3%, respectively, of total annual catch. Cells with substitutions are presented as shaded cells within the sampling intensity table.

The total number of black sea bass landed across both areas combined has declined since 1996 (5.0 million) to a low of 935,000 in 2009 (Figure A34). Total landings since 2010 (1.27 million) have increased steadily to 1.49 million in 2015. Overall mean length in the landings in both areas have increased steadily from around 28 cm in 1989 to 34 cm in 2004 and have remained relatively stable since 2003 around 35 cm (95% CI  $\pm$ 12 cm). The same size trends exist within area, season and gear types (trawl vs non-trawl) (Figures A35-A37). The mean size of landings does not vary much by season, however by gear type there is some area differences with the non-trawl landings smaller in the south.

#### *Commercial discards*

Estimated discards were calculated from four gear types. Otter trawl and sink gillnet discards were calculated using the Standardized Bycatch Reporting Methodology (SBRM) (Wigley et al 2008). SBRM relies on information collected by NMFS observers on a sub-sample of commercial trips as part of a program begun in 1989. Annual discards per half-year were estimated in each region (north/south) as the ratio of recorded discards for the species in question to recorded kept of all species landed, multiplied by the total reported landings of all species in that time strata. The associated CV for the estimate was also calculated (Table A21). The observed trips for hand-line or fish pot gear was limited, therefore the SBRM approach was not used for discard estimates. Pot and hand-line discards from 1994-2015 were estimated from self-reported vessel trip logs (VTR), adjusted to total landings by gear. VTR logs were not required prior to 1994. For the period 1989-1993, the ratio of the trawl discards to pot discards and hand-line discard per half year from 1994-1996 was multiplied by the associated trawl discard estimates. Management began in 1997 so 1994-1996 was used to avoid regulatory discards influencing the results. Pot and hand-line discards were combined into the non-trawl category. Discards from trawls and sink gillnets were assumed to suffer 100% mortality because of depths fished and length of tow or soak time. Discard mortalities of 15% were applied to pot

and hand-line discards. The rationale was that depths fished generally resulted in minimal barotrauma and the volume of fish in a pot catch would result in minimal damage to released fish. Hand-line discard mortality was assumed equivalent to recreational discard mortalities. Discards from sink gillnets was minimal, with an annual average in the time series of less than 0.2 mt. The highest year for discards occurred in 1991 with 1.9 mt, however the CV for the estimate was 0.94.

The largest amount of discard mortality was attributed to the trawl fishery (Table A21). Annual discard losses ranged from 2.0 mt (CV=0.64) in the north during 1991 to 607.2mt (CV=0.62) in the south during 1996. Precision of the trawl discard estimates in the time series varied but CVs averaged 0.53 in the north and 0.42 in the south. Discards in the non-trawl gears was primarily from pots. Discard mortalities from the non-trawl gear averaged 5% of the total discards, however annual proportions ranged from 0.4% to 59.3%. The years with higher proportion of non-trawl discards occur when trawl estimates are low. Total weight of discards in non-trawl gear ranged from 0.7 mt to 23.9 mt (Table A22). In 2015 total trawl discards were 128.9 mt in the north and 186.1 mt in the south. Non-trawl discard losses totaled 17.1 mt in the north and 2.9 mt in the south.

Commercial discard length samples were divided by region (north/south), half year and gear type (trawl/non-trawl) (Table A 22a). Length samples from observed trawl trips prior to 1997 were sporadic, so length samples were pooled (pooled distribution did not include a cluster of fish in 1989 less than 9 cm that were not considered representative) (Table A22b). Length samples from non-trawl gear were limited (although observer data was not used for non-trawl discards estimation, there were limited length samples available). Substitutions for missing lengths were made from other areas, time blocks or gear depending on available alternatives. Most substitutions were made prior to 2000.

Annual commercial discard length distributions show a shift in the size composition over time (Figure A38). Prior to the FMP, discards were composed primarily of sizes below 31 cm. As minimum sizes and quotas went into effect the size distribution increased (likely due to gear changes) and included larger individuals of legal size. The primary reason for discards was quota restrictions or below minimum size (Figure A39).

#### *Recreational Landings and Discards*

The NMFS Marine Recreational Fishery Statistical Survey (MRFSS) provided catch estimates between North Carolina and Maine beginning in 1981. In 2006 following a review by the National Research Council, the design of the survey was changed and the program became the Marine Recreational Information Program (MRIP) which has continued to evolve. Calibrations have been made to adjust the MRFSS estimates to MRIP estimates from 2004 to present. However, pre-2004 estimate adjustments were left to the discretion of individual analysts. The relationship between the calibrated and uncalibrated black sea bass estimates was linear with a slope of 0.975 ( $R^2 = 0.90$ ) (Figure A40). The WG concluded that re-calibration of the MRFSS estimates was unnecessary.

Estimates were downloaded from the website (<http://www.st.nmfs.noaa.gov/recreational-fisheries/data-and-documentation/queries/index.html>) for AB1 fish (fish kept or fish filleted, released dead, disposed in some other way) and B2 fish (released alive). Annual estimates are provided for waves (two-month periods) 2 to 6 and by state. Landings and discards were partitioned into half year blocks and regions, with NY north into the northern region and NJ south into the southern region. Wave 1 (Jan/Feb) is not sampled in the Northeast however wave 1 estimates have been produced for North Carolina since 2004. Recreational catch in wave 1 is primarily party and charter vessels which are required to submit VTR logs while fishing in federal waters. The location is designated by NEFSC statistical area, so totals from VTR records were added to the perspective region based on statistical area. North Carolina catch may occur from either stock (partitioned at Cape Hatteras, NC) therefore annual MRFSS catch estimates were split north and south of Hatteras based on intercept sites. MRFSS estimates are provided as number of fish for AB1, B2 and weight (kg) of AB1 catches. Total weight of discards was derived by applying a length-weight equation to the expanded discard length frequencies.

Stock-wide recreational landings averaged 1,438 mt between 1989 and 2003 then declined to an average of 1,138 mt thereafter (Figure A41). Some of the decline could be attributed to changes in the regulations, particularly minimum size and bag limits. Starting around 1995 there has been a steady shift in the distribution of black sea bass recreational landing from predominantly southern states to the states in the northern region (NY and north) (Table A23). The majority of sea bass landings prior to 2007 were taken in the southern region, primarily New Jersey (Figure A42). Since then, the percentage has shifted north such that 86.2 percent of the landings in 2015

were from the northern region. Total landings in weight in 2015 were 1,588 mt in the north and 276 mt in the south, which is the largest disparity in the time series (Table A24).

Length frequencies of the recreational landings were sampled by MRIP personnel during dockside interviews. Lengths were expanded to total landings by region and half year. Average length of fish in the landings has increased over time likely as a function of changing regulations. Mean length (cm) from 1989 to 1997 ranged from 27 to 31 cm. Size increased between 1997 (29 cm) and 2002 (37 cm) then remained stable (Figure A43). Overall mean length of 2015 landings was 38 cm. North and south differences were not apparent until a divergence in 2000 when northern landings were generally larger fish than in the southern region (Figure A44). On average the northern fish were 4 cm larger and the difference was as much as 7 cm in 2014. It should be noted that until 2010 recreational size limits were relatively consistent. After 2010, states from NJ south remained at 12.5" while northern states were required to adopt larger minimum sizes (up to 15") in an attempt to constrain landings. Although the harvest was not necessarily limited it did produce larger average sizes from New York north.

Regulations of the recreational fishery based on size and bag limits inevitably results in discarding. The survival rate of discarded fish is impacted by multiple factors. Fish captured at depth and brought to the surface undergo pressure changes that can cause barotrauma. In a physoclistous fish like the black sea bass, the most obvious sign of barotrauma is the expanded abdomen and displaced organs by an over-inflated swim bladder. Combinations of angler experience and terminal tackle choice can lead to hook traumas, extended handling and exposure on deck. Conditions at the surface may be significantly warmer or colder than those found at the bottom. Assessments now routinely try to account for discard mortality and there is a growing body of work that addresses it, including several articles that specifically address black sea bass. SAW-43 (NEFSC 2006) assumed a 15% discard mortality rate for black sea bass caught recreationally with a hook and line. Previous assessments (SAW 27) which included recreational discards applied a 25% mortality rate. The assessment cited the Bugley and Shepherd (1991) estimate of 4.7%, generated from black sea bass caught in waters 6-12 meters deep, but noted that fisheries in New Jersey and south accounted for a significant proportion of discards and are often prosecuted in deeper waters. Bugley and Shepherd (1991) noted mildly extended abdomens in some fish but attributed release mortality primarily to anglers with limited experience and the forcible removal of hooks from fish hooked in the esophagus.

The Northeast Data Poor Stocks Working Group (2008) applied a 25% mortality rate to recreational discards but did not offer any justification for the increase from the previous assessment.

SAW-53 (NEFSC 2011) assumed a 15% recreational discard mortality rate, after evaluating the results of black sea bass specific studies (Bugley and Shepherd, 1991 and Rudershausen et al., 2007) and peer reviewed studies for other species. Rudershausen et al. (2007) found that 3.6 % of 199 black sea bass caught in depths from 19-71 meters (28.5 meter average) were unable to orient themselves and swim down. It was assumed that fish in this release condition would not survive. Black sea bass that were bleeding or non-jaw hooked were also more likely to be unable to swim down, but these findings were not statistically significant, perhaps due to low sample sizes. Gastric distention (barotrauma) was visibly evident in 61% of black sea bass but this condition was not associated with immediate discard mortality (poor release condition). The authors assumed that gastric distension resulted in 100% delayed mortality and estimated that 66% of black sea bass would suffer delayed mortality. Later work by these same authors would test and refute this assumption.

SEDAR 25 (2011) applied a 7% mortality rate to discards from commercial and recreational hook and line black sea bass fisheries whereas prior assessments had used 15%. This decision was based heavily upon work detailed in Rudershausen et al. (2010). Fish were caught in waters from 29-37 meters depth, tagged, release condition noted, and then after 72 hours, recaptured. Recapture rates of fish with visible barotrauma, hook trauma, those unable to swim, and those presumed dead were compared with the recapture rates of fish that were able to swim down without any visible signs of trauma (best release condition). It was assumed that fish in the best release condition survived as well as fish never caught. Six additional fishing trips, independent of tag/recapture trips, were then undertaken and a discard mortality rate for hook and line fisheries was estimated based upon prior return rates and the release condition of fish caught. Discard mortality averaged 4.3% and ranged from 0 to 6.9% over these 6 trips. In addition, Rudershausen et al. explored a number of other questions related to: a) predation on released fish swimming down from surface (not significant), b) non-jaw hook trauma (significantly lower survival rates than fish that were jaw hooked), c) post release floating as a proxy for immediate mortality (an over-estimate), d) tagging acting as inadvertent “venting” (tagging resulted in fewer floaters but did impact survival), e) the trend between discard mortality and depth was

positive, but not significant. Over the depths of this study, fish with visible barotrauma that were able to orient and swim down did not have different recapture rates (and estimated survival rates) than fish in the best release condition.

Collins et al. (1999) experimentally tested whether fish able to descend after capture with the assistance of venting had a greater chance of survival. Fish unable to orient and swim down were assumed to not survive and all other fish were held at capture depth in traps. Mortality over 24 hours in the control group (unvented) averaged 13% for black sea bass caught in waters from 20-35 meters deep (n=83). Mortality increased significantly in fish caught between 43-55 meters (39%, n=25). In addition, venting was shown to be an effective way to reduce immediate discard mortality in black sea bass caught across these depths.

Stephen and Harris (2010) collected fishery dependent data on fish release condition during commercial hook and line trips fishing from 20-80 meters depth. They estimated that 66% of released black sea bass (61/92) suffered immediate mortality due to impaired ability to swim down. Their field work occurred at greater depths (majority >40m) than previous studies (e.g. Bugley and Shepherd, 1991; Collins et al., 1999; and Rudershausen et al., 2010) and these depths are likely greater than where the majority of recreational angler activities take place. It is likely that barotraumas are contributing to the greater immediate release mortality, as was also found in Collins et al. (1999). Immediate release mortality in black sea bass did not differ with respect to size.

The data collected and reported on in Rudershausen et al. (2010) was added to, reanalyzed, and later published by Rudershausen et al. (2014), which differs from the 2010 grant report in the addition of a tagging and holding at depth component that acts as a control (no hook, pressure, deck or predation related traumas) to test the assumption that best release condition fish had 100% survival. A total of 5,131 black sea bass were tagged, with an overall tag recapture rate of 23.5%. The model estimated fish in best (surface) release condition to have a survival rate of 87%. Therefore, the fishery related median discard mortality rate (based upon the release condition of fish caught on fishing trips independent of the tag/recapture efforts) over depths from 20-35 meters was estimated to be 19%. The authors noted that in fisheries prosecuted at shallower depths (such as 6-12 meters in Bugley and Shepherd (1991)) the survival rate of surface released fish should be greater.

The five black sea bass specific studies reviewed here include a wide range (5 – 66%) of mortality estimates that could be applied to discards from the recreational hook and line fishery. Important conclusions include: a) visible barotrauma  $\neq$  100% mortality, b) best release condition  $\neq$  100% survival, c) non-jaw hook trauma (and angler experience) reduces survival to a greater extent than visible barotrauma, and d) barotrauma related mortality likely increases at depths greater than 40m.

A large proportion of the recreational fishery for black sea bass in the Mid-Atlantic occurs in depths shallower than those covered in many of these studies. Most of the studies, with the exception of Bugley and Shepherd (1991), were conducted in the South Atlantic where productive fishing grounds are generally further offshore and in deeper water. In addition, water temperatures in the South Atlantic are not reflective of most areas fished in the northern (Mid-Atlantic) stock. In the Mid-Atlantic, 75% of the recreational harvest (in pounds) and 82% of its discards (in numbers of fish) over the last 5 years (2011-2015) have been taken from within state waters (inland waters and  $\leq$  3 miles from the coast).

The hook and line discard mortality rate of 19% estimated by Rudershausen et al. (2014) is well supported by a large sample size, high proportion of recaptures, and an experimental control but the study was conducted in relatively deep water and surface released fish likely exposed to warmer water temperatures. The design of Bugley and Shepherd (1991) also had an experimental control, was conducted in shallower depths in New England, and estimated discard mortality in black sea bass around 5%. The working group concluded that the 15% discard mortality rate used in SAW-43 and SAW-53 is appropriate, slightly decreasing the discard mortality rate estimated in Rudershausen et al. (2014) due to the shallower depths at which most of the Mid-Atlantic recreational black sea bass fishery occurs. A discard mortality rate of 15% is well within the range (2.5 – 20%, avg 11%) of estimates used in recent (2009-2015) assessments for other physoclistous species from the Atlantic and Gulf of Mexico. A review of discard mortality studies related to black sea bass are provided in Table A25.

Overall recreational black sea bass discard losses ( $B2 * 0.15$ ) peaked between 2000 and 2003 with the maximum occurring in 2002 at 771 mt, well above the time series average of 263 mt (Table A24). The 771 mt represented a total number of 1.7 million fish, however the maximum number discarded in the series occurred in 2000 with 1.98 million fish. Discard losses in 2015 were 1.08 million fish equaling 317 mt (Figure A45). Discard trends in the north and south



differed over time (Table A24, A26 and Figure A46). Until 2012 the majority of discards occurred in the south but that trend reversed in 2012 (58% north) likely the result of a large incoming year class and diverging minimum sizes and season lengths between areas. The trends continued to diverge through 2015 with 79% of the discards occurring in the northern region. Discard lengths were compiled from several sources. The majority of the recreational fishery occurs from July to October, so the discard data was applied on an annual rather than seasonal basis. The American Littoral Society is a conservation group that promotes fish tagging of recreationally caught fish to follow their movement. Tagged fish are by definition B2s (caught and released alive). The lengths of the fish tagged between 1989 and 2010 were available, but measured in inches which were converted to length in cm. Additional information came from a Volunteer Angler Survey conducted by NJDEP from 2008 to 2015 involving hook and line gear. Released fish below the minimum size were classified as discards. New York DEP provided discard length information collected from party/charter boats between 1995 and 1999 as well as 2011 through 2015. Finally, the MRFSS program began at-sea sampling of party/charter boats in 2004. Mean length of discards follows a comparable pattern as landings with increasing size over time (Figure A47). Prior to 2002 southern releases tended to be larger and increasing in size over time however the trend could be a function of limited length samples. Post 2004 the regional mean lengths were comparable around 25 cm but diverge in 2011 with larger fish discarded in the north. By 2015 mean discard length in the north was 31.7 cm whereas the southern mean was 24.5 cm.

#### *Total Catch*

The pattern of total catch (commercial landings and discard losses, recreational landings and discard losses) has remained relatively stable with the exception of spikes in the early 2000s and more recently in 2014-2015 (3,631 mt and 3,683 mt, respectively) (Figure A48). The pattern by region is increasing catch in the north while the south remains stable following a decline in the mid-2000s (Figures A49-A50). The commercial catch has been less variable due in part to the quota restrictions monitored in real-time. Mean length in the catch has slowly increased in the north since the mid-1990s while mean length in the south has declined since the early 2000s (Figure A51).

#### *Catch at age development*

The primary source of age data for expansion of commercial landings and discards, recreational landings and discards and survey data was the NEFSC spring and autumn bottom trawl surveys since 1989. Commercial age samples were available since 2012 and age data collected in the NEAMAP survey provided the necessary ages for estimation of indices at age. However the available age information was incomplete for all years and in many instances was not available for all length categories. Alternative approaches were evaluated to make best use of the available ages. Sample sizes of ages by year, season, and source from 1989-2015 are given in Table A27. Ages from all years prior to 2015 are based on scales. Ages from 2015 onwards are based on otoliths.

#### *Age Length Keys*

All age length keys (ALKs) were constructed using multinomial logistic regression (Gerritsen et al. 2006; Stari et al. 2010, Weakfish Stock Assessment Subcommittee 2015) using the *multinom* function in the nnet package for R (Ripley 2016). A set of ALKs was constructed by year for spring (January-June) and fall (July-December) seasons, region (north or south) as well as by gear type (fisheries independent or commercial). Additional ALKs were constructed by year, using additional data from the previous two and subsequent two years, by season, region and gear type. Cumulative ALKs were also made for each season, region and gear type from data pooled among all years. For all ALKs, the plus group was age eight.

Age data from the northern region NEFSC fisheries-independent trawl survey in spring 2015 was insufficient to construct a suitable ALK so the 2015 age data was augmented by including data from the MADMF spring trawl survey and from NYDEC commercial market sampling.

#### *Catch at Age*

Annual catch at age (CAA) was created from recreational catch, recreational discard, commercial trawl catch, commercial trawl discard, commercial non trawl catch, and commercial non trawl discard catch at length for each region and season combination. Catch at age was calculated in two portions for each season, region and gear type combination. The first portion consisted of selecting all BSB <14 cm TL from the catch data for a specific combination. For the spring season, all BSB <14 cm TL were assumed to be age-1. For the fall season, all BSB <14cm TL were assumed to be age-0. The second portion of the CAA was constructed by selecting all BSB

for a given combination  $\geq 14$  cm TL. The ALKs were applied to this portion of the catch at length data based on criteria agreed upon by the SASC (see below).

#### *Criteria for Key Application*

Recreational CAA was created using ALKs constructed from the NEFSC trawl survey.

Commercial CAA was created using ALKs constructed from commercial sampling efforts when available, otherwise the survey ALKs were applied. Criteria for key selection were based on sample size ( $>89$ ) and a maximum age of at least 6. The minimum sample size was based on the median sample size for the fall season, southern region, fisheries independent age data 1984-2015. Median sample size for all other season, region and gear type combinations, with the exception of spring season, northern region, fisheries independent 1984-2015, was greater than 89. In cases where a year, gear, region and season combination ALK met the prescribed criteria, the ALK was applied directly to that catch data. If this criterion was not met, ALKs that included the previous two and subsequent two years of data were applied. If keys were not available that included the previous and subsequent two years, the cumulative key was applied.

Two exploratory exercises were conducted to explore the influence of data pooling and sample size on CAA (Appendix A8). Differences in predicted CAA based on ALKs for individual year vs those pooled among multiple years were examined. This exercise demonstrated that pooling resulted in differences  $<12\%$  in predicted CAA for a given catch data set. Further, a simulation was done to create a population with a known catch and a known age structure, then ages randomly sampled from the population, ALKs constructed and then CAAs calculated from the random samples. The mean age of the catch for each year in the simulation was compared to the known catch. Using 20 truly randomly sampled ages with 1,000 truly randomly sampled lengths produced a distribution of mean age of the catch that had a median close to the true mean age of the catch for all years and the distributions were relatively small e.g. they distinguished among years as true mean age of the catch changed. This work demonstrated that the ALKs allowed the calculation of accurate CAAs at sample sizes  $>20$ , suggesting that the SASC committee's criteria of 89 was appropriate.

A summary of how the age keys were applied is presented in Table A28. The final product of the length data expansion was a catch at age matrix for 1989 to 2015 for trawl and non-trawl fleets (includes recreational catch) for the north, the south and a combined north/south matrix (Tables

A29-A32). Age 0 fish represented 2% (1989 and 1990) or less and were not included in the assessment model. The overall catch at age profile indicated full recruitment around ages 3 or 4, which constituted 63% of the catch. Black sea bass age 6 and greater represented only 4% of the overall catch in the series. More detail on the age composition of the catch and indices are provided in the assessment model diagnostic plots.

Mean weights at age were determined using the total catch at age and total weight at age by area (Table A33 and A34). Comparison of annual weights at age by area show a difference by area with the southern area consistently smaller at age across all years (Figure A52)

**TOR 4.** Consider the consequences of environmental factors on the estimates of abundance or relative indices derived from surveys.

Miller et al (2015) examined the influence of winter oceanographic conditions on distribution and juvenile survival. Beginning around October, black sea bass along the coast begin to migrate offshore towards the Continental Shelf break. Fish in the northern end of the range move south to the shelf edge then follow the shelf further south depending on conditions. Fish in the Mid-Atlantic move southeast or east to the shelf edge. Age 0 sea bass, which are generally less than 14 cm when they leave coastal regions, tend to be more generally distributed across the shelf, perhaps due to slower swimming speeds. Consequently their survival is related to conditions across the shelf. If warm saline Gulf Stream water moves onto the shelf in winter, survival is high. When cold conditions are the norm, survival decreases. As a result of this variable overwinter survival as a function of oceanographic conditions, indices of young of year sea bass collected in the fall do not reflect subsequent year class strength and were not used as indices of abundance.

State survey indices were evaluated with General Linear models to determine if environmental factors influence the signal from empirical data. Model configurations varied among surveys but depth, bottom salinity and bottom temperature were common factors. New Jersey and New York found depth and salinity were significant, Rhode Island was depth and temperature, Massachusetts was temperature and Connecticut was depth (temperature and salinity not considered due to lack of data). Adjustments to the indices using these covariates altered point estimates but the same overall trends remained.

**TOR 5.** Investigate implications of hermaphroditic life history on stock assessment model. If possible, incorporate parameters to account for hermaphroditism.

Black sea bass, like other hermaphroditic fishes, are challenging to assess (Blaylock and Shepherd 2016) because of the dynamics of length and age-specific sex ratio that vary in space, time, and as a result of differential fishing and mortality. The result is that each sex will have unequal mortality patterns (Blaylock and Shepherd 2016). Standard assessment methods that model the population resilience by understanding patterns of female growth, fecundity, and mortality and temporal changes in female biomass may not be appropriate and unique modeling approaches may be warranted (Blaylock and Shepherd 2016) because the reproductive rates at levels of fishing mortality that would not be problematic for a gonochoristic species may be so for a hermaphroditic one (Alonzo et al. 2008).

Variations in reproductive strategies in fishes are diverse and include the ability to change sex from male to female (protandry) or from female to male (protogyny), simultaneous hermaphroditism and gonochoristic sex determination (McBride et al. 2013). Aside from interest in understanding the ecology and evolutionary dynamics of species, sex change as a response to sex ratio has implications for assessment and management. Black sea bass is a protogynous hermaphrodite and its stock dynamics and life-history characteristics can be impacted by size-selective fishing mortality (Provost and Jensen 2015). Such dynamics are challenging to fishery managers because the effects of size- and age-selective fishing on hermaphroditic stocks is not well understood (Provost 2013) though recent work has shown that skewed sex ratios can make a stock more vulnerable to overexploitation (Brooks et al. 2008, Blaylock and Shepherd 2016).

In a review of such challenges and the implications for management, Provost (2013) and Provost and Jensen (2015) documented the impacts to population dynamics that sex change may have and the variety of strategies that are used to model such alterations in age-specific sex ratio in quantitative stock assessments.

There are two potential primary impacts that fishing mortality has on the dynamics of a hermaphroditic stock. The first is that fishing may cause a greater skew in sex ratios and the second is that size-selective fishing mortality may decrease the size or age at sex change (Provost and Jensen 2015). A suite of behavioral strategies including sex-specific aggression and

aggregating mating strategies may reinforce the vulnerability to fishing of one sex at a greater frequency than another and serve to increase stock vulnerability. For protogynous hermaphrodite fishes, sex-specific behavioral responses may serve to change sex ratios. For example, aggressive behavior by male Scamp may have a greater fishing mortality than females because they are more likely to bite hooks (Gilmore and Jones 1992). Another behavioral aspect of size-selective mortality on hermaphroditic fishes is the impact that removing large males from harem groups may have. The interruption of social hierarchies may reduce spawning success. The effects of differential mortality on the stock are not limited to ecological impacts. The removal of males and the resulting increasingly skewed sex ratio may have evolutionary implications as well, resulting in stocks that have fundamentally different population dynamics. Simulation modeling work has indicated that populations that are subject to sex-specific mortality may be at a greater risk of overexploitation (Brooks et al. 2008).

Work to understand, in an assessment context, the impacts of differential sex-, age- and length-specific mortality dynamics (Shepherd and Idoine 1993) has indicated that alterations to the sex ratio should be explicitly included in assessment. Departures from model assumptions from hermaphroditism result in breaking one or more model assumptions in age-structured modeling approaches (Heppell and Heppell 2012). The primary issue is that in sex changing species the catch at age time series and selectivity curve may vary by sex because the fishery differentially targets older males. The implications to management for protogynous species is that fertilization rates may be lowered (sperm limitation) resulting in reduced production which has been explored in simulation (Brooks et al. 2008) and shown to be critical in determination of biological reference points.

A simulation study specific to black sea bass populations (Blaylock and Shepherd 2016) demonstrated the vulnerability to exploitation in this hermaphroditic species. Previous work (SAW 53 WP) has shown that the sex ratio in the northern stock of black sea bass was approximately 30% males at small sizes and 30-40% female at larger sizes. This contrasts with the expectation of a typical hermaphroditic species first maturing as females then a nearly complete transformation to male at older ages. In addition, it was shown (as well as in previous work by Alonzo et al. 2008, Brooks et al. 2008) that the presence of mature secondary males (males without bright coloration or nuchal humps, sometimes called sneaker males during

spawning) increases the resilience of a population to exploitation. If the large dominant male is removed and smaller mature males are available to spawn, the population is more robust than a population totally dependent on the large males or requiring mature females to change sex to replace the male. The black sea bass simulation study concluded that the northern stock of black sea bass was not a typical protogynous hermaphrodite and was more resilient to exploitation. The resilience is a result of a sex ratio which is not completely male at larger sizes and contributions by the secondary males which in combination bring black sea bass life history more in line with gonochoristic species.

Incorporating parameters into the assessment model to account for hermaphroditism would require empirical data that is not available, particularly the type of males present. A study by Brooks et al. (2008) showed that when adequate information about sex composition is unavailable, it is prudent to characterize spawning stock biomass (SSB) as the combined male and female mature biomass. To that end, this assessment incorporated that definition of SSB for modeling and development of biological reference points.

As part of the simulation work conducted using a Stock Synthesis model (Appendix A9) the implication of hermaphroditism was examined. The operating model used to simulate pseudo-data included sex structure and hermaphroditism, but sex-aggregated models performed well for recovering simulated population parameters. Simulation indicates that model results are largely unaffected by the presence or absence of the hermaphroditism criteria. Growth models between sexes are comparable so once sex change occurs, there is little change in productivity. Additionally, with SSB defined as male and female mature biomass, a change of sex does not remove fish from the SSB calculation. Nevertheless, the spawning success in black sea bass is dependent on a balanced sex ratio ensuring the presence of both sexes during spawning.

**TOR 6.** Estimate annual fishing mortality, recruitment and stock biomass (both total and spawning stock), using measures that are appropriate to the assessment model, for the time series (integrating results from TORs-1,-4, & -5 as appropriate), and estimate their uncertainty. Include a historical retrospective analysis and past projection performance evaluation to allow a comparison with most recent assessment results.

### *Model Development*

A statistical catch at age model (ASAP) was developed for the black sea bass assessment reviewed during SAW53. The SARC 53 review panel concluded that the results were not adequate for management purposes. Spatial patterning within the stock was suggested as a reason the model did not perform to their standards. Consequently, there are no historical retrospective or past projections to evaluate, however previous biological reference points are noted under TOR7.

Fishing mortality rates and stock sizes were estimated using the Age Structured Assessment Program (ASAP) statistical catch at age model (Legault and Restrepo 1998) available in the NOAA Fisheries Toolbox (<http://nft.nefsc.noaa.gov/>). ASAP is an age-structured model that uses forward computations assuming the separability of fishing mortality into year and age components to estimate population sizes given observed catches, catch-at-age, and indices of abundance. The separability assumption is partially relaxed by allowing for fleet-specific computations and by allowing the selectivity-at-age to change in blocks of time. The user can choose to estimate or hold fixed the various parameters and input uncertainty estimates associated with data which allows for configurations ranging from relatively simple age-structured production models to fully parameterized statistical catch-at-age models. The objective function is the sum of the negative log-likelihood of the fit to various model components. Catch-at-age and survey age compositions are generally modeled assuming a multinomial distribution, while most other model components are assumed to have lognormal error including penalties when used. Recruitment deviations are also assumed to follow a lognormal distribution, with annual deviations estimated as a bounded vector to force them to sum to zero (this centers the predictions on the expected stock-recruitment relationship). For more technical details refer to the ASAP technical manual.

The current assessment developed an ASAP model with several scenarios. The first scenario was modeling the stock under the assumption of a single unit stock, similar to previous approaches (note: this approach served as a bridge to previous assessments). Second, two unique area-specific models were developed for north and south of Hudson Canyon. Finally, a model was developed to account for exchange between the areas within the trawl fleets as well as within the NEFSC offshore surveys. Based on the performance of the two-area approach relative to more



complicated models, as evaluated by simulation testing (Appendix A6), parsimony and practical familiarity with ASAP, the WG concluded that the use of the two area models combined for status determination provided the best results for use in management. The WG also determined that the conclusions were robust to the modeling approach. The various approaches are presented in the following text, however the Overall and Area exchange approaches are provided for comparison to the final models, which are the North and South area models.

The catch at age data described in TOR 2 (north trawl and non-trawl; south trawl and non-trawl; overall trawl and non-trawl), total catch weight by fleet and survey indices at age described in TOR 3 were input to the model. Maturity at age, weight at age and natural mortality (0.4) described in the Introduction were input for each area. The single stock model (Overall model) represented the sum of catch at age and catch weight by fleet across area.

Model exploration began with a series of trial runs (Table A35) using only NEFSC surveys to establish baseline settings which produced a model solution. The Overall model began with two selectivity blocks with a change occurring in 1997-1998 when management was introduced. Incrementally the full suite of survey indices at age were added, with adjustments to index CVs to produce root mean square error values approaching 1.0. Initial NEFSC indices were an *Albatross* and *Bigelow* combined series. Input effective sample sizes (ESS) were chosen to conform with estimated values which were based on the approach of McAllister and Ianelli (1997). The model evolved (Table A36) as catch selectivities and index selectivity patterns were explored, beginning with selectivities that were freely estimated. The final model selectivities were fixed as flat topped beyond a chosen age and freely estimated in younger ages. The age of full selection was 4 in indices at age, age 4 in the non-trawl fleet as well as the trawl fleet from 1989 to 1997 then increasing to age 5.

The final Overall model incorporated a split series in the NEFSC spring index resulting in an *Albatross* series (1989-2008) and a *Bigelow* series (2009-2015). Additional indices at age included MADMF, NEAMAP, RI, CT, NEFSC winter, Recreational CPA and NJ. Indices from NY, DE, MD and VIMS were limited to age 1 indices only.

Variation in the catch at age, by fleet, is specified by the coefficient of variation values input for each fleet and year. Commercial trawl catch was given a CV equal to 0.05 since it is based on a census of dealer reports (discards are a function of the same reported landings). However, fleet 2

includes recreational catch which is based on a survey design that has undergone multiple changes. The input CV on fleet 2 was increased with an input value of 0.2 from 1989 to 2008 then a value of 0.15 from 2009 to 2015 to reflect better commercial reporting and improved estimates in the MRIP survey.

Recruitment in black sea bass throughout the stock is regionally influenced and not well determined by traditional stock recruitment models. Consequently, recruitment in ASAP was modeled as deviations from the geometric mean (steepness fixed at 1.0). The model allows deviations to be constrained by applying a penalty on the deviations from the stock-recruitment curve, however this constraint was weak due to an input CV of 1.0.

Development of area specific models followed a similar sequence (Tables A37-A40). Selectivity was freely estimated for catch and indices, then fleets and indices with estimates at bounds were fixed. Additional variance was added to indices to produce model estimates consistent with input values and effective sample sizes were adjusted to reflect model output. The final model for the north was model 26 and for the south model 24.

The split into northern and southern areas was an attempt to address the spatial components of the stock as it was understood that the southern offshore area during winter would likely contain fish originating in the north. In an attempt to address this issue, an area exchange model was developed to permit the exchange of individuals between northern and southern areas. For black sea bass, the intent of this exchange was to account for fish caught in the southern area during the winter/spring offshore trawl fishery and NEFSC bottom trawl surveys that originated in the north. The modeling framework did not explicitly incorporate movement into the dynamics of the stock, but instead moved both fishery dependent and independent catch from the southern to the northern component before running two separate, area-specific ASAP models. The exchange of individuals impacted only total catch and aggregate indices, and did not alter age compositions.

Two components impacted the amount of catch exchanged between the areas. The first component was the overall proportion of southern catch that was assumed to originate from the northern area. The second component was related to annual variability in this assumed exchange proportion. The extent of black sea bass seasonal movement is thought to be related to water temperature. Therefore, to account for the impact of temperature on seasonal movement,

average bottom temperature from the NEFSC spring bottom trawl survey was used as an annual covariate that modified the assumed exchange proportion. The magnitude of the covariate influence on the exchange proportion was the second component that impacted the amount of catch exchanged. This magnitude was dictated by an assumed range in possible annual fluctuations of the exchange proportion and effectively acted as a covariate scalar. As the range of possible fluctuations in the exchange proportion increased, the more the annual proportions could vary from the assumed average exchange proportion and therefore, the greater the overall impact of the covariate.

For a given average exchange proportion ( $\bar{p}$ ) and range in annual fluctuations ( $r$ ), the annual exchange proportions ( $p$ ) were calculated as:

$$p = lb + \frac{ub - lb}{1 + e^{cov}}$$

where  $cov$  represented the covariate as standardized (z-scored) average bottom temperature and  $lb$  and  $ub$  represented the lower and upper bounds, respectively, of the annual exchange proportion calculated as:

$$lb \text{ or } ub = \bar{p} \pm 0.5r.$$

In warmer years, it is believed that black sea bass do not migrate as far south (Miller et al. 2016); therefore, increased covariate values resulted in decreased annual exchange proportions. These annual exchange proportions were then used to develop adjusted time series of catches for the area-specific models that just represented individuals originating from that specific area.

For a given set of fishery dependent and independent catches, a series of exchange proportions and ranges were evaluated. Exchange proportion and range value combinations were only investigated if all possible annual proportions fell between zero and one. Candidate models based on adjusted time series were evaluated using the resulting combined (unweighted sum) total likelihood values of the area-specific models. The best exchange proportion and range combination was selected as the model set that produced the lowest combined likelihood. The robustness of assessment outputs, including recruitment, fishing mortality and spawning stock biomass, was evaluated across changes in exchange proportion and range values.

The final North and South models were incorporated into the area exchange model. The model was run for exchange rates between 0% (base models) and 50% by 5% increments and with

annual ranges from 0 to 50% in 10% increments. The model likelihoods for each area and the sum were evaluated to determine the optimal north-south combination.

### *Model Results*

The predicted total index proportions at age in the OVERALL model generally matched the observed values among all the indices. The predicted indices tended to match the trend among all the observed indices but generally were unable to match the large deviations associated with strong cohorts such as the 2011 year class among northern state indices. The root mean square error (RMSE) values ranged from 0.47 for REC CPA to 1.97 for MADMF indices (Figure A53). The model resulted in annual catch estimates that matched fleet 1 (standardized residuals ranged from 0.31 to -0.11) but varied more for fleet 2 (standardized residuals from 1.95 to -2.85). Both fleets tended to have positive residuals prior to 2002 and negative after 2002 (Figure A54-A55). Effective sample size for fleet 1 was constant at 60 while fleet 2 equaled 100. Catchability estimates of all the indices were less than 1, however the *Albatross*  $q$  was larger than the *Bigelow*  $q$  which was opposite of expectations. Detailed diagnostics from the model are presented in Appendix A10.

The model provided total stock number estimates showing a slowly increasing population from a low in 2005 until 2011, at which point the total abundance increased dramatically due to an incoming 2011 cohort (Figure A56). Abundance declined thereafter but remained larger in 2015 than pre-2011 abundance. Average fishing mortality (ages 4-7) was very high in the early 1990s (as high as 1.44) but declined following management implementation in 1997. Estimated  $F$  in 2007 was 0.74 but declined steadily to 0.24 by 2011 (Figure A56). The 2015 average fishing mortality was estimated as 0.20.

Spawning stock biomass (male and female combined) steadily increased from 1994 to 2002 when it reached 9,796 mt (Figure A56). There was a decline until 2007 (4,081 mt) followed by a steady increase to the 2015 estimate of 19,585 mt. Total biomass and exploitable biomass followed a similar pattern with 2015 total biomass equal to 24,155 mt and exploitable biomass of 23,658 mt (Figure A56). Average recruitment over the time series equaled 25.9 million age 1 fish, with strong cohorts in 1999, 2001 and an exceptional year class in 2011 (Figure A56).

A retrospective analysis of the model, using a 7 year peel, indicated very little impact on the results from additional years in the time series (Figures A57-A62). However, using a 7 year peel resulted in the loss of the Bigelow index and the reduction of the NEAMAP indices to one year. Consequently, the analysis uses a different model once the 7 year peel is completed so the retrospective results should be viewed with caution. The F estimates, with a Mohn's rho equal to -0.11, showed no obvious pattern. Biomass, total stock number and recruitment were similar and had Mohn's rho values ranging from 0.055 (recruitment) to 0.091 (SSB).

A Monte Carlo Markov Chain (MCMC) was implemented using 1000 iterations and a thinning rate of 200. The 2016 recruitment was estimated as the geometric mean recruitment from 2000 to 2015. The 90% confidence interval of the median terminal year Fmult (0.20) ranged from 0.151 to 0.264 (Figure A63-A64). The median 2015 SSB (20,092 mt) was estimated with 90% CI between 16,501 mt to 24,675 mt (Figure A65-A66). The confidence interval for January 1 median biomass in 2015 (24,693 mt) ranged from 20,788 mt to 29,674 mt (Figure A67-A68). The terminal year values of F, SSB and total biomass adjusted for retrospective were all within the 90% CI of the base values.

The NORTH model was developed using trawl and non-trawl catches taken north of Hudson Canyon and indices from the northern strata for NEFSC spring (*Albatross* and *Bigelow* series) and NEAMAP. The NEFSC winter northern index was not included in the final model due to poor diagnostics. Also included were indices from MA, RI, CT, NY and Rec CPA of states NY and north. Model results and diagnostics are provided in Figures A69-A97. The predicted total index proportions at age and the observed values in the model matched reasonably well among indices but annual matches varied among indices. The annual proportions at age from the NEFSC *Albatross* time series fit poorly as did several other indices in the early part of the series. In comparison, indices collected since the mid to late 2000s captured the age composition reasonably well. The predicted indices tended to match the trend among all the observed indices and were generally able to match the strong 2011 year class although not always of the magnitude within an index. The root mean square error (RMSE) values ranged from 0.74 for the Rec CPA to 1.88 for NY age 1 indices. The final model configuration resulted in annual catch estimates that matched fleet 1 (standardized residuals ranged from 0.06 to -0.18) and less so for fleet 2 (standardized residuals from 0.90 to -0.67). Both fleets tended to have positive residuals

after 2009 and negative prior to 1997. Effective sample size for fleet 1 was constant at 50 while fleet 2 equaled 100. Catchability estimates of all the indices were less than 1. In addition estimated *Albatross* catchability was two times smaller than the Bigelow catchability which was comparable to the between vessel calibration coefficient (3.4), especially when considering the uncertainty in the  $q$  estimates from the model and changes in selectivity. Additional diagnostics from the model are presented in Appendix A11.

The model provided total stock abundance estimates showing a steadily increasing population beginning around 1993 until 2011, at which point the total abundance increased dramatically due to an incoming 2011 cohort (Figure A98). Abundance declined thereafter and 2015 total abundance was equivalent to abundance in 2011. Average fishing mortality (ages 4-7) was very high in the early 1990s (as high as 1.32 in 1992) but steadily declined to 0.17 by 2011 (Figure A98). Estimated fishing mortality since 2012 has remained steady at around 0.23.

Spawning stock biomass (male and female combined) steadily increased from 1994 (259 mt) to 13,119 mt in 2014 (Figure A98). There was slight decline to 11,719 mt in 2015. Total biomass and exploitable biomass followed a similar pattern with a peak total biomass equal in 2014 of 21,376 mt and exploitable biomass of 14,737 mt (Figure A98). Total biomass in 2015 decreased to 17,306 mt while exploitable biomass increased slightly to 15,079 mt. Average recruitment over the time series equaled 8.353 million age 1 fish, with an exceptional year class in 2011 of 49.88 million age 1 fish (Figure A98). Recruitment in 2013 was also above average followed by a low recruitment year in 2014.

A retrospective analysis of the model, using a 7 year peel, indicated a general mis-specification of fishing mortality, abundance and biomass. However, using a 7 year peel resulted in the loss of the Bigelow index and the reduction of the NEAMAP indices to one year. Consequently, the analysis uses a different model once the 7 year peel is completed so the retrospective results should be viewed with caution. The  $F$  estimates, with a Mohn's  $\rho$  equal to 0.703, showed a pattern of over-estimation (Figure A99). Biomass, total stock number and recruitment were under-estimated and had Mohn's  $\rho$  values ranging from -0.484 (recruitment) to -0.325 (exploitable biomass) (Figures A100-A104). The direction of the  $\rho$  values are opposite to the pattern typically seen and results in decreased  $F$  and increased SSB following retrospective adjustments.

A Monte Carlo Markov Chain (MCMC) was implemented using 1000 iterations and a thinning rate of 200. The 2016 recruitment was estimated as the geometric mean recruitment from 2000 to 2015. The 90% confidence interval of the median terminal year  $F_{mult}$  (0.247) ranged from 0.181 to 0.337 (Figures A105-A106). The median 2015 SSB (11,959 mt) was estimated with 90% CI between 9,606 mt to 15,003 mt (Figures A107-A108). The confidence interval for January 1 median biomass in 2015 (17,656 mt) ranged from 14,569 mt to 21,449 mt (Figures A109-A110). The terminal year (2015) values of  $F$ , SSB and total biomass were adjusted for retrospective patterns. The retro adjusted value of  $F$  (0.14), SSB (19,211 mt), total Jan 1 biomass (27,733 mt) fell outside the 90% confidence bounds of the terminal year estimate (Figure A110a)

The final SOUTH model was developed using trawl and non-trawl catches taken south of Hudson Canyon and indices from the NEFSC spring (*Albatross* and *Bigelow* series), NEAMAP (age 1 only because of poor diagnostics in older fish), and NEFSC winter southern strata. Also included were indices from NJ, DE, MD, VIMS and Rec CPA from states NJ and south. The only indices beyond age 1 were from the NEFSC *Albatross* and *Bigelow*, NEFSC winter, NJ and Rec CPA. The remainder provided indices of age 1 abundance. Model results and diagnostics are provided in Figures A111-A138. The predicted total index proportions at age and the observed values matched reasonably well among indices but annual matches varied among indices. The annual proportions at age from the NEFSC *Albatross* time series fit poorly. The predicted indices tended to match the trend among all the observed indices and were generally able to match the strong 1999 and 2001 year classes although not always of the magnitude within an index. The 2011 cohort was apparent as above average in several indices but not of the magnitude seen in the northern model. The root mean square error (RMSE) values ranged from 0.68 for the Rec CPA to 1.45 for NEFSC *Bigelow* indices. The final model configuration resulted in annual catch estimates that matched fleet 1 (standardized residuals ranged from 0.27 (1996) to -0.13) and less so for fleet 2 (standardized residuals from 1.73 to -1.74). Both fleets had positive residuals prior to 2002 and negative from 2003 to 2015. Effective sample size for fleet 1 was constant at 30 while fleet 2 equaled 40. Catchability estimates of all the indices were less than 1. In addition estimated *Albatross* catchability was two times smaller than the *Bigelow* catchability which was comparable to the between vessel calibration coefficient of 3.4, especially when considering the

uncertainty in the  $q$  estimates from the model and changes in selectivity. Additional diagnostics from the model are presented in Appendix A12.

The model provided total stock number estimates showing a relatively stable population beginning between 1989 and 2002, at which point the total abundance declined reaching a low point in 2006 (Figure A139). Abundance steadily increased thereafter and 2015 total abundance approached the abundance seen in 2002. Average fishing mortality (ages 4-7) was very high in the early 1990s (as high as 1.43 in 1993) but declined to 0.55 by 1999 (Figure A139). Estimated fishing mortality increased again to 0.94 in 2007 but steadily declined to 0.23 by 2015.

Spawning stock biomass (male and female combined) steadily increased from 1989 (2,088 mt) to 5,801 mt in 2002 (Figure A139). There was a decline to 1,732 mt by 2007 followed by another steady increase reaching 4,834 mt by 2015. Total biomass and exploitable biomass followed a similar pattern with a peak total biomass in 2002 equal to 9,728 mt and exploitable biomass of 6,718 mt (Figure A139). Total biomass decreased to 3,614 mt in 2007 with an exploitable biomass of 2,171 mt in 2008 but both have since increased to 6,838 mt and 5,168 mt, respectively, in 2015. Average recruitment over the time series equaled 15.92 million age 1 fish, with higher recruitment between 1989 and 2002 (Figure A139). Recruitment in 2015 was above average with 21.57 million age 1 recruits.

A retrospective analysis of the model, using a 7 year peel, indicated a general mis-specification of fishing mortality, abundance and biomass (Figures A140-A145). However, using a 7 year peel resulted in the loss of the *Bigelow* index and the reduction of the NEAMAP indices to one year. Consequently, the analysis uses a different model once the 7 year peel is completed so the retrospective results should be viewed with caution. The  $F$  estimates, with a Mohn's rho equal to -0.419, showed a pattern of under-estimation. Biomass, total stock number and recruitment were over-estimated and had Mohn's rho values ranging from 0.550 (exploitable biomass) to 0.873 (recruitment).

A Monte Carlo Markov Chain (MCMC) was implemented using 1000 iterations and a thinning rate of 200. The 2016 recruitment was estimated as the geometric mean recruitment from 2000 to 2015. The 90% confidence interval of the median terminal year  $F_{mult}$  (0.233) ranged from 0.162 to 0.343 (Figures A146-A147). The median 2015 SSB (4,901 mt) was not as well estimated with 90% CI between 3,628 mt to 6,529 mt (Figures A148-A149). The 90% confidence interval for



January 1 median biomass in 2015 (6,920 mt) ranged from 5,432 mt to 8,845 mt (Figures A150-A151).

The terminal year values of F, SSB and total biomass were adjusted for retrospective patterns. The retro adjusted value of F (0.39), SSB (2,966 mt), total Jan 1 biomass (4,276 mt) fell outside the 90% confidence bounds of the terminal year estimate (Figure A110a)

#### *Area Exchange Model*

A model using 40% exchange in trawl catch and indices with a 10% range in annual exchange proportions produced the lowest value for the combined likelihoods (Table A41). The best model for the southern area occurred with a 50% exchange and 50% range which were at the upper bound for both exchange rate and range. The northern model minimum likelihood occurred by moving 35% of southern catch and indices to the north and no annual fluctuations in this exchange rate (0% range).

Results from the optimal area exchange model showed minimal impact on the final overall estimates but did produce some regional differences (Figures A152). The catch in the north increased substantially in the early part of the time series (1993 total catch increased 127%) but by 2015 the exchange increased the northern total catch by only 3.7%. The effect of the catch increased abundance estimates an average of 22% but only resulted in reducing F by 3%. The average trawl catch reduction in the south of 40% reduced total catch by 9% which resulted in an 8-10% decrease in total abundance, SSB, recruitment and exploitable biomass. Fishing mortality decreased by only 1%. Overall, temporal trends in area-specific assessment outputs were robust to both the selected exchange proportion and range, though they were more robust to changes in the assumed range than the exchange proportion. Additional diagnostic plots are available in Appendices A13-A14.

The area exchange approach had minimal impact on the area-specific retrospective values with the exception of the fishing mortality in the north. The Mohn's rho in the north decreased 25% from 0.70 to 0.53 but adjusted F only decreased from 0.14 to 0.13. Rho only decreased 15% in exploitable biomass, 9% in total stock number and 6% in age one recruitment. Rho values for the south model increased but only by 3% or less.

#### *Model Comparisons*

Although models were developed for a northern area and a southern area, the final product for determining stock status was a single combined estimate for the unit stock. These combined area results as well as the combined area exchange estimates were compared with results from the single overall model (Table A42). The F values from North and South were averaged to compare to the Overall model results.

In general, the overall model produced larger estimates of abundance and biomass and lower estimates of F than the area-specific models and the combined areas estimates from the exchange model were nearly identical to the overall model (Figure A153). Despite the difference in point estimates among models, the trends were nearly identical. The WG concluded that the area exchange model as configured did not produce any benefits beyond the simple area combined or the overall model. However, it did prove informative as a sensitivity run to evaluate the impact of catch misspecifications to each area and showed that the model was robust to migration effects.

There existed a trade-off between the use of the overall model and the combined area-specific models. The area-specific models resulted in retrospective patterns which were in opposite directions. However because of the reduced areas in each, the local indices of abundance were more informative (i.e. fit better) than in the overall model. In the overall model the residuals in the catch and indices were greater, however the retrospective patterns of each area essentially cancelled each other resulting in low rho values. The WG concluded that the combined area specific models provided the most information for management purposes. There were clearly local variations in recruitment strength of the 2011 cohort which was most apparent in the area models. Fortunately the overall, combined area and area exchange models all produced similar results.

#### *Sensitivity analysis*

A sensitivity analysis was conducted to evaluate the influence of individual indices on the model results for each area. The models were run with each index removed from the model (only one index missing per run). The results show that the model was robust and not dependent on any single index. Model results with any particular index dropped remained within the 90% confidence interval of the base model. Results are presented in Figures A154-A163.

The models were profiled over a range of possible values of natural mortality. Each model was run using a new  $M$  value ranging from 0.01 to 1.0. The analysis was conducted on the two area models, the areas combined and the overall model and calculated the difference between each model negative log-likelihood and the model with the minimum negative log-likelihood (Figure A164). In the southern model, the minimum negative log-likelihood occurred for  $M$  at 0.8 whereas the northern model minimum occurred at 0.15. The combination of the north and south produced a minimum at an  $M=0.4$  while the overall model minimum was at  $M=0.55$ .

An additional sensitivity test was made to examine the implications of the higher CV values used for the non-trawl catches. The CVs in the base model were 0.2 from 1989-2008 and 0.15 in 2009-2015). These values were varied from (0.05, 0.025) to (0.35, 0.3). The results (Figure A165) suggest that the North models  $F$  and  $SSB$  were robust to the range of values explored. The South model was impacted in the early 2000s when the  $SSB$  was highest, but remained robust to the range of CV over the past decade.

In addition to developing a model using the ASAP approach, the fishery and survey data was also explored using the Stock Synthesis framework (Methot and Wetzel, 2013). The model (Appendix A9) incorporated tagging data from a project in 2003-2005 (Moser and Shepherd, 2009) and modeled the population seasonally by length and sex, accounting for hermaphroditism. Although the model was not fully vetted by the BSB WG, and therefore not presented as a working model, the results were similar to ASAP. The exploratory model suggested that the stock has increased in the last decade in both areas from strong recruitment and that fishing mortality decreased to relatively low values in the last decade. Sensitivity analyses indicate that these results are robust to alternative model configurations.

**TOR 7.** Estimate biological reference points (BRPs; point estimates or proxies for  $BMSY$ ,  $BTHRESHOLD$ ,  $FMSY$ , and  $MSY$ ), including defining BRPs for spatially explicit areas if appropriate, and provide estimates of their uncertainty. If analytic model-based estimates are unavailable, consider recommending alternative measurable proxies for BRPs. Comment on the appropriateness of existing BRPs and the “new” (i.e., updated, redefined, or alternative) BRPs.

Black sea bass north of Cape Hatteras, NC have been managed as a single stock since the inception of the management plan in 1997. Although the evidence presented for TOR 1 suggests regional differences within the stock, there is no evidence that more than one stock exists north of Cape Hatteras. Although the preferred model approach of the WG has been the two area model with no exchange, development of reference points and subsequent evaluation of stock status was made on the unit stock (Cape Hatteras to the Gulf of Maine).

The WG examined the two-area models for a relationship between SSB and subsequent recruitment and concluded there was no evidence of any stock/recruitment relationship in the range of estimated stock size in the assessment time series (Appendices A11 & A12).

Consequently there was no direct calculation of MSY and associated F or biomass. Instead, an average value of F40% between the two areas was chosen as a proxy for  $F_{MSY}$ . In similar situations, and for species with similar life histories, the South Atlantic snapper-grouper fishery management plan uses F30% or F40% as  $F_{MSY}$  proxies when  $F_{MSY}$  cannot be directly estimated. Thorson et al. (2012) suggest that F35% is an appropriate proxy for  $F_{MSY}$  based on a meta-analysis of stock-recruitment information from Perciformes. Zhou et al. (2012) suggested that  $F_{MSY}$  is approximately  $0.922 \times M$  for Perciformes, which would correspond to  $F_{MSY} \sim 0.37$  for the assumed  $M=0.4$ . The estimate of F40% in the north equaled 0.355 and was 0.365 in the south. An average of 0.36 was adopted as the proxy  $F_{MSY}$ .

An approach was adopted for developing biomass reference points using empirical recruitment in a long term projection. Inputs for projections included averages from 2013 to 2015 total (Jan 1) and SSB weights at age, selectivity at age weighted by contributing catch, and natural mortality =0.4. Recruitment was chosen from a random draw for the combined area recruits between 2000 and 2015. The WG felt that recruitment over this time period was more reflective of current population dynamics. Catch input for 2016 was the Allowable Biological Catch (ABC) established by the MAFMC for 2016. Distributions of abundance at age from each model MCMC run were combined for a single stock projection. A 100 year projection with 1000 bootstrap iterations was run and the average of the final 20 years calculated as the biological reference points ( $SSB_{MSY}$  proxy,  $B_{MSY}$  proxy and  $MSY_{proxy}$ ) associated with the  $F_{MSY}$  proxy. In previous assessments using a length based model (SCALE (Northeast Data Poor Stocks Working Group 2009))  $B_{MSY}$  proxy was chosen as a biomass reference point. Traditionally spawning stock biomass is preferred because of the potential relationship with subsequent recruitment, however

hermaphroditism presents unique challenges in defining female SSB due to sex changes.

Analyses by Brooks et al. (2009) showed that SSB defined by both sexes provided a reasonable alternative. Consequently combined male and female SSB is recommended for use as a biomass reference point for black sea bass although both SSB and total biomass estimates are provided.

The final estimate of  $SSB_{MSY}$  proxy equaled 9,667 mt with  $\pm 2$  SD of 4,149 mt,  $B_{MSY}$  proxy of 17,256 mt with  $\pm 2$  SD of 7,453 mt, and an  $MSY$  proxy estimate of 3,097 mt with  $\pm 2$  SD of 1,299 mt (Table A43). The previous basis for determining if a stock is overfished is  $\frac{1}{2}$  of  $B_{MSY}$  proxy which in this stock equals 8,628 mt ( $\pm 2$  SD of 3,722 mt). An alternative approach for biomass reference point estimation using the single overall model produced results similar to the combined area model, although generally higher. Those estimates are  $SSB_{MSY}$  proxy = 10,296 mt,  $B_{MSY}$  proxy = 18,379 mt, and  $MSY$  = 3,298 mt.

Biological reference point estimates from the 2008 assessment black sea bass assessments were an  $F_{MSY}$  proxy = 0.42,  $SSB_{MSY}$  proxy of 12,537 mt and an  $MSY$  of 3,903 mt and the 2012 assessment (which was not accepted in SAW53) produced reference points of an  $F_{MSY}$  proxy = 0.32,  $SSB_{MSY}$  proxy of 8,128 mt and an  $MSY$  of 3,197 mt.

**TOR 8.** Evaluate overall stock status with respect to a new model or new models that considered spatial units developed for this peer review.

Retrospectively adjusted values of SSB and F from the ASAP model are being used for stock status determination. The stock of black sea bass north of Cape Hatteras, NC is not overfished, nor is it experiencing over fishing (Figures A169-A170). The retrospective adjusted estimates of F, SSB and B from each area were combined for comparison to biological reference points. The retro-adjusted 2015 biomass (32,010 mt) is currently 86% above  $B_{MSY}$  proxy and retro-adjusted 2015 SSB (22,176 mt) is 129% above the  $SSB_{MSY}$  proxy (9,667 mt). Retro-adjusted fishing mortality in 2015 ( $F_{MULT} = F_{Ages\ 4-7} = 0.27$ ) is 25% below the  $F_{MSY}$  proxy = 0.36.

The 2015 stock biomass prior to adjustment (24,143 mt) is 40% above  $B_{MSY}$  proxy (17,256 mt) and 2015 SSB (16,552 mt) is 71% over  $SSB_{MSY}$  proxy (9,667 mt). Fishing mortality in 2015 before adjusting for the retrospective bias ( $F_{MULT} = 0.24$ ) is 35% below  $F_{MSY}$  proxy = 0.36.

It should be noted that the area specific conditions are different. Relative to F40% (0.355 in the north, 0.365 in the south), the retro adjusted F in the north (0.14) is well below the threshold whereas retro adjusted F in the south (0.39) is slightly above (although well within the 80% CI of 0.30 to 0.52). Biomass reference points were not independently estimated for each area.

**TOR 9.** Develop approaches and apply them to conduct stock projections.

- a. Provide numerical annual projections (3-5 years) and the statistical distribution (e.g., probability density function) of the OFL (overfishing level) that fully incorporates observation, process and model uncertainty (see Appendix to the SAW TORs). Each projection should estimate and report annual probabilities of exceeding threshold BRPs for F, and probabilities of falling below threshold BRPs for biomass. Use a sensitivity analysis approach in which a range of assumptions about the most important uncertainties in the assessment are considered (e.g., terminal year abundance, variability in recruitment, and definition of BRPs for black sea bass).
- b. Comment on which projections seem most realistic. Consider major uncertainties in the assessment as well as the sensitivity of the projections to various assumptions.
- c. Describe this stock's vulnerability (see "Appendix to the SAW TORs") to becoming overfished, and how this could affect the choice of ABC.

Catch projections are presented for the unit stock, however they represent the sum of each area specific projection. Projections for the north and south area were made for years 2017 through 2020 under the assumption of either  $F_{MSY\ proxy}$  or status quo F. Since 2016 catch estimates were not available, 2016 catch was interpreted as the ABC (3,024 mt) split into area based on the 3 year average ratio of catch by area. The 2013 to 2015 average catch ratio was 71% in the north and 29% in the south, therefore 2016 catch in the north was 2,150 mt and the south as 874 mt. Additional runs were included which allowed a 10% and a 20% overage in the north 2016 catch since recent years have exceeded the ABC. The 2013-2015 average weights at age, selectivity (fleets combined, weighted by catch at age), and empirical recruitment from 2000 to 2015 were input to the stochastic projection software AGEPRO. A distribution of starting numbers at ages 2 to 8+ were produced by the MCMC run of each final model. Projections were also made in each area using rho adjusted estimates. The results of each area were summed for the final catch

projections. The final products were projections for area combined, area combined rho adjusted and a comparison to projections using the overall model under  $F_{MSY}$  and  $F$  status quo ( $F_{SQ}$ ).

Results without rho adjustments from the area combined model under  $F_{MSY\ proxy}$  (0.36) shows a substantial increase in total catch beginning in 2017 at 4,446 mt ( $\pm 2$  SD 1,687 mt) (Table A44, Figure A171). However as the 2011 cohort abundance declines, catch declines such that 2018 projected catch would drop to 3,822 mt ( $\pm 2$  SD 1,299 mt) and 3,493 mt ( $\pm 2$  SD 1,481 mt) in 2019. Total biomass would also decline beginning in 2016 (23,374 mt  $\pm 2$  SD 7,351 mt) and declining to 18,764 mt ( $\pm 2$  SD 8,977 mt) by 2019 which is above  $B_{MSY\ proxy}$  (17,256 mt, Table A44, Figure A172). Spawning biomass follows a similar trajectory, declining from 15,055 mt ( $\pm 2$  SD 5,621 mt) in 2016 to 10,782 mt ( $\pm 2$  SD 5,083 mt) by 2019 (Table A44, Figure A173).

Results without rho adjustments from the area combined model under  $F_{SQ}$  (0.24) shows a substantial increase in total catch beginning in 2017 at 3,127 mt ( $\pm 2$  SD 1,189 mt) (Table A45, Figure A174). However as the 2011 cohort abundance declines, catch declines such that 2018 projected catch would drop to 2,904 mt ( $\pm 2$  SD 996 mt) and 2,809 mt ( $\pm 2$  SD 1,147 mt) in 2019. Total biomass would also decline beginning in 2016 (23,374 mt  $\pm 2$  SD 7,351 mt) and declining to 20,848 mt ( $\pm 2$  SD 9,423 mt) by 2019 which is above  $B_{MSY\ proxy}$  (17,256 mt, Table A45, Figure A175). Spawning biomass follows a similar trajectory, declining from 15,055 mt ( $\pm 2$  SD 5,168 mt) in 2016 to 12,805 mt ( $\pm 2$  SD 5,663 mt) by 2019 (Table A45, Figure A176).

Results with the rho adjustments from the area combined model under  $F_{MSY\ proxy}$  (0.36) have the same trend with higher abundance (Table A43). A substantial increase in total catch would occur beginning in 2017 at 5,467 mt ( $\pm 2$  SD 1,984 mt) (Table A46, Figure A171). However as the 2011 cohort abundance declines, catch declines such that 2018 projected catch would drop to 4,494 mt ( $\pm 2$  SD 1,457 mt) and 3,901 MT ( $\pm 2$  SD 1,502 mt) in 2019. Total biomass would decrease from 29,350 mt ( $\pm 2$  SD 9,028 mt) in 2016 to 20,788 mt ( $\pm 2$  SD 9,063 mt) by 2019 which is above  $B_{MSY\ proxy}$  (Table A46, Figure A172). Spawning biomass follows a similar trajectory, declining from 18,670 mt ( $\pm 2$  SD 6,721 mt) in 2016 to 11,849 mt ( $\pm 2$  SD 5,143 mt) by 2019 (Table A46, Figure A173).

Results with the rho adjustments from the area combined model under  $F_{SQ}$  (0.27) have the same trend with higher abundance (Table A47). A decrease in total catch would occur beginning in 2017 at 3,006 mt ( $\pm 2$  SD 1,126 mt) (Table A47, Figure A174). Catch continues to decline

through 2018 as projected catch would drop to 2,859 mt ( $\pm 2$  SD 953 mt) and 2,764 mt ( $\pm 2$  SD 986 mt) in 2019. Total biomass would decrease from 29,246 mt ( $\pm 2$  SD 8,993 mt) in 2016 to 24,704 mt ( $\pm 2$  SD 9,839 mt) by 2019 which is above  $B_{MSY}$  proxy (Table A47, Figure A175). Spawning biomass follows a similar trajectory, declining from 18,587 mt ( $\pm 2$  SD 6,690 mt) in 2016 to 15,349 mt ( $\pm 2$  SD 6,099 mt) by 2019 (Table A47, Figure A176).

The Overall model, with no rho adjustments, was similar to the area combined estimates with the rho adjustment. Projected catch in 2017 at  $F_{MSY}$  would be 6,615 mt compared to the rho adjusted combined area model with 5,467 mt (Figure A171). SSB in 2017 in the overall model equaled 19,125 mt compared to the rho adjusted combined estimate of 15,918 mt (Figure A172).

Given the convergence of estimates from the overall model and the rho adjusted area combined model, the rho adjusted values appear to provide reasonable projection estimates. The projections assume the 2016 ABC was not exceeded. If the 2016 ABC was exceeded by 20%, the projected rho-adjusted 2017 catch at  $F_{SQ}$  would only decrease 2% to 2,993 mt. The projections also assume the selectivity estimated for 2013-2015 would apply to the future catches. Due to overages in 2015, states from NJ north adjusted regulations to reduce harvest in 2016, which may have influenced selectivity to some degree.

The assessment model and consequently the projections do not explicitly account for hermaphroditism in sea bass beyond using male and female SSB. However, hermaphroditism is unlikely to have a large impact on the assessment results but should be considered in developing management options. Finally, there is evidence in 2016 survey indices that in some areas the 2015 cohort is above average. This incoming year class would not have much influence on projections until 2018, at which time it will be included in the assessment estimates.

**TOR 10.** Review, evaluate and report on the status of the SARC and Working Group research recommendations listed in recent SARC reviewed assessments and review panel reports.

Identify new research recommendations.

Research recommendations from SAW 53:

**1.** *The panel recommends multiple age-structured models be evaluated for use in a future model. Specifically, we recommend:*

*a. A simple model such as a separable model with smoothing on  $F$  among years.*

Not attempted for this assessment, however evaluation of catch curves is a routine diagnostic.



b. *A more complex, spatially structured model with 6 month time step within independent stock areas in spring and mixing in winter with natal homing, if data area adequate to support such a model.*

Completed in the form of the area exchange model and explored within the SS3 model.

c. *Consideration should be given to including tag return data in an age-structured (and possibly spatially-structured) assessment model.*

Tag results were considered in developing the area exchange bounds but not explicitly within the model. Explored within the SS3 model.

2. *The Panel recommends evaluation of a species specific survey, such as a pot survey to provide increased information on abundances and biological characteristics.*

Previously existed but was terminated by the MAFMC following a review of the program design.

3. *Continue and expand the tagging program to provide:*

a. *increased age information.*

b. *increased resolution on mixing rates among putative populations.*

No additional designed tagging projects were conducted.

4. *Continue and expand genetic studies to evaluate the potential of population structure north of Cape Hatteras.*

Some genetic work being evaluated for the Gulf of Maine by Marissa McMahon, Northeastern University.

5. *Continued research on rate, timing and occurrence of sex-change in this species. Recent research findings discussed at the SARC lead to the hypothesis that protogyny is not obligate in this species – some individuals may never have been female before maturing as a male.*

Some further work conducted by NEFSC as well as Rutgers University.

6. *The validity of the age data used in the assessment requires further evaluation, in particular the reliability of scale-based ageing needs to be determined. A scale- otolith intercalibration exercise might be of utility.*

Working paper included which explores this issue.

SAW 62 Black Sea Bass Working Group research recommendations:

1. Expand on previous genetic studies with smaller spatial increments in sampling.
2. Consider the impact of climate change on black sea bass, particularly in the Gulf of Maine.
3. Evaluate population sex change and sex ratio, particularly comparing dynamics among communities.
4. Study black sea bass catchability in a variety of survey gear types.
5. Investigate and document social and spawning dynamics of black sea bass.
6. Increased work to understand habitat use in sea bass and seasonal changes.
7. Evaluate use of samples collected by industry study fleets.

## Acknowledgments

Special thanks to Joshua Dayton of the NEFSC Population Biology Branch for supplying age results, Alicia Miller for developing distribution maps and Kiersten Curti for development of the area exchange model and index uncertainty evaluations.

## Literature (Cited or Background literature)

Alonzo, S.H., T. Ish, M. Key, A.D. MacCall, and M. Mangel. 2008. The importance of incorporating protogynous sex change into stock assessments. *Bulletin of Marine Science* 83(1): 163-179.

Bell, R.J., D.E. Richardson, J.A. Hare, P.D. Lynch and P.S. Fratantoni. 2014. Disentangling the effects of climate, abundance, and size on the distribution of marine fish: an example based on four stocks from the Northeast US shelf. *ICES Journal of Marine Science*, 72(5):1311-1322.

Blaylock, J. and G.R. Shepherd. 2016. Evaluating the vulnerability of an atypical protogynous hermaphrodite to fishery exploitation: results from a population model for black sea bass (*Centropristis striata*). *Fishery Bulletin* 114(4): 476-489.

Brooks, L., K.W. Shertzer, T. Gedamke, and D.S. Vaughan. 2008. Stock assessment of protogynous fish: evaluating measures of spawning biomass used to estimate biological reference points. *Fishery Bulletin* 106(1):12-23.

Bugley, K. and G. Shepherd. 1991. Effect of catch-and-release angling on the survival of black sea bass. *North American Journal of Fisheries Management*. 11:468-471.

Collins, M.R., J.C. McGovern, G.R. Sedberry, H.S. Meister and R. Pardieck. 1999. Swim bladder deflation in black sea bass and vermilion snapper: potential for increasing post-release survival. *North American Journal of Fisheries Management* 19:828-832.

Drohan, A. F., J. P. Manderson, and D. B. Packer. 2007. *Essential Fish Habitat Source Document: Black Sea Bass, *Centropristis striata*, Life History and Habitat Characteristics. Second Edition. NOAA Technical Memorandum NMFS-NE-200.*

Gerritsen, H.D., D. McGrath and C. Lordan. 2006. A simple method for comparing age-length keys reveals significant regional differences within a single stock of haddock (*Melanogrammus aeglefinus*). *ICES Journal of Marine Sciences*. Vol 63 (6): 1096-1100.

Gilmore, G.R. and R.S. Jones. 1992. Color variation and associated behavior in the Epinephelinae Groupers, *Mycteroperca microlepis* (Good and Bean) and *M. phenax* Jordan and Swain. *Bulletin of Marine Science* 51(1):83-103.

Hamilton, S.L. J.E. Caselle, J.D. Standish, D.M. Schroeder, M.S. Love, J.A. Rosales-Casian and O.Sosa-Nishizaki. 2007. Size-selective harvesting alters life histories of a temperate sex-changing fish. *Ecological Applications*. 17(8): 2268-2280.

Hartman, K.J. and S.B. Brandt. 1995. Trophic resource partitioning, diets, and growth of sympatric estuarine predators. *Transactions of the American Fisheries Society*. 124: 520-537.

Heppell, S.S., S.A. Heppell, F.C. Coleman and C.C. Koenig. 2006. Models to compare management options for a protogynous fish. *Ecological Applications*. 16(1):238-249.

Heppell, S. and S. Heppell. 2013. Issues in assessment modeling for protogynous fishes. In: Shepherd, G., K. Shertzer, J. Coakley, and M. Caldwell (Editors). 2013. Proceedings from a workshop on modeling protogynous hermaphrodite fishes, Raleigh, NC. 33 p. Available from: Mid-Atlantic Fishery Management Council, 800 North State Street, Suite 201, Dover, DE 19901, or online at <http://www.mafmc.org>

Hewitt, D.A. and J.J. Hoenig. 2005. Comparison of two approaches for estimating natural mortality based on longevity. *Fishery Bulletin*. 103:433-437.

Hoenig, J.M. 1983. Empirical use of longevity data to estimate mortality rates. *Fishery Bulletin*. 82:898-903.

Hoenig, J.M., N.J. Barrowman, W.S. Hearn and K.H. Pollock. 1991. Multiyear tagging studies incorporating fishing effort data. *Canadian Journal of Fisheries and Aquatic Science*. 55:1466-1476.

Huntsman, G.E. and W.E. Schaaf. 1994. Simulation of the impact of fishing on reproduction of a protogynous grouper, the graysby. *North American Journal of Fisheries Management*. 14:41-52.

Jaccard, Paul (1912), "The distribution of the flora in the alpine zone", *New Phytologist* 11: 37-50.

Keigwin B, G.R. Shepherd, M. J. Wuenschel. 2016. Geomorphometric analysis indicates overlap in body shape between sexes of black sea bass (*Centropristis striata*). US Dept Commer, Northeast Fish Sci Cent Ref Doc. 16-07; 26 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026, or online at <http://www.nefsc.noaa.gov/publications/>

Lavenda, N. 1949. Sexual difference and normal protogynous hermaphroditism in the Atlantic sea bass, *Centropristes striatus*. *Copeia* 3:185-194.

Legault, C.M. and V.R. Restrepo. 1998. A flexible forward age-structured assessment program. ICCAT Col. Vol. Sci. Pap. 49:246-253.

Lindquist, D.G., L.B. Cahoon, I.E. Clavijo, M.H. Posey, S.K. Bolden, L.A. Pike, S.W. Burk and P.A. Cardullo. 1994. Reef fish stomach contents and prey abundance on reef and sand substrata associated with adjacent artificial and natural reefs in Onslow Bay, North Carolina. *Bulletin of Marine Science*, 55(2-3): 308-318.

Lorenzen, K. 1996. The relationship between body weight and natural mortality in juvenile and adult fish: a comparison of natural ecosystems and aquaculture. *Journal of Fish Biology*. 49:627-647.

MAFMC 1996. Amendment 9 to the Summer Flounder Fishery Management Plan: Fishery Management Plan and Final Environmental Impact Statement for the Black Sea Bass Fishery. 17 June 1996. <http://www.mafmc.org/fisheries/fmp/sf-s-bsb>

McAllister, M.K and J.N.Ianelli. 1997. Bayesian stock assessment using catch–age data and the sampling –importance resampling algorithm. *Canadian Journal of Fisheries and Aquatic Science* 54:284-300.

McBride, R., M.Wuenschel and G. Fitzhugh. 2013. The How and Why of Hermaphroditism. *In*: Shepherd, G., K. Shertzer, J. Coakley, and M. Caldwell (Editors). 2013. Proceedings from a workshop on modeling protogynous hermaphrodite fishes, Raleigh, NC. 33 p. Available from: Mid-Atlantic Fishery Management Council, 800 North State Street, Suite 201, Dover, DE 19901, or online at <http://www.mafmc.org>

McCartney, M. and M.L.Burton. 2011. Population genetic structure of black seabass (*Centropristis striata*) on the eastern U.S. coast, with an analysis of mixing between stocks north and south of Cape Hatteras, North Carolina. SEDAR25-RD42 working paper (cited with permission of the author).

Mercer, L.P. 1978. The reproductive biology and population dynamics of black sea bass, *Centropristis striata*. Ph.D. thesis, Va. Inst. Mar. Sci., College of William and Mary, Gloucester Point, VA 23062.

Methot, R.D. Jr and Wetzel, C.R. 2013. Stock Synthesis: A biological and statistical framework for fish stock assessment and fishery management. *Fisheries Research* 142: 86–99.

Miller, A. S., G.R. Shepherd and P.S. Fratantoni. 2016. Offshore habitat preference of overwintering juvenile and adult black sea bass, *Centropristis striata*, and the relationship to year-class success. *PLoS one* 11(5):e0156355.

Moser, J. and G.R. Shepherd. 2009. Seasonal Distribution and Movement of Black Sea Bass (*Centropristis striata*) in the Northwest Atlantic as Determined from a Mark-Recapture Experiment. *Journal of Northwest Atlantic Fisheries Science* 40:17-28.

Musick, J.A., and L.P. Mercer. 1977. Seasonal distribution of black sea bass, *Centropristis striata*, in the Mid-Atlantic Bight with comments on the ecology and fisheries of the species. *Transactions of the American Fisheries Society* 106(1):12-25.

Nelson, D.A., D. Perry, and E. Baker. 2003. Natural spawning of black sea bass, *Centropristis striata*, at the NMFS Milford Laboratory and the UMASS Dartmouth Laboratory with observations on spawning behavior. *Journal of Shellfish Research* Vol. 22, no. 1, pp. 297-298.

Northeast Data Poor Stocks Working Group. 2009. The Northeast Data Poor Stocks Working Group Report, December 8-12, 2008 Meeting. Part A. Skate species complex, deep sea red crab, Atlantic wolffish, scup, and black sea bass. *US Dept Commer, Northeast Fish Sci Cent Ref Doc.* 09-02; 496 p.

Northeast Fisheries Science Center (NEFSC). 2006. 43rd Northeast Regional Stock Assessment Workshop (43rd SAW): 43rd SAW assessment report. *US Dep Commer, Northeast Fish Sci Cent Ref Doc* 06-25; 400 p.

Northeast Fisheries Science Center (NEFSC). 2012. 53rd Northeast Regional Stock Assessment Workshop (53rd SAW) Assessment Report. *US Dept Commer, Northeast Fish Sci Cent Ref Doc.* 12-05; 559 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026, or online at <http://nefsc.noaa.gov/publications/>

Provost, M.M. 2013. Understanding sex change in exploited fish populations: A review of east coast fish stocks and assessment of selectivity and sex change in black sea bass (*Centropristis striata*) in New Jersey. MS thesis, Rutgers University. 77 pp.

Provost.M.M and O.P.Jensen. 2015. The impacts of fishing on hermaphroditic and treatment of sex change in stock assessments. *Fisheries* 40(11), 2015.

Rountree, R.A. 1999. Diets of NW Atlantic fishes and squid. <http://www.fishecolology.org>. Accessed October 2016

Rudershausen, P.J. and J.A.Buckel. 2007. Discard composition and release fate in the snapper and grouper commercial hook and line fishery in North Carolina, USA. *Fisheries Management and Ecology*, 14:103-113.

Rudershausen P.J., J.A. Buckel, and T. Burgess. 2010. Estimating discard mortality of black sea bass (*Centropristis striata*) and other reef fish in North Carolina using a tag-return approach (SEDAR25-RD10). North Carolina Sea Grant FRG #'s 07-FEG--01 and 09-FEG-04

Rudershausen, P.J., J.A. Buckel, and J.E. Hightower. 2014. Estimating reef fish discard mortality using surface and bottom tagging: effects of hook injury and barotrauma. *Canadian Journal of Fisheries and Aquatic Science*. 71: 514–520.

Sattar, S.A., C. Jorgensen, and O. Fiksen. 2008. Fisheries-induced evolution of energy and sex allocation. *Bulletin of Marine Science*. 83(1): 235-250.

Sedberry. George R. 1988. Food and feeding of black sea bass, *Centropristis striata*, in live bottom habitats in the south Atlantic bight. *The Journal of the Elisha Mitchell Scientific Society*, 104 (2), 1988, pp 35-50.

Shepherd, G. 1991. Meristic and morphometric variation in black sea bass north of Cape Hatteras, North Carolina. *North American Journal of Fisheries Management* 11:139-148.

Shepherd, G.R. and J. Idoine. 1993. Length based analysis of yield and spawning biomass per recruit for black sea bass, *Centropristis striata*, a protogynous hermaphrodite. *Fishery Bulletin*. 91:328-337.

Shepherd, G.R. and M. Terceiro. 1994. The summer flounder, scup and black sea bass fishery of the Middle Atlantic Bight and Southern New England waters. *NOAA Technical Report NMFS 122*.

Shepherd, G., K. Shertzer, J. Coakley, and M. Caldwell (Editors). 2013. Proceedings from a workshop on modeling protogynous hermaphrodite fishes, Raleigh, NC. 33 p. Available from: Mid-Atlantic Fishery Management Council, 800 North State Street, Suite 201, Dover, DE 19901, or online at <http://www.mafmc.org>

Smith, B.E. and J.S Link. 2010. The trophic dynamics of 50 finfish and 2 squid species on the northeast US continental shelf. *NOAA Technical Memorandum NMFS-NE-216(640)*; 646p.

Southeast Data, Assessment and Review (SEDAR). 2011. SEDAR 25 Stock Assessment Report: Atlantic Menhaden. North Charleston, SC. 480 pp.

Southeast Data, Assessment and Review (SEDAR). 2015. SEDAR 40 Stock Assessment Report: South Atlantic Black Sea Bass. North Charleston, SC. 643 pp.

Stari, T., K. Preedy, E. McKenzie, W. Gurney, M. Heath, P. Kunzlik, and D.C. Speirs. 2010. *Fisheries Research*. 105(1):2-12.

Steimle, F.W. and W. Figley. 1996. The Importance of Artificial Reef Epifauna to Black Sea Bass Diets in the Middle Atlantic Bight. *North American Journal of Fisheries Management*. 16:2, 433-439.

Steimle, F.W., C.A. Zetlin, P.L. Berrien, and S. Chang. 1999. Essential Fish Habitat source document: Black Sea Bass, *Centropristis striata*, Life History and Habitat Characteristics. *NOAA Technical Memo*. NMFS-NE-143.

Steimle, F.W., R.A. Pikanowski, D.G. McMillan; C.A. Zetlin and S.J. Wilk. 2000. Demersal fish and American lobster diets in the Lower Hudson-Raritan Estuary. *NOAA Technical Memo*. NMFS-NE-161; 118p.

Stephen, J.A. and P. J. Harris. 2010. Commercial catch composition with discard and immediate release mortality proportions off the southeastern coast of the United States. *Fisheries Research*. 103:18-24

Thorson, J.T., J.M. Cope, T.A. Branch and O.P. Jensen. 2012. Spawning biomass reference points for exploited marine fishes, incorporating taxonomic and body size information. *Canadian Journal of Fisheries and Aquatic Science*. 69: 1556-1568.

Vuong, Q. H. (1989). Likelihood ratio tests for model selection and non-nested hypotheses. *Econometrica*, 57, 307-333.

Wigley, S.E., M.E. Palmer, J. Blaylock, and P.J. Rago. 2008. A brief description of the discard estimation for the National Bycatch Report. *NEFSC Reference Doc.* 08-02.

Zhou, S., S. Yin, J.T. Thorson, A.D.M. Smith and M. Fuller. 2012. Linking fishing mortality reference points to life history traits: an empirical study. *Canadian Journal of Fisheries and Aquatic Science*. 69: 1292-1301.

## TABLES

Table A1. Summary of life history and habitat characteristics for black sea bass, *Centropristis striata*. (YOY = young-of-the-year; SNE = southern New England; MAB = Middle Atlantic Bight; GOM = Gulf of Maine. *In*: Steimle et al. 1999.

Life Stage	Time of Year	Size and Growth	Geographic Location	Habitat	Substrate
<i>Eggs</i>	May-Oct; appear earlier in south and later in north	0.9-1.0 mm; incubation 2-5 days	Coastal MAB; rarely in estuaries	Upper water column, shore to > 200 m depth off Virginia	Buoyant in upper water column
<i>Larvae</i>	May-Nov; peak June-July; appear earlier in south and later in north	Hatch at ~2.1 mm; stage lasts to ~15 mm transition	MAB, near shore, mouths of some estuaries, but rarely in them	As for eggs, < 100 m until transition to juveniles	Upper water column until transition to juveniles
<i>Juveniles YOY</i>	April-Dec; most settle June-Nov	~10-16 mm to 100 mm TL by Nov	MAB into GOM, inshore and into estuaries mid-late summer	Estuarine - coastal; ~1-38 m; salt marsh edges & channels; high habitat fidelity	Rough bottom, shellfish, sponge, and eelgrass beds, nearshore shell patches, man-made objects
<i>Juveniles Winter</i>	Dec-April	~2-12 cm; growth rate reduced	MAB: Most move offshore and south of New Jersey to warmer, deeper waters	Mostly deeper than 38 m; may prefer 90-100 m; mid and outer continental shelf and Chesapeake Bay	Nearshore shell patches and other shelter on sandy bottoms
<i>Adults Summer</i>	April-Dec	> 19 cm FL; growth sexually dimorphic	Coastal: MAB into GOM	~2-38 m; larger fish stay in deeper waters	Mussel beds, rock, artificial reefs, wrecks and other structures
<i>Adults Winter</i>	Nov-March	> 19 cm FL	Most move offshore and south of New Jersey to warmer (> 6°C) waters.	30-240 m depths; mostly 60-150 m mid/outer continental shelf; otherwise poorly known	Poorly known, possibly available shelter on offshore silty sand (e.g., pits)
<i>Spawning Adults</i>	May-Oct, peak in June; begins in the south and progresses north	> 19 cm FL; mature at age 1+	Inshore MAB, south to north, during migration	~20-50 m	Over sand, sand with rock, and reefs



Table A2. Summary of results from a variety of methods used to estimate black sea bass natural mortality.

Method	Equation	L_inf	K	t_max	P 1	P 2	a_50	GI	temp	instantaneous	annual
Alverson and Carney 1975	$M = 3K / (\exp[0.38 * K * t_{max}] - 1)$		X	X						0.42	0.35
Rikhter and Efanov 1977	$M = [1.521 / (a_{50}^{0.720})] - 0.155$						X			0.94	0.61
Gunderson 1980	$M = -0.370 + 4.64GI$							X		0.56	0.43
Pauly 1980 <sup>1</sup>	$M = \exp[-0.0152 + 0.6543 * \ln(K) - 0.279 * \ln(L_{inf}/10) + 0.4634 * \ln(Temp)]$	X	X						X	0.32	0.28
Hoinig 1983 (rule-of-thumb)	$M = -\ln(P) / t_{max}$			X	X					0.35	0.30
Hoinig 1983 (rule-of-thumb)				X		X				0.25	0.22
Roff's 1st 1984	$M = 3 * K / [\exp(t_{max} * K) - 1]$		X	X						0.07	0.07
Ralston 1987 <sup>4</sup> (linear regression)	$M = 0.0189 + 2.06 * K$		X							0.39	0.32
Gunderson and Dygert 1988	$M = 0.03 + 1.68 * GI$							X		0.37	0.31
Charnov & Berrigan 1990	$M = 2.2 / t_{max}$			X						0.18	0.17
Jensen's 1st 1996	$M = 1.65 * t_{max}$			X						0.14	0.13
Jensen 1996 (theoretical)	$M = 1.50 * K$		X							0.27	0.24
Jensen 1996 (derived from Pauly 1980)	$M = 1.60 * K$		X							0.29	0.25
Gunderson 1997	$M = 1.79 * GI$							X		0.36	0.30
Hewitt and Hoinig 2005	$M = 4.22 / t_{max}$			X						0.35	0.30
Updated Tmax estimator (Then et al. 2015)	$M = 5.109 / t_{max}$			X						0.43	0.35
Updated Hoinig 1983 (Then et al. 2015)	$M = \exp[1.717 - 1.01 * \ln(t_{max})]$			X						0.45	0.36
Updated Hoinig Non-linear least squares (Then et al. 2015)	$M = 4.899 * t_{max} - 0.916$			X						0.50	0.40
Updated Pauly Non-linear least squares-T (Then et al. 2015)	$M = 4.118 * K^{0.73} * L_{inf}^{0.33}$	X	X							0.14	0.13
Beverton <sup>5</sup>	$M = 3 * K / [\exp(a_{50} * K) - 1]$		X				X			1.65	0.81

Notes

- 1 Pauly's (1980) equation converted from base 10 to natural logarithms by Quinn and Deriso (1999); L\_inf divided by 10 to adjust for mm
- 2 Values 1.44 (intercept) and -0.982 (slope) were those recommended by Hoinig (1983)
- 3 Equation  $\approx 3 / t_{max}$  when P = 0.05
- 4 Ralston's (1987) relationship was developed for snappers and groupers
- 5 Was not able to verify this equation or identify reference

Table A3. Brief summary of the history of black sea bass stock assessments at NEFSC.

- 1985 SARC 1 – Simple data review. Indices and catch.
- 1987 SARC 5 – Simple data review. Indices and catch.
- 1988 SARC 6 – Simple data review. Indices and catch.
- 1989 SARC 9 – Simple data review. Indices and catch.
- 1990 SARC 11 – length based mortality. Over-exploited.
- 1991 SARC 13 – length based mortality. Over-exploited.
- 1995 SARC 20 – ADAPT VPA accepted. High F (>1.0)
- 1997 SARC 25 – exploratory VPA, general trends accepted. Inadequate data.
- 1998 SARC 27 – not approved. Inadequate data for age based or production model
- 2004 SARC 39 – Tag based R/M, F replacement model. Exploitation from tagging ok.
- 2006 SARC 43 – Tag analysis. Not accepted as valid.
- 2008 Northeast Data Poor Stocks Working Group – Length based model (SCALE) approved for status and BRPs. Use with caution.
- 2009 SSC – Length based model considered too uncertain to use.
- 2010 SSC (a) – Model results reconsidered. Still too uncertain.
- 2010 SSC (b) – Model improved. Still considered unreliable.
- 2011 SARC 53- Not accepted for management. Data deficiencies and spatial issues.
- 2013 SSC – Data update
- 2014 SSC – Data update
- 2015 SSC – Data update

Table A4 . Summary of black sea bass survey indices used in overall and regional models.

Index Name	Assessed			Years	Fishery			Filter Method	Derivation Method	Data	Spatial	
	Area	Season	Location		Independent	Type	Age Range				Range	Rank
NEFSC Winter Bottom Trawl Survey	S	Feb-Mar	Offshore	1992-2007	Yes	Absolute	1-8+	Strata subset	Stratified mean	Len,Wt & Age	Wide	3
NEFSC Spring Bottom Trawl Survey ALBATROSS	N + S	Mar-May	Offshore	1989-2008	Yes	Absolute	1-8+	Strata subset	Stratified mean	Len,Wt & Age	Wide	2
NEFSC Spring Bottom Trawl Survey BIGELOW	N + S	Mar-May	Offshore	2009-2015	Yes	Absolute	1-8+	Strata subset	Stratified mean	Len,Wt & Age	Wide	3
MA Resource Assessment Trawl Survey	N	May	Inshore	1989-2015	Yes	Relative	1-8+	Strata subset	GLM	Length	Narrow	4
RI Seasonal Bottom Trawl Survey	N	Apr-May	Inshore	1989-2015	Yes	Relative	1-8+	All Strata	GLM	Length	Narrow	4
CT Long Island Sound Trawl Survey	N	May-Jun	Inshore	1989-2015	Yes	Relative	1-8+	All Strata	GLM	Length	Narrow	4
NY Small Mesh Trawl Survey	N	May-Jul	Inshore	1990-2015	Yes	Relative	1	Max length	GLM	Length	Narrow	5
NJ Ocean Trawl Survey	S	Jun	Inshore	1989-2015	Yes	Relative	1-8+	All Strata	GLM	Length	Medium	4
DE 16ft Trawl Survey	S	Apr-Jun	Inshore	1989-2015	Yes	Relative	1	Max length	Stratified mean	Length	Narrow	5
MD Coastal Bays 16ft Trawl Survey	S	May-Jun	Inshore	1989-2015	Yes	Relative	1	Max length	Stratified mean	Length	Narrow	5
VIMS Juvenile Fish and Blue Crab Trawl Survey	S	May-Jul	Inshore	1989-2015	Yes	Relative	1	Max length	Stratified mean	Length	Narrow	5
NEAMAP Ocean Trawl Survey	N + S	Apr-May	Coastal	2008-2015	Yes	Relative	1-8+ (N), 1 (S)	All Strata	Stratified mean	Len,Wt & Age	Medium	3
Recreational CPUE	N + S	Mar-Dec	Coastal	1989-2015	No	Relative	1-8+	Multi-Species effort	GLM	Length	Wide	1

Ranked by  
Area coverage  
Time series length  
Ages represented

Table A5. Regional guild species used for calculation of black sea bass effort, based on Jaccard similarity.

<b>Region</b>	<b>Species</b>	<b><i>J</i> value</b>
North	Black sea bass	1
	Scup	0.216
	Summer flounder	0.0826
	Sea robins	0.0749
	Tautog	0.0615
South	Black sea bass	1
	Summer flounder	0.1164
	Sea robins	0.0931
	Atlantic croaker	0.0864
	Weakfish	0.0636
	Tautog	0.0634
	Oyster toadfish	0.0597
	Bluefish	0.0572
	Spot	0.0538
Scup	0.05	
Coast	Black sea bass	1
	Scup	0.1294
	Summer flounder	0.1047
	Sea robins	0.0852
	Atlantic croaker	0.0702
	Tautog	0.064

Table A6. Time series of recreational effort by region. Effort is calculated as the number of intercepted trips with that caught one or more guild species.

<b>Year</b>	<b>North</b>	<b>South</b>	<b>Coast</b>
1989	3491	5668	6585
1990	4168	5200	7272
1991	4626	6294	9069
1992	4563	5551	8621
1993	3989	4212	7313
1994	3323	6122	8016
1995	1788	5000	5599
1996	2446	5002	6135
1997	2550	6227	6998
1998	2711	6551	7649
1999	2879	6025	7453
2000	3297	6433	8055
2001	3935	9233	11070
2002	3800	7912	9738
2003	5089	8822	11740
2004	3580	8123	9799
2005	3308	7952	9463
2006	3471	5890	7688
2007	3557	7846	9428
2008	3169	7922	9338
2009	2551	7031	8132
2010	3060	7093	8470
2011	2667	5642	6942
2012	3316	4799	6834
2013	4226	7238	9712
2014	4261	7188	9345
2015	5442	6417	10046

Table A7. Final models used to fit regional recreational black sea bass fishery indices of abundance.

	<b>Positive</b>	<b>Binary</b>
North (MA-NY)	$\ln\text{CPUE} \sim \text{YEAR} + \text{AREA} + \text{WAVE} + \text{STATE} + \text{HRSF}$	$\text{success} \sim \text{YEAR} + \text{AREA} + \text{WAVE} + \text{STATE} + \text{MODE}$
South (NJ-VA)	$\ln\text{CPUE} \sim \text{YEAR} + \text{AREA} + \text{WAVE} + \text{HRSF}$	$\text{success} \sim \text{YEAR} + \text{AREA} + \text{WAVE} + \text{MODE}$
Coast (MA-VA)	$\ln\text{CPUE} \sim \text{YEAR} + \text{AREA} + \text{WAVE} + \text{STATE} + \text{HRSF}$	$\text{success} \sim \text{YEAR} + \text{AREA} + \text{WAVE} + \text{STATE} + \text{MODE}$

Table A8. Commercial black sea bass landings (mt) by state.

Year	State												Total
	ME	NH	MA	RI	CT	NY	NJ	PA	DE	MD	VA	NC	
1989	1.8	0.0	157.4	94.4	0.0	40.6	381.3	0.0	0.0	134.2	294.7	0.0	1104.5
1990	0.9	0.0	193.6	98.7	0.0	34.3	449.1	0.0	0.0	223.3	401.7	0.0	1401.6
1991	0.0	0.0	109.2	35.3	0.0	45.7	469.1	0.0	85.9	218.2	226.3	0.0	1189.6
1992	0.0	0.0	19.6	64.0	0.0	52.8	564.7	0.0	0.0	300.2	263.1	0.0	1264.3
1993	0.0	0.0	16.5	102.5	0.0	58.5	626.2	0.0	0.0	202.9	346.0	0.0	1352.6
1994	0.0	0.0	7.8	42.4	0.0	55.4	433.4	0.0	0.0	132.4	176.9	0.0	848.4
1995	0.0	0.0	14.6	54.4	5.2	76.3	365.1	0.0	0.0	213.8	159.8	0.0	889.1
1996	0.0	0.0	17.2	71.9	7.6	117.6	551.7	0.0	75.4	248.7	358.2	0.0	1448.4
1997	0.0	0.0	39.5	83.5	6.5	116.7	320.0	0.0	69.2	222.8	239.1	99.6	1197.0
1998	0.0	0.0	125.7	61.6	4.0	62.3	263.9	0.0	40.3	144.6	368.9	80.3	1151.9
1999	0.0	0.0	250.6	83.0	11.6	95.0	227.3	0.0	51.6	199.4	335.6	36.3	1290.3
2000	0.0	0.0	241.0	59.1	32.7	66.6	301.8	0.0	42.6	136.8	259.7	45.8	1186.2
2001	0.2	0.0	232.5	170.0	31.4	117.1	317.6	0.0	11.5	70.0	274.0	54.5	1278.8
2002	0.0	0.0	395.6	155.3	33.3	151.1	310.1	0.0	10.9	90.2	352.6	64.6	1563.7
2003	0.0	0.0	188.1	166.1	10.1	104.4	323.6	3.3	43.6	143.8	208.2	156.1	1347.3
2004	0.0	0.2	130.3	199.4	14.0	143.3	331.7	5.1	24.4	151.1	223.3	182.7	1405.4
2005	0.0	0.0	231.3	128.9	11.4	111.9	201.9	0.0	33.8	185.0	223.9	168.6	1296.7
2006	0.0	0.0	228.2	154.1	10.8	147.9	224.6	0.0	40.1	159.2	147.6	172.9	1285.4
2007	0.0	0.0	144.1	146.7	9.3	189.4	220.2	0.0	29.5	100.7	94.6	102.2	1036.9
2008	0.0	0.0	98.6	83.7	18.2	144.9	192.9	0.0	23.2	96.0	119.3	98.2	875.1
2009	0.0	0.0	46.6	57.0	10.9	75.1	91.7	0.0	20.3	61.7	79.1	80.9	523.2
2010	0.0	0.0	62.0	109.0	44.0	113.1	136.5	0.0	34.8	70.3	134.2	47.4	751.4
2011	0.0	0.0	76.9	84.0	46.1	111.7	131.6	0.0	37.2	85.2	141.6	50.6	765.0
2012	0.0	0.0	41.5	80.2	75.5	105.7	138.1	0.0	37.9	66.9	196.5	39.9	782.1
2013	0.0	0.0	73.9	106.8	46.9	134.9	194.9	0.0	48.7	96.0	253.5	71.5	1027.0
2014	0.0	0.0	62.0	101.1	50.4	161.1	226.9	0.0	46.8	113.7	187.8	138.0	1087.8
2015	0.0	0.0	157.1	72.2	22.2	68.4	249.2	0.0	55.9	131.7	203.1	153.8	1113.4

Table A9. Commercial black sea bass landings (mt) by statistical area.

year	Stat area																				Grand Tota		
	511-534	537	538	539	541-600	611	612	613	614	615	616	621	622	623-624	625	626	627-629	631	632	633-634		635	636-639
1989	5.8	42.3	162.4	25.3	0.0	15.5	9.2	14.1	8.6	2.1	40.4	239.5	183.8	0.0	11.8	210.5	0.0	25.4	107.2	0.0	0.7	0.0	1104.5
1990	6.0	53.2	201.7	17.9	0.0	9.0	12.6	8.5	16.8	3.6	39.2	367.7	193.6	3.2	54.1	289.6	0.0	33.4	91.0	0.0	0.4	0.0	1401.6
1991	1.2	24.8	109.7	10.4	0.0	12.0	24.4	13.0	44.3	5.5	39.4	576.0	69.9	0.8	53.7	114.0	0.0	51.0	38.8	0.0	0.4	0.0	1189.6
1992	2.5	25.5	19.7	17.4	0.0	18.4	25.6	13.8	40.3	4.4	111.5	532.0	100.8	0.0	73.3	192.7	0.0	67.7	17.7	0.0	1.1	0.0	1264.3
1993	2.3	17.7	15.9	29.0	0.0	8.9	31.3	21.8	33.2	2.4	65.2	400.1	287.6	2.9	45.7	338.1	0.3	30.2	19.1	0.0	0.7	0.0	1352.6
1994	5.4	14.0	7.0	15.4	0.3	11.6	20.1	20.7	34.4	5.0	48.3	274.4	103.5	9.0	32.9	195.6	2.6	37.5	8.6	0.1	1.0	0.9	848.4
1995	9.3	18.6	14.2	22.4	0.0	9.6	30.1	29.3	47.0	12.8	47.4	327.6	63.5	9.7	30.0	109.0	15.1	60.1	22.5	0.0	0.0	10.9	889.1
1996	6.6	30.3	17.6	28.2	73.8	19.2	37.6	43.9	45.0	19.8	96.9	298.3	250.8	6.0	89.8	218.5	0.4	83.6	80.4	0.3	1.4	0.0	1448.4
1997	3.5	34.9	40.5	25.2	65.6	24.0	19.3	54.3	34.9	11.7	85.8	289.8	49.3	2.6	69.1	149.6	0.0	81.5	55.0	0.0	100.2	0.0	1197.0
1998	14.5	29.4	114.0	30.2	0.0	13.4	6.3	26.5	31.0	5.8	35.4	225.2	80.4	0.7	111.0	129.8	0.3	63.3	149.6	0.2	82.4	2.4	1151.9
1999	11.3	53.0	243.7	37.0	0.1	23.6	10.7	43.3	33.6	9.3	38.0	283.2	71.5	1.3	81.8	163.9	0.3	88.6	55.6	1.2	35.9	3.3	1290.3
2000	4.3	31.5	228.0	39.6	0.4	43.9	9.4	39.0	29.0	15.9	33.1	186.8	77.7	2.5	63.1	196.8	3.0	84.6	56.3	0.0	41.2	0.2	1186.2
2001	8.7	49.2	226.6	61.8	0.1	50.6	11.2	57.4	30.9	15.1	131.2	79.3	178.1	22.0	60.1	115.3	5.7	58.6	56.5	0.9	59.4	0.1	1278.8
2002	15.9	88.7	355.6	63.7	2.9	58.9	32.9	68.6	50.8	30.8	94.2	104.2	229.1	7.7	74.6	92.5	0.4	89.0	47.3	2.5	53.2	0.0	1563.7
2003	6.1	27.9	178.2	119.0	0.8	63.8	20.2	41.6	44.7	38.2	49.7	219.0	157.6	14.4	68.0	121.5	0.4	35.6	49.0	0.0	91.8	0.0	1347.3
2004	11.1	57.1	140.5	73.8	1.5	47.8	61.5	55.2	13.0	40.1	110.0	239.9	238.2	17.7	48.8	80.5	11.4	50.3	34.7	0.0	65.7	6.6	1405.4
2005	13.7	69.9	194.6	55.1	0.2	49.7	22.5	41.7	13.4	13.1	88.8	246.4	235.0	20.1	28.5	117.5	17.9	31.5	12.8	0.0	20.1	4.2	1296.7
2006	17.8	81.0	178.0	50.5	0.4	55.2	27.5	71.6	45.8	39.4	113.1	298.5	126.2	3.7	14.5	54.4	0.5	48.8	28.7	0.0	8.7	21.2	1285.4
2007	11.9	68.9	123.6	61.6	0.6	42.3	34.6	112.2	24.0	42.8	106.5	208.0	65.5	10.7	9.8	56.0	1.8	32.9	15.8	0.0	3.0	4.4	1036.9
2008	32.9	52.5	68.7	31.6	0.8	37.7	21.8	96.4	23.0	39.0	42.2	141.7	89.9	4.1	17.0	133.0	0.0	24.7	17.6	0.0	0.1	0.5	875.1
2009	2.5	30.1	36.4	21.2	0.2	23.9	24.4	49.1	14.3	9.1	40.6	82.4	90.7	1.3	15.0	26.7	0.6	20.4	16.4	0.0	15.9	1.9	523.2
2010	20.0	31.4	74.8	39.6	0.3	65.8	29.9	72.2	23.3	29.1	63.4	83.9	91.4	7.0	6.7	36.5	0.1	13.9	14.6	0.1	47.1	0.0	751.4
2011	7.9	43.6	53.9	39.8	0.4	83.0	18.8	68.9	10.3	38.3	83.0	102.8	110.5	7.1	16.7	34.3	0.3	17.4	21.1	0.0	6.6	0.2	765.0
2012	8.3	41.0	25.8	56.5	0.3	108.4	17.5	84.8	9.4	62.5	114.2	98.5	50.0	4.2	14.3	28.9	0.3	25.0	26.3	0.0	3.5	2.2	782.1
2013	36.4	80.4	49.8	57.4	1.3	78.4	34.6	107.2	11.6	125.9	124.5	101.4	62.1	10.2	21.6	35.5	0.3	42.8	21.1	0.0	19.3	5.0	1027.0
2014	10.2	59.8	42.4	52.4	0.4	83.9	25.5	131.0	13.7	109.5	273.9	87.3	42.5	9.3	19.0	28.1	0.5	62.3	15.2	0.0	15.3	5.5	1087.8
2015	17.9	84.7	91.3	32.6	0.2	48.9	40.3	122.1	13.4	43.8	313.9	120.6	13.4	6.3	28.1	34.9	0.7	57.7	15.2	0.0	25.7	1.8	1113.4



Table A10. Commercial black sea bass landings (kg) by market category from trawl gear in the northern area for January-June.

Period = QTR 1,2  
 Region = NORTH  
 Gear = TRAWL

Year	LANDINGS (KG)					Total
	Unclassified	Large	Jumbo	Medium	Small	
1989	28675	16843	2615	13158	15228	76519
1990	31993	11187	1889	22243	28837	96149
1991	23455	21387	3614	12184	10748	71388
1992	39865	10060	1424	10649	80440	142437
1993	25095	6888	958	19166	35916	88023
1994	22443	16355	2267	14639	19200	74904
1995	14183	14153	3952	13659	25996	71943
1996	25338	24867	7235	24469	60336	142245
1997	15727	20409	5668	32282	52504	126590
1998	10718	13086	4511	14572	14265	57153
1999	18046	21245	5300	32387	27354	104332
2000	17513	19225	8988	13506	23091	82324
2001	23235	55059	11002	79028	56013	224335
2002	32828	76508	29962	52371	39271	230940
2003	19208	35693	37138	18635	6777	117451
2004	16864	50347	47304	43962	16861	175338
2005	15995	42028	38344	24850	3835	125052
2006	13295	36547	64482	26085	4612	145021
2007	13277	42350	44642	36079	6878	143226
2008	9826	25013	13302	44387	13545	106073
2009	4806	24543	12600	16787	5478	64215
2010	9197	34863	28068	28185	9397	109710
2011	3561	41725	38975	26781	7601	118642
2012	4137	65760	58232	28318	5672	162117
2013	2515	69512	70375	34314	9026	185742
2014	3512	89112	127820	62501	14548	297493
2015	2875	144201	128594	71413	7204	354285

Table A11. Commercial black sea bass landings (kg) by market category from non-trawl gear in the northern area for January-June.

Period = QTR 1,2  
 Region = NORTH  
 Gear = NON-TRAWL

Year	LANDINGS (KG)					Total
	Unclassified	Large	Jumbo	Medium	Small	
1989	40977	39006	3742	16757	254	100735
1990	14561	60892	8987	100025	330	184796
1991	73546	23085	4905	2820	43	104398
1992	14034	5324	2719	556	209	22841
1993	8109	2955	1199	3316	193	15772
1994	16795	7296	876	1948	1587	28502
1995	13921	8101	2573	6420	3480	34494
1996	14967	8468	1577	3145	1295	29451
1997	3968	17115	7556	9140	4967	42746
1998	22309	17577	6511	11962	2475	60833
1999	13569	82579	25576	72089	4337	198151
2000	21497	59956	68466	10768	2168	162855
2001	16869	37322	21291	39018	2033	116533
2002	24886	77985	77735	47373	1876	229855
2003	7650	51805	73897	20039	1019	154411
2004	14894	31986	61037	18601	6437	132955
2005	35408	35588	52978	9085	2125	135183
2006	19939	29875	110783	11842	1791	174230
2007	18320	47816	62619	24695	2768	156217
2008	16736	37765	29146	28159	3117	114922
2009	14935	33949	28058	11364	1064	89369
2010	21377	49585	59496	32408	1427	164292
2011	8490	32951	35370	12686	1569	91067
2012	18164	50168	84517	18438	2301	173588
2013	6405	17636	24244	10059	1838	60182
2014	4339	12832	23063	12877	3689	56800
2015	2044	18415	17473	9310	357	47599

Table A12. Commercial black sea bass landings (kg) by market category from trawl gear in the southern area for January-June.

Period = QTR 1,2  
 Region = SOUTH  
 Gear = TRAWL

Year	LANDINGS (KG)					Total
	Unclassified	Large	Jumbo	Medium	Small	
1989	5387	95133	16086	110267	188775	415647
1990	427	93940	11582	166844	216572	489364
1991	568	43733	7602	54017	69721	175641
1992	618	49306	8186	68824	116077	243011
1993	24770	86183	15274	139992	337853	604071
1994	3984	44096	4194	89568	178520	320362
1995	3409	38983	6077	70474	62572	181515
1996	3036	86229	16495	116683	255777	478219
1997	12108	27728	6742	59246	83684	189507
1998	1050	73864	10067	138553	108068	331601
1999	1330	69687	10919	94119	62228	238284
2000	1059	123069	28741	69245	39544	261658
2001	398	111704	25777	152740	57878	348497
2002	1016	165608	76966	122384	9970	375944
2003	2237	136516	80817	92408	13511	325490
2004	59360	130764	105078	78557	4226	377985
2005	10163	134853	134339	61979	2664	343998
2006	6707	51994	76854	53469	3886	192910
2007	121	47755	30916	33097	386	112275
2008	502	72202	27140	95381	2319	197543
2009	79	55257	34594	23288	973	114191
2010	9	58825	44542	38768	948	143092
2011	4364	67844	45344	27324	327	145202
2012	220	43649	31571	17448	103	92990
2013	122	58143	68872	20806	290	148232
2014		34672	51044	17566	112	103395
2015	26	22820	16655	9647	7	49155

Table A13. Commercial black sea bass landings (kg) by market category from non-trawl gear in the southern area for January-June.

Period = QTR 1,2  
 Region = SOUTH  
 Gear = NON-TRAWL

Year	LANDINGS (KG)					Total
	Unclassified	Large	Jumbo	Medium	Small	
1989	4758	17378	12682	28920	90512	154250
1990	1982	24348	11027	39003	108214	184574
1991	12466	33706	7270	52190	156297	261929
1992	23591	29240	10277	54646	170133	287886
1993	45299	18667	6938	28454	98301	197659
1994	2211	18959	5678	42641	83444	152933
1995	8525	37192	9628	58593	66004	179943
1996	33732	42989	13921	61578	134524	286743
1997	37878	50001	31986	73296	120852	314013
1998	9539	49888	43692	66997	74984	245100
1999	16739	46857	40197	57898	62409	224100
2000	15895	76314	81692	63963	67868	305730
2001	6474	51110	76697	43812	34885	212976
2002	5562	46488	85331	67241	28124	232746
2003	750	58535	60322	102838	21961	244406
2004	2697	50605	43819	57750	19734	174604
2005	17590	41949	27485	76879	13896	177800
2006	956	44836	21196	107307	15824	190119
2007	4101	39411	21235	53864	14629	133240
2008	11655	23333	13851	48486	13456	110780
2009	9398	12551	8431	47742	14637	92759
2010	7334	13572	12431	40766	8482	82585
2011	3728	15035	9187	39667	7908	75527
2012	2942	19911	17795	61425	9554	111627
2013	449	23799	17077	46510	13222	101056
2014	8668	31564	17409	52852	13267	123759
2015	5824	31483	18785	57178	11157	124427

Table A14. Commercial black sea bass landings (kg) by market category from trawl gear in the northern area for July-December.

Period = QTR 3,4  
 Region = NORTH  
 Gear = TRAWL

Year	LANDINGS (KG)					Total
	Unclassified	Large	Jumbo	Medium	Small	
1989	31837	8327	1542	12015	9925	63645
1990	19995	7902	2897	3120	1543	35457
1991	18704	2311	1030	515	3109	25669
1992	21365	2075	706	4334	5521	34001
1993	13656	8472	2295	6678	1266	32366
1994	4422	2103	1154	1776	1034	10489
1995	8349	6680	1520	4958	8247	29753
1996	13012	9710	3561	14536	9204	50024
1997	8473	21576	6739	14545	7147	58481
1998	4868	10106	2474	13656	8307	39410
1999	3539	15938	9174	5975	1978	36604
2000	1601	2351	1098	1720	1622	8391
2001	4489	6233	5493	3622	1560	21397
2002	12754	14922	11502	4702	1081	44961
2003	13130	16089	25211	10401	2328	67158
2004	13569	26363	30154	16522	3522	90130
2005	7409	14998	51921	3731	1048	79108
2006	5143	13357	34230	10016	3620	66366
2007	10753	11809	21229	11249	4869	59908
2008	2395	11525	10233	3786	1579	29519
2009	2330	5553	9729	3081	1808	22501
2010	3009	7081	13126	3856	1203	28274
2011	3476	15867	20985	8569	1379	50276
2012	1621	14095	25331	6843	1644	49534
2013	2393	28593	46541	12932	2779	93237
2014	3300	33283	39559	16765	2797	95704
2015	2502	64011	55270	9507	354	131643

Table A15. Commercial black sea bass landings (kg) by market category from non-trawl gear in the northern area for July-December.

Period = QTR 3,4  
 Region = NORTH  
 Gear = NON-TRAWL

Year	LANDINGS (KG)						Total
	Unclassified	Large	Jumbo	Medium	Small		
1989	58976	5652	821	7945	613	74006	
1990	10818	9817	1702	8906	587	31829	
1991	25875	3995	661	2619	507	33658	
1992	30156	1396	1365	888	1281	35086	
1993	32204	15275	294	6959	1338	56069	
1994	15720	3330	950	1975	6961	28935	
1995	7408	12701	3235	12702	8639	44685	
1996	31637	7237	2522	9601	7595	58593	
1997	13118	18632	5715	15115	7158	59736	
1998	33702	31199	6659	32237	8574	112372	
1999	8238	52410	22704	31590	6729	121669	
2000	24563	50739	15532	69572	15199	175604	
2001	27675	91976	26427	72119	16273	234470	
2002	32137	111361	54513	72834	4871	275716	
2003	9957	58356	50252	32872	16729	168167	
2004	15673	47006	39583	47049	10757	160067	
2005	23996	71914	86556	13466	1038	196969	
2006	23556	56651	75076	46493	7703	209479	
2007	24991	67499	33561	54191	22616	202858	
2008	18641	47841	24993	32481	10023	133980	
2009	4438	17388	11212	14530	4835	52403	
2010	5709	31212	26940	28195	3103	95157	
2011	8977	53890	34426	38396	3521	139210	
2012	5344	20581	29629	14675	1450	71679	
2013	16770	81610	75249	52097	5147	230873	
2014	22797	72582	69026	55785	9382	229572	
2015	27193	90035	69433	31365	313	218339	

Table A16. Commercial black sea bass landings (kg) by market category from trawl gear in the southern area for July-December.

Period = QTR 3,4  
 Region = SOUTH  
 Gear = TRAWL

Year	LANDINGS (KG)					Total
	Unclassified	Large	Jumbo	Medium	Small	
1989	534	3779	552	2780	6588	14233
1990	156	1718	165	1404	2060	5503
1991	265	4701	723	3655	7315	16658
1992	143	7596	1126	8929	21357	39150
1993	219	1921	164	2695	4988	9987
1994	444	4993	888	6609	2060	14995
1995	145	3896	1423	1964	2910	10338
1996	632	4590	1329	8265	18374	33191
1997	222	1198	230	1206	1836	4692
1998	521	1879	793	1533	889	5616
1999	67	1899	1092	916	587	4561
2000	13	362	164	419	285	1243
2001	40	162	109	39	134	485
2002		734	322	442	241	1739
2003	26	20419	24037	4653	58	49194
2004	2001	15612	52285	4674	43	74615
2005	158	5153	12734	456	31	18532
2006	1012	2365	18116	706	74	22273
2007	15	3350	2275	1516	16	7171
2008		7261	10548	948	3	18760
2009		1067	5578	371	16	7031
2010	6	7020	9897	553	8	17483
2011	4	7100	13426	453	8	20990
2012	79	6442	9009	1251	7	16788
2013		30462	30763	7274	60	68559
2014		24402	30158	6029	11	60600
2015		11256	8189	1493		20938

Table A17. Commercial black sea bass landings (kg) by market category from non-trawl gear in the southern area for July-December.

Period = QTR 3,4  
 Region = SOUTH  
 Gear = NON-TRAWL

Year	LANDINGS (KG)					Total
	Unclassified	Large	Jumbo	Medium	Small	
1989	13576	23608	2812	42813	122679	205487
1990	104753	33603	4866	65804	164894	373919
1991	110251	33797	6570	73331	276277	500227
1992	137809	28704	3238	58883	231287	459921
1993	80972	20256	4846	43274	199324	348672
1994	6803	20480	3967	47073	138910	217232
1995	16932	48725	7509	78413	184834	336413
1996	54550	35356	11108	61183	207687	369884
1997	46242	47974	18387	89718	198881	401201
1998	35020	46450	16276	74573	127482	299802
1999	58411	68761	22878	88342	124214	362606
2000	10660	35625	14778	50231	77067	188361
2001	4126	20791	15608	29273	50312	120110
2002	10874	32276	26581	77074	24946	171752
2003	660	55353	24150	124153	16735	221051
2004	3349	49117	20394	110222	36621	219703
2005	13079	29373	14925	138438	24273	220087
2006	20133	58542	17655	140089	48617	285035
2007	18755	54919	17096	106892	24327	221989
2008	14851	24879	14276	81075	28434	163514
2009	5433	9187	6110	47124	12860	80713
2010	7975	12265	6115	72816	11610	110781
2011	13724	16382	6769	78088	9103	124067
2012	4002	22357	11518	54271	11598	103745
2013	2257	30325	12043	80077	14392	139093
2014	4799	22901	7683	71498	13619	120499
2015	49409	28039	10152	66253	13207	167061



Table A18. Summary of black sea bass commercial length measurements by half year and market category, 1989-2015.

Qtr 1,2 year	# lengths					Qtr 3,4 year	# lengths				
	Unclassified	Small	Medium	Large	Jumbo		Unclassified	Small	Medium	Large	Jumbo
1989	202	1767	1341	864	132	1989	38	84	39	48	26
1990	181	3269	2537	1934	260	1990	168	328	150	10	
1991	226	568	452	172		1991	117	130	12	105	67
1992	33	993	827	614	90	1992	37	280	142	31	
1993	76	1927	1816	920	74	1993		56		37	
1994	188	1471	1150	387	2	1994		67	39	42	3
1995	482	562	906	551	98	1995		476	215	151	
1996	24	1077	984	624	107	1996	495	1099	408	491	10
1997	384	1485	1432	830		1997		355	325	150	17
1998		1217	1559	661	152	1998	69	671	362	18	16
1999	221	1157	1268	288	103	1999		752	612	275	37
2000	2	954	1127	802	295	2000	4	621	185		
2001	188	1040	1580	932	50	2001		500	309	127	
2002	128	474	1359	1423	739	2002		303	417	367	269
2003	250	1086	2900	2890	2288	2003	50	99	520	564	368
2004	63	219	3387	2661	2010	2004	219	109	409	212	228
2005	133	318	2991	3208	2348	2005	67	9	853	444	422
2006	445	132	2704	2731	2163	2006	181	250	1639	657	346
2007	174	334	3186	2707	1107	2007	332	299	1548	750	435
2008	271	767	4196	3626	1752	2008	317	807	992	403	497
2009	156	583	2727	2635	1444	2009	183	442	657	307	313
2010	57	440	3623	2501	1861	2010		302	302	274	469
2011	76	260	2880	2603	1305	2011	9	227	1225	879	440
2012	193	743	1993	1173	715	2012	338	578	1375	964	414
2013	320	471	1520	1276	904	2013	204	227	1030	1018	908
2014	282	458	2664	2107	1081	2014	201	328	848	1108	350
2015		400	1124	900	236	2015		256	402	537	83

Table A19. Summary of black sea bass commercial length sample intensity (mt per 100 lengths measured) by market category from January to June, 1989-2015.

year	MT/100 lengths				
	Unclassified	Small	Medium	Large	Jumbo
1989	40	17	13	19	27
1990	27	11	13	10	13
1991	49	42	27	71	0
1992	237	37	16	15	25
1993	136	25	11	12	33
1994	24	19	13	22	0
1995	8	28	16	18	23
1996	321	42	21	26	37
1997	18	18	12	14	0
1998	0	16	15	23	43
1999	22	14	20	77	80
2000	0	14	14	35	64
2001	25	15	20	27	270
2002	50	17	21	26	37
2003	12	4	8	10	11
2004	149	22	6	10	13
2005	60	7	6	8	11
2006	9	20	7	6	13
2007	21	7	5	7	14
2008	14	4	5	4	5
2009	19	4	4	5	6
2010	67	5	4	6	8
2011	27	7	4	6	10
2012	13	2	6	15	27
2013	3	5	7	13	20
2014	6	7	5	8	20
2015	0	5	13	24	77

Table A20. Summary of black sea bass commercial length sample intensity (mt per 100 lengths measured) by market category from July to December, 1989-2015.

year	MT/100 lengths Qtr 3,4				
	Unclassified	Small	Medium	Large	Jumbo
1989	276	166	168	86	22
1990	81	52	53	530	0
1991	133	221	668	43	13
1992	512	93	51	128	0
1993	0	369	0	124	0
1994	0	222	147	74	0
1995	0	43	46	48	0
1996	20	22	23	12	185
1997	0	61	37	60	183
1998	107	22	34	498	164
1999	0	18	21	51	151
2000	0	15	66	0	0
2001	0	14	34	94	0
2002	0	10	37	43	35
2003	48	36	33	27	34
2004	16	47	44	65	62
2005	67	293	18	27	39
2006	28	24	12	20	42
2007	16	17	11	18	17
2008	11	5	12	23	12
2009	7	4	10	11	10
2010	0	5	35	21	12
2011	0	6	10	11	17
2012	3	3	6	7	18
2013	10	10	15	17	18
2014	15	8	18	14	42
2015	0	5	27	36	172

Table A21. Annual black sea bass commercial discard estimates (mt) by region from otter trawl gear, 1989-2015

Year	Otter Trawl Discards			
	NORTH		SOUTH	
	MT	CV	MT	CV
1989	21.5	0.64	61.4	0.40
1990	7.5	0.40	51.3	0.40
1991	2.0	0.64	20.6	0.27
1992	12.7	0.71	84.4	0.42
1993	13.9	0.69	219.7	1.00
1994	5.7	0.68	36.7	0.55
1995	16.3	0.38	21.6	0.65
1996	39.5	0.44	607.2	0.62
1997	75.1	0.51	11.8	1.67
1998	13.1	0.86	150.8	0.43
1999	11.8	0.32	23.9	0.32
2000	13.3	0.65	33.1	0.28
2001	56.4	0.65	144.1	0.32
2002	25.0	1.39	12.0	0.31
2003	67.7	0.59	50.1	0.45
2004	40.2	0.44	375.7	0.25
2005	8.3	0.38	27.3	0.26
2006	12.6	0.44	11.2	0.50
2007	127.8	0.44	29.2	0.51
2008	26.6	0.49	32.8	0.31
2009	70.9	0.48	133.7	0.40
2010	40.6	0.28	96.5	0.17
2011	58.0	0.40	92.3	0.17
2012	43.5	0.43	54.0	0.17
2013	80.7	0.34	119.3	0.19
2014	260.2	0.20	146.0	0.14
2015	128.9	0.31	186.1	0.12

Table A22. Estimated black sea bass commercial discards (mt) by region, gear and season.

	North Total (mt)				South Total (mt)			
	Trawl		Non-trawl		Trawl		Non-trawl	
	1st half	2nd half	1st half	2nd half	1st half	2nd half	1st half	2nd half
1989	18.9	2.6	0.2	0.1	31.2	30.2	0.4	0.4
1990	5.6	1.9	0.1	0.0	23.0	28.2	0.3	0.4
1991	1.4	0.6	0.0	0.0	10.3	10.3	1.1	1.1
1992	9.4	3.4	0.1	0.1	41.4	43.0	0.5	0.6
1993	10.7	3.2	0.1	0.1	92.4	127.3	1.1	1.6
1994	4.5	1.3	0.4	0.1	15.9	20.8	1.2	0.5
1995	11.4	4.9	0.2	0.2	9.6	12.0	0.4	1.5
1996	32.9	6.6	0.1	0.3	278.2	329.0	1.1	1.3
1997	55.7	19.4	0.1	0.3	4.9	7.0	0.7	2.0
1998	7.9	5.2	0.6	1.9	72.2	78.6	2.5	6.8
1999	7.0	4.8	4.3	2.7	11.0	12.9	2.5	5.3
2000	7.4	6.0	5.8	5.1	15.8	17.3	3.1	4.7
2001	33.5	22.9	4.8	12.0	71.5	72.7	3.0	4.1
2002	11.1	13.9	5.2	10.8	5.7	6.4	2.5	3.5
2003	52.0	15.7	1.6	5.8	25.5	24.6	5.0	3.3
2004	24.6	15.6	1.5	2.3	175.4	200.3	2.1	3.9
2005	5.8	2.5	0.5	1.2	12.6	14.7	1.7	2.9
2006	8.9	3.7	1.1	1.4	5.7	5.5	1.8	2.2
2007	106.4	21.4	1.2	1.6	12.7	16.5	0.9	2.9
2008	12.5	14.0	0.7	2.2	14.0	18.8	1.3	2.3
2009	47.1	23.8	1.0	1.0	62.0	71.7	1.1	1.7
2010	21.7	18.9	0.8	2.2	52.4	44.1	0.8	1.4
2011	42.5	15.5	2.6	2.5	44.9	47.4	0.7	1.3
2012	28.7	14.8	2.8	1.5	29.6	24.5	0.9	0.7
2013	32.7	48.1	0.2	8.6	57.1	62.2	1.0	1.5
2014	192.4	67.7	0.3	6.4	70.9	75.1	1.1	1.6
2015	120.5	8.4	0.4	16.7	73.7	112.4	1.4	1.5

Table A22a. Number of black sea bass commercial discard lengths by gear, area and season, 1989-2015.

	Trawl North 1st half	Non-Trawl North 1st half	Trawl North 2nd half	Non-Trawl North 2nd half	Trawl South 1st half	Non-Trawl South 1st half	Trawl South 2nd half	Non-Trawl South 2nd half
<b>1989</b>	96				345			
<b>1990</b>					8			
<b>1991</b>					6			
<b>1992</b>								
<b>1993</b>								
<b>1994</b>			4	4			20	20
<b>1995</b>	8	8	9	9	7	5	70	70
<b>1996</b>	51	51	35	35	379	6	38	4
<b>1997</b>	198	199	62	62	40	42	62	62
<b>1998</b>	61	61	61	61	393	394	3	61
<b>1999</b>	12	12	33	33	12	12	12	33
<b>2000</b>	23	26	92	92	54	54	5	92
<b>2001</b>	102	102	38	38	27	27	53	53
<b>2002</b>	3	3	87	87	49	54	8	8
<b>2003</b>	129	129	124	44	635	53	37	37
<b>2004</b>	609	614	236	236	1107	190	54	54
<b>2005</b>	149	149	58	11	596	603	11	11
<b>2006</b>	148	148	94	9	277	277	32	32
<b>2007</b>	488	109	52	143	198	12	4	4
<b>2008</b>	102	47	620	190	794	22	18	18
<b>2009</b>	581	584	166	18	605	40	17	17
<b>2010</b>	1079	8	473	473	627	75	8	8
<b>2011</b>	635	31	144	144	716	136	8	8
<b>2012</b>	717	9	653	653	196	54	65	65
<b>2013</b>	1264	30	463	18	600	223	136	1
<b>2014</b>	2153	229	621	207	355	119	215	215
<b>2015</b>	1738	64	1581	960	137	98	540	540

Table A22b. Data source for black sea bass commercial length substitutions by gear, area and season, 1989-2015.

	Trawl North 1st half	Non-Trawl North 1st half	Trawl North 2nd half	Non-Trawl North 2nd half	Trawl South 1st half	Non-Trawl South 1st half	Trawl South 2nd half	Non-Trawl South 2nd half
1989								
1990								
1991								
1992								
1993								
1994								
1995								
1996								
1997								
1998								
1999								
2000								
2001								
2002								
2003								
2004								
2005								
2006								
2007								
2008								
2009								
2010								
2011								
2012								
2013								
2014								
2015								




Table A23. Recreational black sea bass landings (AB1) number (000s) (note: does not include offshore wave 1 landings).

	AB1 #										
	ME	NH	MA	RI	CT	NY	NJ	DE	MD	VA	NC
1989	0.0	0.0	3.6	2.0	2.8	65.8	220.8	20.6	40.9	65.5	0.0
1990	0.0	0.0	3.1	0.8	0.1	35.7	150.5	11.3	49.4	134.4	2.8
1991	0.0	0.0	2.5	1.0	0.2	19.8	248.6	39.2	64.1	144.6	2.4
1992	0.0	0.0	0.6	1.2	0.8	16.4	157.9	19.6	75.8	78.3	2.2
1993	0.0	0.0	1.1	2.6	1.0	21.9	421.1	23.7	59.3	67.2	1.2
1994	0.0	0.0	0.7	1.7	0.0	21.8	191.3	6.7	27.3	91.3	1.3
1995	0.0	0.0	0.8	4.2	0.5	9.0	395.2	20.9	149.8	91.9	1.3
1996	0.0	0.0	1.7	4.3	0.5	7.9	232.0	5.9	28.7	79.9	0.9
1997	0.0	0.0	1.6	3.5	0.2	21.7	335.2	9.1	37.2	63.5	1.6
1998	0.0	0.0	0.7	2.6	0.3	1.2	27.3	5.2	35.4	39.8	2.0
1999	0.0	0.0	2.1	2.5	0.2	8.9	44.9	4.1	15.9	53.6	3.2
2000	0.0	0.0	6.6	19.7	1.5	33.5	196.1	15.1	43.3	44.8	3.6
2001	0.0	0.3	5.9	12.3	0.9	16.4	191.8	20.3	11.9	23.1	1.3
2002	0.0	0.0	18.4	7.8	1.8	22.1	175.9	60.7	33.7	13.2	3.4
2003	0.0	0.0	11.7	7.0	0.5	31.7	190.3	30.7	24.1	26.5	3.4
2004	0.0	0.0	15.8	2.7	2.6	13.3	107.8	4.4	1.6	4.6	0.7
2005	0.0	0.0	17.5	8.5	0.0	14.3	66.0	6.8	9.1	3.4	1.4
2006	0.0	0.0	10.5	4.1	0.3	26.8	53.1	11.4	12.1	8.3	1.8
2007	0.0	0.0	14.9	4.4	2.4	41.0	72.4	9.3	3.9	3.6	0.7
2008	0.0	0.0	24.6	5.2	6.0	25.9	57.9	2.3	2.6	3.8	1.7
2009	0.0	0.0	43.1	3.6	0.0	56.6	58.3	3.7	3.3	11.5	1.1
2010	0.0	0.0	70.2	16.0	1.6	54.3	68.7	2.1	3.6	3.0	1.1
2011	0.0	0.0	19.5	5.0	0.8	27.4	14.8	4.3	4.7	1.9	3.1
2012	0.0	0.3	52.0	10.3	11.1	32.1	73.5	4.0	3.3	0.4	0.4
2013	0.0	1.2	29.2	7.5	10.8	35.3	34.5	3.7	3.0	2.1	0.8
2014	0.0	0.0	45.7	21.4	40.7	42.3	46.8	2.4	6.8	1.4	0.1
2015	0.0	0.0	34.2	23.4	33.1	87.6	31.0	2.3	5.8	3.9	0.2



Table A24. Recreational black sea bass landings (AB1) and discard losses (B2\*0.15) number and total weight, by region.

	AB1 # (000s)				Total AB1 wt (MT)	B2 # (000s)				Total B2 loss wt (MT)
	North	South	Total			North	South	Total		
1989	742.0	3479.0	4221.1		1193.0	36.1	283.2	319.3		45.9
1990	396.8	3484.2	3881.1		1159.5	11.2	777.2	788.3		124.5
1991	233.9	4990.2	5224.1		1621.7	42.2	799.4	841.5		108.9
1992	189.7	3339.5	3529.2		1133.3	19.1	626.5	645.7		96.3
1993	265.8	5727.8	5993.6		1735.5	7.4	476.0	483.4		69.5
1994	241.6	3179.9	3421.5		1128.4	50.3	545.2	595.5		116.8
1995	145.4	6594.3	6739.7		2119.4	61.4	1072.0	1133.4		233.9
1996	144.4	3478.7	3623.1		1541.2	65.0	617.8	682.9		155.1
1997	270.5	4476.8	4747.4		1730.5	90.9	811.4	902.3		161.5
1998	52.7	1100.7	1153.4		473.4	24.4	560.9	585.3		178.7
1999	142.3	1223.7	1366.0		729.3	158.0	705.1	863.1		261.0
2000	615.7	3041.4	3657.0		1702.0	297.8	1684.9	1982.7		540.9
2001	360.2	2508.7	2868.9		1522.1	142.8	1493.6	1636.5		595.4
2002	506.1	2890.4	3396.4		1952.6	333.1	1366.2	1699.2		771.1
2003	515.2	2766.2	3281.4		1890.9	179.8	1152.9	1332.8		401.2
2004	346.8	1205.0	1551.8		871.2	113.7	743.7	857.4		201.1
2005	404.9	883.5	1288.4		845.7	200.3	667.7	868.0		193.1
2006	428.6	892.5	1321.0		801.6	262.6	686.3	949.0		202.6
2007	638.2	924.0	1562.2		947.2	326.5	757.0	1083.4		200.5
2008	626.0	713.0	1339.0		908.6	357.8	1049.2	1407.0		257.0
2009	1049.9	802.2	1852.1		1158.7	340.6	777.7	1118.3		240.6
2010	1421.6	785.1	2206.6		1420.5	361.6	804.3	1165.9		272.6
2011	527.8	288.8	816.6		507.4	231.3	804.0	1035.3		232.2
2012	1059.1	816.0	1875.1		1479.5	805.1	801.7	1606.8		364.4
2013	885.3	467.6	1352.9		1198.5	729.5	576.8	1306.3		296.0
2014	1537.7	580.2	2118.0		1745.1	775.2	538.2	1313.3		382.4
2015	1783.4	431.5	2215.0		1863.6	655.9	426.7	1082.6		371.5

Table A25. Review of discard mortality estimates for hook and line fisheries among species comparable to black sea bass.

Species	Recreational Discard		Basis
	Mortality Estimate	Assessment	
Scup	0.15	60th SAW Assesment Report (2015)	Howell and Simpson 1985
Tautog	0.025	ASMFC 2015 Benchmark Peer Review	Simpson and Gates 1999, CT DEP white paper
Spotted Sea Trout	0.08	State assessments (FL FWC)	Gearhart 2002, Murphy et al 1995, Matlock et al 1993
Weakfish	0.1	ASMFC 2016 Benchmark	Murphy et al 1995, Malchoff and Heins 1997, Swihart et al 2000, Duffy 2002, Gearhart 2002
Black Drum	0.08	ASMFC 2015 Assessment	SEDAR 18 (2009) for Red Drum
Atlantic Croaker	0.1	ASMFC 2010 Benchmark	unsupported
Red Drum	0.08	SEDAR 18 (2009)	Vecchio and Wenner 2008, Gearhart 2002, Aguilar 2003
Bluefish	0.15	60th SAW Assesment Report (2015)	Lit. review (working paper B1)
Vermillion Snapper	0.15	not considered in SEDAR 45 (2016)	Guccione 2005 - shallow depths
Gray Triggerfish	0.05	SEDAR 43 (2015)	
Gray Triggerfish	0.125	SEDAR 41 (2014)	Thorough species specific literature review
Red Grouper (GofMex)	0.116	SEDAR 42 (2015)	FL FWC work: Sauls et al. 2014
Atlantic Striped Bass	0.09	57th SAW (2013)	Diodati and Richards (1996) in addition to broader literature review
Black Grouper	0.2	SEDAR 19 (2010)	lacking empirical data, infered from similar species
Red Grouper (SA)	0.2	SEDAR 19 (2010)	Literature review indicated 0-15%, potentially related to depth of capture
Black Sea Bass	0.07	SEDAR 25 (2011)	Literature review, Rudershausen et al 2010 (4.3% from 29-37m depth)
	0.047		Bugley and Shepherd, 1991 (6-12m depths)
	0.13		Collins et al., 1999 (for depths 20-35m)
	0.036		Rudershausen et al., 2007 (immediate mortality from capture depths averageing 28.5m, ranging from 19.1-71.2m)
	0.043		Rudershausen et al., 2010 (ranged from 0 - 6.9% across 6 trips from 29-37m depth)
	0.19		Rudershausen et al., 2014 (29-34m depth)

Table A26. Recreational black sea bass discard (B2) mortalities (000s) assuming 15% discard mortality (note: does not include offshore wave 1 landings).

	ME	NH	MA	RI	CT	NY	NJ	DE	MD	VA	NC
1989	0	0	3	3	3	28	88	22	66	107	0
1990	0	0	2	1	0	8	92	19	219	433	14
1991	0	0	1	1	6	34	303	41	192	243	20
1992	0	0	0	1	0	17	160	33	164	231	39
1993	0	0	1	1	0	4	143	27	82	188	36
1994	0	0	1	0	0	49	198	31	57	207	53
1995	0	0	2	6	1	51	386	47	275	302	63
1996	0	0	3	12	3	47	211	18	87	290	11
1997	0	0	4	5	3	78	422	41	88	230	31
1998	0	0	6	4	2	12	185	43	113	200	20
1999	0	0	4	18	26	110	259	32	223	186	4
2000	0	0	40	60	13	184	832	123	484	236	10
2001	0	0	21	23	3	96	657	151	349	328	8
2002	0	0	40	36	44	212	649	185	139	367	27
2003	0	0	30	31	8	111	645	125	116	261	6
2004	0	0	12	4	5	94	469	42	39	187	7
2005	0	0	22	10	8	161	359	41	84	168	15
2006	0	0	26	24	13	199	313	49	97	204	23
2007	0	0	60	18	15	233	365	88	87	191	27
2008	0	0	61	19	29	249	666	70	101	188	24
2009	0	0	104	20	31	186	473	44	68	174	18
2010	0	0	140	32	14	175	582	35	101	79	10
2011	0	0	101	58	12	236	344	56	93	117	18
2012	0	1	162	115	152	372	575	31	44	133	21
2013	0	1	213	104	197	210	391	38	54	91	8
2014	0	0	224	131	214	198	343	35	76	86	5
2015	0	0	129	111	73	341	337	23	22	40	7

Table A27. Black sea bass age samples, by region, season and source, available for development of age length keys.

	Commercial				NEFSC Survey			
	NORTH 1	SOUTH 1	NORTH 2	SOUTH 2	NORTH 1	SOUTH 1	NORTH 2	SOUTH 2
1989	25					106	72	72
1990		76			34	97	119	110
1991					18	81	97	97
1992					4	148	56	52
1993					8	59	47	46
1994		70			12	53	45	45
1995	15	113	75	133	18	50	90	89
1996	51	417	179	520	8	59	37	37
1997					22		14	
1998					10	55	100	95
1999					35	61	101	71
2000					68	31	54	
2001					75	89	106	76
2002					69	113	192	117
2003					51	147	63	
2004					45	130	70	39
2005					48	86	99	78
2006					24	53	102	70
2007					82	33	97	69
2008	362	237	237	184	111	47	141	83
2009	229	271	321	22	141	113	208	155
2010	84	270	133	78	192	259	247	251
2011	501	661	355	510	66	133	192	194
2012	538	680	616	646	347	157	297	320
2013	764	446	505	119	214	400	283	306
2014	905	252	575	251	352	125	358	307
2015	659	179	180		49	116	321	236

Table A28. Key to application of black sea bass age length data for expansion of commercial and recreational length frequencies.

				1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
North	Commercial	Landings	Trawl	Spring	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
North	Commercial	Landings	Trawl	Fall	A	A	A	A	A	A	A	D	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
North	Commercial	Landings	Non-Trawl	Spring	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
North	Commercial	Landings	Non-Trawl	Fall	A	A	A	A	A	A	A	D	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
North	Commercial	Discards	Trawl	Spring	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
North	Commercial	Discards	Trawl	Fall	A	A	A	A	A	A	A	D	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
North	Commercial	Discards	Non-Trawl	Spring	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
North	Commercial	Discards	Non-Trawl	Fall	A	A	A	A	A	A	A	D	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
North	Recreational	Landings		Spring	A	A	A	B	B	B	B	A	A	A	B	B	B	B	B	B	B	B	C	C	C	B	C	C	C	C
North	Recreational	Landings		Fall	B	A	A	A	A	A	A	A	A	A	B	B	C	B	B	B	B	C	C	C	B	C	C	C	C	C
North	Recreational	Discards		Spring	A	A	A	B	B	B	B	A	A	A	B	B	B	B	B	B	B	B	C	C	C	B	C	C	C	C
North	Recreational	Discards		Fall	B	A	A	A	A	A	A	A	A	A	B	B	C	B	B	B	B	C	C	C	B	C	C	C	C	C
South	Commercial	Landings	Trawl	Spring	A	A	A	A	A	A	D	D	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
South	Commercial	Landings	Trawl	Fall	A	A	A	A	A	A	D	D	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
South	Commercial	Landings	Non-Trawl	Spring	A	A	A	A	A	A	D	D	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
South	Commercial	Landings	Non-Trawl	Fall	A	A	A	A	A	A	D	D	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
South	Commercial	Discards	Trawl	Spring	A	A	A	A	A	A	D	D	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
South	Commercial	Discards	Trawl	Fall	A	A	A	A	A	A	D	D	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
South	Commercial	Discards	Non-Trawl	Spring	A	A	A	A	A	A	D	D	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
South	Commercial	Discards	Non-Trawl	Fall	A	A	A	A	A	A	D	D	A	A	A	A	A	A	A	A	A	A	D	D	D	D	D	D	D	D
South	Recreational	Landings		Spring	C	B	B	C	B	B	A	A	A	A	B	B	C	C	C	C	B	B	B	B	C	C	C	C	C	C
South	Recreational	Landings		Fall	B	C	B	B	B	B	D	B	B	B	B	B	C	B	B	A	B	B	C	C	C	C	B	B	A	C
South	Recreational	Discards		Spring	C	B	B	C	B	B	A	A	A	A	B	B	C	C	C	B	B	B	B	B	C	C	C	C	C	C
South	Recreational	Discards		Fall	B	C	B	B	B	B	D	B	B	B	B	B	C	B	B	A	B	B	C	C	C	C	B	B	A	C

Key	
A	Master
B	NEFSC ± 2 years
C	NEFSC Survey
D	Commercial Key
E	Commercial ± 2 years

Table A29. Black sea bass catch number (000s) at age from trawl gear in the northern region, 1989-2015.

000s Year	Age1	Age2	Age3	Age4	Age5	Age6	Age7	Age8+	total
1989	56.1	296.8	159.8	57.3	17.1	5.3	1.4	0.4	594.8
1990	44.7	171.9	178.9	45.8	10.7	5.2	1.9	0.7	460.2
1991	44.0	86.4	112.2	43.3	9.6	2.8	0.8	0.3	299.5
1992	54.9	417.0	335.2	42.0	7.9	2.0	0.4	0.1	860.2
1993	32.3	223.6	200.1	42.5	8.0	2.7	1.0	0.4	511.2
1994	19.9	120.8	140.3	32.9	6.4	2.2	0.6	0.2	323.8
1995	86.9	118.4	185.1	39.9	6.7	1.6	0.3	0.0	443.5
1996	71.1	392.0	304.6	58.6	12.6	5.1	1.3	0.3	845.7
1997	84.7	438.1	371.0	82.0	15.1	3.7	0.8	0.2	995.6
1998	24.6	71.2	125.6	41.6	10.5	3.8	1.2	0.6	279.2
1999	7.8	38.8	180.3	72.0	17.4	5.6	1.5	0.5	323.9
2000	3.6	21.0	103.0	48.0	12.8	5.1	1.4	0.4	195.3
2001	24.1	108.0	353.0	156.9	31.2	9.4	2.8	1.5	687.5
2002	7.3	82.2	284.8	153.2	41.4	14.5	3.7	0.9	606.7
2003	11.1	235.5	159.2	88.9	37.4	20.3	7.9	2.9	563.4
2004	9.5	39.9	215.9	143.3	48.2	21.9	7.8	3.1	489.7
2005	0.5	12.8	98.3	87.3	38.4	20.1	8.9	4.7	271.2
2006	2.6	26.5	115.2	82.4	36.7	22.4	10.0	5.6	301.5
2007	28.4	331.9	342.6	113.0	34.0	17.3	7.9	5.6	880.8
2008	1.1	21.6	261.5	41.1	5.5	2.0	3.9	2.6	339.3
2009	3.4	37.2	195.3	71.4	15.7	2.8	1.6	3.2	331.0
2010	0.0	48.1	124.8	75.5	30.5	10.5	3.3	1.4	294.2
2011	7.7	33.5	132.3	142.1	41.3	20.4	4.5	1.1	382.9
2012	0.2	88.4	159.4	85.0	55.0	26.6	9.6	2.0	427.8
2013	0.3	185.5	127.0	125.6	72.8	48.2	11.9	7.3	579.0
2014	0.0	35.9	806.7	141.1	98.6	52.4	29.9	9.5	1174.8
2015	0.9	32.1	167.6	476.8	86.2	44.8	14.8	19.6	843.1

Table A30. Black sea bass catch number (000s) at age from non-trawl gear in the northern region, 1989-2015.

Year	Age1	Age2	Age3	Age4	Age5	Age6	Age7	Age8+	total
1989	27.4	405.6	474.8	146.0	40.3	36.5	2.6	0.6	1133.8
1990	44.1	212.8	456.3	160.0	28.2	8.7	2.7	1.3	914.3
1991	100.1	211.5	211.4	86.6	20.2	5.8	1.5	0.5	639.9
1992	46.5	148.0	83.6	42.0	24.0	15.1	5.7	5.4	371.1
1993	43.1	202.5	123.0	42.2	10.4	3.3	1.3	0.5	431.8
1994	161.9	177.3	103.2	25.6	5.1	1.5	0.4	0.1	477.5
1995	106.0	138.0	135.6	38.9	8.1	2.0	0.4	0.0	429.4
1996	67.7	184.4	105.1	40.4	11.7	4.6	1.2	0.3	415.5
1997	121.8	222.1	147.4	69.4	17.3	5.0	1.2	0.3	584.8
1998	44.5	117.3	170.1	81.2	24.3	7.7	1.9	0.5	447.7
1999	116.6	180.3	334.2	181.9	49.9	21.8	4.6	1.5	890.7
2000	145.2	477.9	529.2	255.4	57.7	27.6	8.0	2.7	1504.2
2001	66.6	345.4	521.0	194.0	43.8	31.8	4.1	1.3	1207.9
2002	94.7	363.7	609.0	460.6	164.0	54.4	8.7	2.6	1757.6
2003	71.6	299.3	425.3	269.1	79.1	37.5	11.0	4.0	1196.9
2004	22.1	204.5	388.4	211.9	64.9	27.6	9.6	3.3	932.9
2005	50.7	195.9	307.2	274.4	98.4	64.9	22.2	7.0	1023.6
2006	129.5	276.0	288.5	235.8	162.0	54.6	32.9	22.5	1207.1
2007	65.6	531.4	454.3	330.2	54.1	46.9	11.4	21.8	1517.1
2008	112.4	327.0	687.7	244.5	12.0	5.1	17.0	13.1	1420.4
2009	62.7	407.4	618.8	411.4	79.4	15.6	4.4	21.4	1623.2
2010	44.6	409.4	946.1	435.4	238.1	64.1	22.9	13.0	2185.7
2011	39.4	276.5	359.6	330.6	88.3	33.7	5.6	2.2	1136.2
2012	237.2	473.9	558.2	496.7	253.9	128.2	24.5	13.6	2196.4
2013	78.8	654.4	383.8	359.5	267.1	143.1	92.6	59.7	2040.2
2014	40.9	281.1	1460.7	449.9	180.2	148.7	92.8	72.7	2729.9
2015	12.4	180.7	492.2	1363.9	404.1	177.4	122.5	38.6	2792.1

Table A31. Black sea bass catch number (000s) at age from trawl gear in the southern region, 1989-2015.

Year	Age1	Age2	Age3	Age4	Age5	Age6	Age7	Age8+	total
1989	448.1	852.4	569.6	259.9	90.0	26.2	7.1	3.5	2437.8
1990	160.6	845.1	801.9	337.4	93.6	20.3	4.4	2.1	2270.0
1991	76.8	358.7	229.9	118.5	44.0	11.5	2.6	1.3	845.1
1992	282.5	626.2	489.6	197.2	58.3	13.7	2.8	1.0	1678.3
1993	736.3	1790.0	1123.9	412.2	111.4	24.0	5.3	2.3	4226.2
1994	161.2	635.7	557.8	204.8	54.7	12.6	2.8	1.3	1630.9
1995	129.3	256.0	284.3	108.2	21.3	4.8	0.0	0.0	815.7
1996	2175.1	3213.7	1012.9	228.8	49.4	47.7	7.1	7.1	6741.8
1997	10.9	185.9	298.0	143.4	41.3	8.8	1.7	0.7	690.8
1998	216.9	453.0	593.7	319.3	98.9	22.2	4.2	1.4	1709.6
1999	13.9	78.6	266.3	201.9	73.6	19.7	4.6	1.9	660.5
2000	60.1	50.9	170.5	190.4	96.6	33.1	9.2	4.3	615.1
2001	182.7	312.4	386.6	364.8	145.6	36.8	8.1	3.4	1476.8
2002	21.3	35.4	137.8	250.0	143.2	52.5	14.8	6.5	683.9
2003	27.7	44.0	141.9	242.2	144.9	59.0	19.1	10.3	689.9
2004	483.7	287.4	368.9	362.0	219.7	110.6	41.1	25.0	1909.3
2005	15.2	27.0	96.2	178.4	121.1	58.0	24.2	17.7	538.0
2006	2.7	5.7	61.0	103.7	65.2	32.8	14.5	11.4	297.1
2007	39.1	57.9	74.0	83.9	42.0	15.1	5.5	4.6	322.2
2008	0.1	194.4	155.6	120.0	42.3	26.2	15.1	6.1	559.8
2009	234.9	199.2	141.7	139.5	28.3	17.5	5.7	11.3	780.3
2010	0.3	103.9	223.8	130.3	77.4	19.3	3.4	4.4	565.3
2011	0.2	275.7	100.7	187.1	82.2	24.8	4.2	3.6	678.6
2012	109.5	13.2	173.9	75.2	70.7	21.4	7.2	5.2	476.3
2013	0.0	55.2	155.8	168.0	107.4	63.8	44.0	5.8	600.0
2014	0.0	34.7	499.8	78.7	49.3	43.1	27.0	2.3	735.0
2015	584.3	271.9	109.0	78.5	38.1	18.6	7.4	4.9	1139.1



Table A32. Black sea bass catch number (000s) at age from non-trawl gear in the southern region, 1989-2015.

Year	Age1	Age2	Age3	Age4	Age5	Age6	Age7	Age8+	total
1989	328.8	2845.8	1216.4	457.8	79.1	38.7	4.9	5.8	5004.5
1990	714.1	2499.6	1912.3	666.2	211.2	49.7	8.8	12.4	6274.2
1991	1271.1	4230.7	2213.8	864.9	206.0	83.5	5.5	81.5	9020.5
1992	730.8	3175.5	1889.9	678.8	187.0	42.6	8.9	6.3	6724.7
1993	643.4	3809.3	2637.1	771.9	192.2	38.5	5.8	21.5	8170.9
1994	758.9	1495.5	2288.2	457.0	109.4	28.9	3.2	1.6	5171.0
1995	2244.3	3963.7	2040.2	620.3	72.6	16.2	1.4	384.2	9437.5
1996	805.5	2950.4	2030.8	476.0	98.9	55.4	8.3	10.6	6452.7
1997	966.9	2789.8	1950.7	1131.2	367.9	77.0	14.8	179.1	7516.2
1998	193.7	865.8	1049.0	709.4	241.7	60.5	13.8	8.3	3143.3
1999	195.9	694.9	1133.8	781.6	368.5	98.0	17.7	11.4	3313.5
2000	612.1	1604.8	2017.0	1163.8	243.1	91.8	34.2	11.7	5782.3
2001	104.7	1598.7	1740.1	741.4	209.2	229.3	31.2	13.1	4668.1
2002	91.2	578.5	2677.4	1321.1	208.7	75.5	64.6	25.6	5043.8
2003	271.1	701.1	1883.2	1372.1	479.6	108.3	30.9	7.3	4854.3
2004	179.8	433.8	1477.6	357.4	245.7	43.9	11.0	9.0	2763.3
2005	120.4	607.7	624.8	608.6	342.4	53.0	11.4	6.2	2383.1
2006	216.4	380.5	719.4	679.4	478.7	78.1	13.8	7.2	2584.6
2007	315.7	425.2	596.9	561.8	239.0	162.4	55.1	5.7	2375.4
2008	484.5	821.5	650.4	206.2	99.8	47.3	22.8	15.0	2378.1
2009	205.4	344.5	720.7	490.7	129.7	47.8	13.1	16.0	1981.0
2010	215.8	308.7	753.4	577.0	105.4	30.0	2.0	17.4	2018.8
2011	252.6	265.4	381.5	426.8	180.3	24.9	3.3	1.0	1547.1
2012	315.1	212.7	488.9	446.3	511.3	35.4	12.8	6.8	2071.8
2013	232.3	186.7	294.1	403.6	259.7	47.2	86.2	1.4	1526.7
2014	122.8	229.3	677.4	169.6	332.5	42.6	11.9	1.1	1595.1
2015	126.1	182.3	279.0	579.7	133.2	60.1	11.2	3.9	1384.7

Table A33. Black sea bass mean weight at age (kg) for the Northern region.

Year	Age 1	Age2	Age3	Age4	Age5	Age6	Age7	Age8+
1989	0.101	0.178	0.399	0.688	0.999	1.254	1.471	1.763
1990	0.157	0.225	0.369	0.560	0.844	1.293	1.593	2.236
1991	0.138	0.200	0.361	0.617	0.845	1.145	1.473	2.031
1992	0.144	0.175	0.276	0.716	1.208	1.569	1.775	2.095
1993	0.140	0.199	0.341	0.580	0.940	1.329	1.695	1.969
1994	0.124	0.185	0.321	0.542	0.858	1.193	1.514	1.944
1995	0.119	0.215	0.329	0.556	0.813	1.043	1.235	1.495
1996	0.140	0.231	0.357	0.617	0.964	1.223	1.397	1.837
1997	0.158	0.180	0.333	0.597	0.849	1.116	1.388	1.980
1998	0.188	0.269	0.390	0.641	0.929	1.203	1.545	2.199
1999	0.125	0.276	0.435	0.648	0.896	1.250	1.513	1.967
2000	0.132	0.329	0.521	0.786	0.976	1.321	1.553	1.947
2001	0.134	0.297	0.464	0.644	0.869	1.372	1.595	2.133
2002	0.118	0.319	0.476	0.819	1.099	1.525	1.475	1.950
2003	0.124	0.253	0.486	0.782	1.106	1.426	1.655	1.953
2004	0.171	0.349	0.473	0.719	1.023	1.397	1.676	2.011
2005	0.104	0.316	0.489	0.766	1.128	1.514	1.829	2.212
2006	0.125	0.312	0.455	0.680	0.999	1.442	1.814	2.691
2007	0.131	0.273	0.403	0.651	1.009	1.318	1.869	2.410
2008	0.140	0.279	0.509	0.920	1.048	1.595	1.737	2.227
2009	0.117	0.283	0.455	0.763	1.042	1.195	1.314	2.432
2010	0.083	0.358	0.536	0.772	0.956	1.272	1.334	1.767
2011	0.112	0.296	0.467	0.670	0.976	1.353	1.523	2.111
2012	0.102	0.319	0.514	0.753	1.156	1.293	1.715	2.304
2013	0.108	0.243	0.539	0.698	1.034	1.373	1.537	2.135
2014	0.112	0.256	0.451	0.819	1.086	1.559	1.753	2.299
2015	0.106	0.233	0.455	0.697	1.025	1.276	1.691	1.967

Table A34. Black sea bass mean weight at age (kg) from the southern region.

Year	Age 1	Age2	Age3	Age4	Age5	Age6	Age7	Age8+
1989	0.066	0.173	0.307	0.483	0.582	0.843	1.189	1.516
1990	0.097	0.180	0.306	0.428	0.540	0.874	1.108	1.516
1991	0.123	0.168	0.314	0.501	0.621	1.018	1.080	1.954
1992	0.126	0.208	0.290	0.432	0.637	0.840	1.077	1.317
1993	0.114	0.193	0.300	0.398	0.544	0.872	1.173	1.123
1994	0.113	0.204	0.347	0.397	0.554	0.872	1.127	1.456
1995	0.123	0.229	0.371	0.574	0.702	1.029	1.060	1.031
1996	0.116	0.181	0.450	0.462	0.669	0.860	0.925	1.637
1997	0.143	0.222	0.354	0.469	0.593	0.852	1.079	1.043
1998	0.150	0.219	0.319	0.458	0.612	0.851	1.118	1.304
1999	0.143	0.250	0.377	0.573	0.626	0.921	1.137	1.322
2000	0.122	0.242	0.385	0.604	0.690	1.063	1.306	1.496
2001	0.114	0.278	0.412	0.562	0.676	1.104	1.389	1.523
2002	0.106	0.266	0.404	0.649	0.748	1.020	1.399	1.421
2003	0.107	0.279	0.438	0.583	0.734	1.015	1.347	1.561
2004	0.109	0.217	0.434	0.521	0.723	1.078	1.381	1.576
2005	0.111	0.295	0.376	0.532	0.776	1.027	1.406	1.742
2006	0.097	0.269	0.387	0.510	0.616	0.954	1.356	1.710
2007	0.099	0.224	0.399	0.537	0.653	1.019	1.083	1.904
2008	0.105	0.275	0.462	0.566	0.742	0.865	1.311	1.493
2009	0.114	0.206	0.366	0.545	0.797	0.997	1.090	1.193
2010	0.098	0.277	0.363	0.463	0.672	1.067	1.309	1.484
2011	0.110	0.190	0.322	0.455	0.590	0.990	1.395	1.947
2012	0.081	0.216	0.318	0.455	0.743	0.871	1.239	1.553
2013	0.087	0.184	0.313	0.463	0.593	0.962	1.294	1.793
2014	0.092	0.226	0.333	0.505	0.712	1.006	1.483	1.565
2015	0.080	0.181	0.372	0.586	0.770	1.085	1.426	1.800

Table A35. Sequence of black sea bass ASAP model configurations considered for the Overall single area model.

Run	Fleets	Selectivity Blocks	Indices -Age	Indices-one age	Lambda 1	phases
A series of runs initially made to produce functional model.			Previous runs used NEFSC series split with Bigelow. Did not work, some combined with length calibration.			
Overall-1	1	2 (1989-1997, 1998-2015)	2	0	0.01	no F-Mult in first year
Overall-2	1	2 (1989-1997, 1998-2015)	2	0	0.01	no F-Mult in first year
Overall-3	1	2 (1989-1997, 1998-2015)	3	0	0.01	no F-Mult in first year
Overall-4	1	2 (1989-1997, 1998-2015)	3	7	0.01	no F-Mult in first year
Overall-4A	1	2 (1989-1997, 1998-2015)	3	7	0.01	no F-Mult in first year
Overall-4B	1	2 (1989-1997, 1998-2015)	3	7	0.01	no F-Mult in first year
Overall-4C	1	2 (1989-1997, 1998-2015)	3	7	0.01	no F-Mult in first year
Overall-4D	1	2 (1989-1997, 1998-2015)	3	7	0.05	no F-Mult in first year
Overall-5	1	2 (1989-1997, 1998-2015)	3	7	0.05	no F-Mult in first year
Overall-5(2)	1	2 (1989-1997, 1998-2015)	5	6	0.05	no F-Mult in first year
Overall-6	1	2 (1989-1997, 1998-2015)	6	5	0.05	no F-Mult in first year
Overall-7	1	2 (1989-1997, 1998-2015)	7	4	0.05	F-Mult in first year
Overall-8	1	2 (1989-1997, 1998-2015)	7	4	0.05	F-Mult in first year
Overall-5_2FLEET	2	4 (1989-1997, 1998-2015)	5	6	0.05	F-Mult in first year
Overall-6_2FLEET	2	4 (1989-1997, 1998-2015)	7	4	0.05	F-Mult in first year
Overall-10	2	3(1989-1997, 1998-2015, 1989-2015)	8	4	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
Overall-11	2	3(1989-1997, 1998-2015, 1989-2015)	8	5	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
Overall-12	2	3(1989-1997, 1998-2015, 1989-2015)	8	5	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
Overall-13_SPLIT	2	3(1989-1997, 1998-2015, 1989-2015)	9	5	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
Overall-18_SPLIT	2	3(1989-1997, 1998-2015, 1989-2015)	9	5	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
Overall-19_SPLIT	2	3(1989-1997, 1998-2015, 1989-2015)	9	5	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
Overall-20_SPLIT	2	3(1989-1997, 1998-2015, 1989-2015)	9	5	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year

Table A36. Changes within black sea bass ASAP overall single area model considered prior to selection of final model.

<u>Run</u>	<u>changes</u>
<i>A series of runs initially made to produce functional model.</i>	
Overall-1	Selectivity fixed at 1 >age 5
Overall-2	estimate catch selectivity for older ages
Overall-3	add NEAMAP and estimate index selectivity for older ages
Overall-4	
Overall-4A	decreased 2000-2002 NEFSC spring CV to 0.2
Overall-4B	Alter NY CV 0.4 to 0.6
Overall-4C	Return to empirical NEFSC spring CVs
Overall-4D	Return to fixed NEFSC spring CVs 0.6
Overall-5	added Rivard weight at age matrix
Overall-5(2)	added MADMF age indices and REC CPA age index      linked REC CPA to fleet selectivity
Overall-6	added NJ indices at age
Overall-7	added RI indices at age
Overall-8	made NEFSC flat topped selectivity
Overall-5_2FLEET	split fleets into trawl, non-trawl
Overall-6_2FLEET	added MADMF,RI age indices and REC CPA age index      linked REC CPA to fleet selectivity
Overall-10	added CT indices at age
Overall-11	aded VIMS age 1, fixed catch selectivity as flat top
Overall-12	
Overall-13_SPLIT	added Bigelow indices at age
Overall-18_SPLIT	all index phases 3
Overall-19_SPLIT	Fix age 5 and over flat topped in catch selectivity
Overall-20_SPLIT	Fix age 5 and over flat topped in catch selectivity

Table A37. Sequence of black sea bass ASAP model configurations considered for the North area model.

North						
Run	Fleets	Selectivity Blocks	Indices -Age	Indices-one age	Lambda 1	phases
ASAP-North1	1	1 (1989-2015)	3	4	0.1	no F-Mult in first year
ASAP-North2	1	1 (1989-2015)	4	3	0.1	no F-Mult in first year
ASAP-North3	1	1 (1989-2015)	4	3	0.1	no F-Mult in first year
ASAP-North4	1	1 (1989-2015)	4	3	0.1	no F-Mult in first year
ASAP-North5	1	1 (1989-2015)	4	3	0.1	no F-Mult in first year
ASAP-North6	1	1 (1989-2015)	4	3	0.05	no F-Mult in first year
NORTH-2Fleet	2	1 (1989-2015)	4	3	0.05	no F-Mult in first year
NORTH-3Fleet	2	1 (1989-2015)	3	4	0.05	no F-Mult in first year
NORTH-4Fleet	2	1 (1989-2015)	4	3	0.05	no F-Mult in first year
NORTH-4aFleet	2	1 (1989-2015)	4	3	0.05	no F-Mult in first year
NORTH-5Fleet	2	1 (1989-2015)	6	2	0.05	no F-Mult in first year
NORTH-ALB_Double_lc	2	2 (1989-2015)	5	3	0.05	no F-Mult in first year
NORTH-6Fleet	2	1 (1989-2015)	6	2	0.05	no F-Mult in first year
NORTH-7Fleet	2	1 (1989-2015)	6	2	0.05	F-Mult in first year
NORTH-8Fleet	2	2 (1989-2015)	6	2	0.05	F-Mult in first year
NORTH-9Fleet	2	2 (1989-2015)	6	2	0.05	F-Mult in first year
NORTH-10Fleet	2	2 (1989-2015)	6	2	0.05	F-Mult in first year
NORTH-11Fleet	2	2 (1989-2015)	7	1	0.05	F-Mult in first year
NORTH-12Fleet	2	2 (1989-2015)	7	1	0.05	F-Mult in first year
NORTH-13Fleet	2	2 (1989-2015)	7	1	0.05	F-Mult in first year
NORTH-14Fleet	2	2 (1989-2015)	7	1	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
NORTH-15Fleet	2	2 (1989-2015)	7	1	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
NORTH-16Fleet	2	2 (1989-2015)	7	1	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
NORTH-17Fleet	2	2 (1989-2015)	7	1	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
NORTH-18Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	7	1	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
NORTH-19Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	7	1	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
NORTH-20Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	6	1	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
NORTH-21Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	7	1	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
NORTH-23Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	7	1	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
NORTH-24Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	8	1	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
NORTH-24Fleet_SPLIT	2	3(1989-1997, 1998-2015, 1989-2015)	8	1	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
NORTH-26Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	8	1	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015 fleet 2	F-Mult in first year
NORTH-26	Final					

Table A38. Changes within black sea bass ASAP North area model considered prior to selection of final model.

North

<b>Run</b>	<b>changes</b>
ASAP-North1	NEFSC spring, NEAMAP,MADMF,RI,CT,NY,NEFSC winter
ASAP-North2	added MADMF aged indices
ASAP-North3	allow catch some dome in age 7,8
ASAP-North4	decrease index selectivity CV
ASAP-North5	index phase 5 to 3, increase NEFSC selectivity CV
ASAP-North6	lambda 1 reduced
NORTH-2Fleet	trawl and non-trawl split
NORTH-3Fleet	removed age comp from NEFSC spring
NORTH-4Fleet	reduced NEFSC indices to reduce catchability estimate
NORTH-4aFleet	reduced NEFSC indices to reduce catchability estimate
NORTH-5Fleet	add RI trawl indices and REC CPA
NORTH-ALB_Double_lc	tried NEFSC selectivity as double logistic
NORTH-6Fleet	reduced NEFSC q to match south
NORTH-7Fleet	new NY indices
NORTH-8Fleet	added 2nd selectivity block
NORTH-9Fleet	created flat top NEFSC indices
NORTH-10Fleet	linked REC CPA selectivity to fleet
NORTH-11Fleet	added CT ages
NORTH-12Fleet	removed REC CPA
NORTH-13Fleet	restored REC CPA
NORTH-14Fleet	change lamdda 1 CVs for fleet 2
NORTH-15Fleet	moved fixed age at catch selectivity up from 4 to 6
NORTH-16Fleet	changed catch selectivity lambdas to 0
NORTH-17Fleet	made catch selectivities equal between fleets to start
NORTH-18Fleet	added third selectivity block
NORTH-19Fleet	Fix age 5 and over flat topped in catch selectivity
NORTH-20Fleet	Fixed index selectivities as flat top, turned off ALB spring ages, unlinked REC CPUE to fleet selectivity
NORTH-21Fleet	restored ALB spring ages, made index selectivity lambda 0
NORTH-23Fleet	dropped NEFSC winter
NORTH-24Fleet	split NEFSC spring to include Bigelow; increase ESS of fleet
NORTH-24Fleet_SPLIT	index data corrections
NORTH-26Fleet	index phase adjustments to 3
NORTH-26	

Table A39. Changes within black sea bass ASAP South area model considered prior to selection of final model.

Run	Fleets	Selectivity Blocks	Indices -Age	Indices-one age	Lambda 1	phases
ASAP-South1	1	1 (1989-2015)	3	4	0.01	no F-Mult in first year
ASAP-South2Fleet	2	1 (1989-2015)	3	4	0.01	no F-Mult in first year
SOUTH-2Fleet	2	1 (1989-2015)	3	5	0.01	no F-Mult in first year
SOUTH-2AFleet	2	1 (1989-2015)	4	4	0.05	no F-Mult in first year
SOUTH-3Fleet	2	1 (1989-2015)	5	3	0.05	no F-Mult in first year
SOUTH-4Fleet	2	1 (1989-2015)	5	3	0.05	F-Mult in first year
SOUTH-5Fleet	2	2 (1989-2015)	4	4	0.05	F-Mult in first year
SOUTH-6Fleet	2	2 (1989-2015)	4	4	0.05	F-Mult in first year
SOUTH-7Fleet	2	2 (1989-2015)	5	3	0.05	F-Mult in first year
SOUTH-8Fleet	2	2 (1989-2015)	5	3	0.1	F-Mult in first year
SOUTH-10Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	5	3	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015	fleet 2 F-Mult in first year
SOUTH-11Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	5	3	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015	fleet 2 F-Mult in first year
SOUTH-12Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	5	3	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015	fleet 2 F-Mult in first year
SOUTH-13Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	5	3	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015	fleet 2 F-Mult in first year
SOUTH-14Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	5	3	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015	fleet 2 F-Mult in first year
SOUTH-15Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	5	3	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015	fleet 2 F-Mult in first year
SOUTH-16Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	5	3	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015	fleet 2 F-Mult in first year
SOUTH-17Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	5	3	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015	fleet 2 F-Mult in first year
SOUTH-18Fleet	2	3(1989-1997, 1998-2015, 1989-2015)	4	4	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015	fleet 2 F-Mult in first year
SOUTH-18Fleet_SPLIT	2	3(1989-1997, 1998-2015, 1989-2015)	5	3	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015	fleet 3 F-Mult in first year
SOUTH-20Fleet_SPLIT	2	3(1989-1997, 1998-2015, 1989-2015)	5	3	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015	fleet 3 F-Mult in first year
SOUTH-21Fleet_SPLIT	2	3(1989-1997, 1998-2015, 1989-2015)	5	3	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015	fleet 3 F-Mult in first year
SOUTH-22Fleet_SPLIT	2	3(1989-1997, 1998-2015, 1989-2015)	5	3	0.05 (fleet 1), 0.2-2008, 0.15 2009-2015	fleet 3 F-Mult in first year
SOUTH24.dat	final					



Table A40. Changes within black sea bass ASAP South area model considered prior to selection of final model.

<b>Run</b>	<b>changes</b>
ASAP-South1	altered index selectivity CVs
ASAP-South2Fleet	altered index selectivity CVs
SOUTH-2Fleet	includes REC CPA w/o ages
SOUTH-2AFleet	added REC CPA ages
SOUTH-3Fleet	added NJ indices at age
SOUTH-4Fleet	reduced NEFSC qs
SOUTH-5Fleet	removed NEFSC winter age comps
SOUTH-6Fleet	REC CPA selectivity linked to fleet
SOUTH-7Fleet	winter age comps back in
SOUTH-8Fleet	increase catch CV
SOUTH-10Fleet	add catch selectivity block, decrease index selectivity cv, lambda 1 changes
SOUTH-11Fleet	fix flat top catch selectivity
SOUTH-12Fleet	reduce at at full selectivity for NJ index
SOUTH-13Fleet	increase rec cpa ESS
SOUTH-14Fleet	reduce NEAMAP full selection to 4
SOUTH-15Fleet	remove index select lambdas
SOUTH-16Fleet	revised NJ indices
SOUTH-17Fleet	revise maturity data
SOUTH-18Fleet	limit NEAMAP to age 1 only
SOUTH-18Fleet_SPLIT	split NEFSC to Alb and Bigelow
SOUTH-20Fleet_SPLIT	reduce REC CPA CV, increase fleet ESS from 20 to 30
SOUTH-21Fleet_SPLIT	correct NEFSC indices
SOUTH-22Fleet_SPLIT	revise NEAMAP cv, phase change index 2
SOUTH24.dat	final

Table A41. Likelihood values from black sea bass area exchange model. Bold cell indicates minimum value.

Range	Area	Percent Movement										
		p_0	p_0.05	p_0.1	p_0.15	p_0.2	p_0.25	p_0.3	p_0.35	p_0.4	p_0.45	p_0.5
r_0	combined	16058.7	16047.9	16044.9	16043.4	16042.6	16042.1	16041.8	16041.6	16041.6	16041.7	16041.9
r_0.1	combined		16049.5	16045.6	16043.8	16042.7	16042.1	16041.8	16041.6	<b>16041.5</b>	16041.5	16041.6
r_0.2	combined			16046.9	16044.5	16043.2	16042.4	16042.0	16041.7	16041.6	16041.6	16041.6
r_0.3	combined				16045.6	16043.9	16043.0	16042.4	16042.0	16041.8	16041.8	16041.8
r_0.4	combined					16045.0	16043.8	16043.0	16042.5	16042.3	16042.2	16042.1
r_0.5	combined						16044.8	16043.8	16043.3	16042.9	16042.7	16042.7

Range	Area	Percent Movement										
		p_0	p_0.05	p_0.1	p_0.15	p_0.2	p_0.25	p_0.3	p_0.35	p_0.4	p_0.45	p_0.5
r_0	north	9734.8	9724.0	9721.0	9719.5	9718.7	9718.2	9717.9	<b>9717.8</b>	9717.8	9717.9	9718.1
r_0.1	north		9726.0	9722.1	9720.3	9719.3	9718.8	9718.5	9718.4	9718.4	9718.5	9718.6
r_0.2	north			9723.8	9721.4	9720.2	9719.5	9719.2	9719.0	9719.0	9719.0	9719.2
r_0.3	north				9722.9	9721.3	9720.5	9720.0	9719.7	9719.7	9719.7	9719.9
r_0.4	north					9722.7	9721.6	9720.9	9720.6	9720.5	9720.5	9720.6
r_0.5	north						9722.9	9722.0	9721.6	9721.3	9721.3	9721.3

Range	Area	Percent Movement										
		p_0	p_0.05	p_0.1	p_0.15	p_0.2	p_0.25	p_0.3	p_0.35	p_0.4	p_0.45	p_0.5
r_0	south	6324.0	6324.0	6323.9	6323.9	6323.9	6323.9	6323.8	6323.8	6323.8	6323.8	6323.7
r_0.1	south		6323.5	6323.5	6323.5	6323.4	6323.3	6323.3	6323.2	6323.2	6323.1	6323.0
r_0.2	south			6323.1	6323.0	6323.0	6322.9	6322.8	6322.7	6322.6	6322.5	6322.4
r_0.3	south				6322.7	6322.6	6322.5	6322.4	6322.3	6322.2	6322.1	6321.9
r_0.4	south					6322.3	6322.2	6322.1	6322.0	6321.8	6321.7	6321.6
r_0.5	south						6321.9	6321.8	6321.7	6321.6	6321.5	<b>6321.4</b>

Table A42. Results of the black sea bass ASAP models for 2015. The North and South model combine to form the Combined Area model, the Overall model is the single area model and the Area exchange model moves fish from the south back to the north during the offshore migratory period.

2015	North model	North adjusted	South model	South adjusted	Two Area models		Overall Single Area model	Area Exchange Model
					Combined	Combined-adjusted		
Catch (MT)	2767.5	2767.5	915.9	915.9	3683.4	3683.4	3683.4	3683.4
F	0.24	0.14	0.23	0.39	0.24	0.27	0.20	0.21
SSB (MT)	11,719	19,211	4,834	2,966	16,552	22,176	19,585	19,828
Exploitable Biomass (MT)	15,079	22,340	5,168	3,336	20,247	25,676	23,658	24,206
Total Biomass (MT)	17,306	27,733	6,838	4,276	24,143	32,010	24,155	29,339
Recruitment (000s)	3,342	6,484	21,574	11,519	24,917	18,002	27,863	24,066

Table A43. Proxy values of black sea bass biological reference points developed assuming  $F_{MSY} = F_{40\%}$ .

	SSB <sub>msy</sub>	B <sub>msy</sub>	F <sub>msy</sub>	MSY	1/2 SSB <sub>msy</sub>	1/2 B <sub>msy</sub>
Combined north and south	9,667 mt	17,256 mt	0.36	3,097 mt	4,834	8,628 mt
-2 SD	5,517	9,801		1,797	2,759	4,901
+2 SD	13,816	24,709		4,396	6,908	12,355
Overall model	10,296 mt	18,379 mt	0.36	3,298 mt	5,148	9,189 mt
-2 SD	5,615	10,026		1,831	2,808	7,101
+2 SD	14,977	26,733		4,765	7,489	11,278

Table A44. Short term black sea bass projections at  $F_{MSY}$  for the **Combined** area models. Values based on estimates not adjusted for retrospective bias.

Total Area Combined		<u>no</u> Rho adjust at $F_{MSY}$		
		- 2 STD DEV	Average	+ 2 STD DEV
SSB (mt)	2015		16,552	
	2016	9,434	15,055	20,676
	2017	8,380	13,299	18,217
	2018	7,483	11,618	15,753
	2019	5,699	10,782	15,864
Recruits (000s)	2015		24,917	
	2016	2,025	25,081	57,593
	2017	1,987	25,126	57,664
	2018	1,998	25,096	57,597
	2019	2,012	25,133	57,846
Jan 1 biomass (mt)	2015		24,143	
	2016	16,023	23,374	30,725
	2017	14,438	22,246	30,054
	2018	12,202	19,848	27,495
	2019	9,787	18,764	27,741
Catch (mt)	2015		3,683	
	2016		3,024	
	2017	2,759	4,446	6,132
	2018	2,523	3,822	5,121
	2019	2,012	3,493	4,974

Table A45. Short term black sea bass projections at  $F_{SQ}$  for the **Combined** area models. Values based on estimates not adjusted for retrospective bias.

Total		<u>no</u> Rho adjust		
Area Combined		$F_{status\ quo} = 0.24$		
		- 2 STD DEV	Average	+ 2 STD DEV
SSB (mt)	2015		16,552	
	2016	9,434	15,055	20,676
	2017	8,778	13,946	19,114
	2018	8,509	13,129	17,748
	2019	7,141	12,805	18,468
Recruits (000s)	2015		24,917	
	2016	2,025	25,081	57,593
	2017	1,987	25,126	57,664
	2018	1,998	25,096	57,597
	2019	2,012	25,133	57,846
Jan 1 biomass (mt)	2015		24,143	
	2016	16,023	23,374	30,725
	2017	14,438	22,246	30,054
	2018	13,181	21,154	29,127
	2019	11,425	20,848	30,271
Catch (mt)	2015		3,683	
	2016		3,024	
	2017	1,938	3,127	4,316
	2018	1,909	2,904	3,900
	2019	1,662	2,809	3,957

Table A46. Short term black sea bass projections at  $F_{MSY}$  for the **Combined** area models. Values based on estimates adjusted for retrospective bias.

Total Area Combined		<b>Rho adjusted</b> at $F_{MSY}$		
		- 2 STD DEV	Average	+ 2 STD DEV
SSB (mt)	2015		22,176	
	2016	11,950	18,670	25,391
	2017	10,270	15,918	21,565
	2018	8,914	13,407	17,900
	2019	6,706	11,849	16,991
Recruits (000s)	2015		18,002	
	2016	2,025	25,081	57,593
	2017	1,987	25,126	57,664
	2018	1,998	25,096	57,597
	2019	2,012	25,133	57,846
Jan 1 biomass (mt)	2015		32,010	
	2016	20,322	29,350	38,379
	2017	18,461	27,540	36,619
	2018	15,255	23,315	31,375
	2019	11,725	20,788	29,851
Catch (mt)	2015		3,683	
	2016		3,024	
	2017	3,484	5,467	7,451
	2018	3,037	4,494	5,950
	2019	2,398	3,901	5,403

Table A47. Short term black sea bass projections at F<sub>SQ</sub> for the **Combined** area models. Values based on estimates adjusted for retrospective bias.

Total Area Combined		<b>Rho adjusted</b> at Fstatus quo = 0.27		
		- 2 STD DEV	Average	+ 2 STD DEV
SSB (mt)	2015		22,176	
	2016	11,897	18,587	25,278
	2017	11,052	17,131	23,210
	2018	10,792	16,121	21,451
	2019	9,250	15,349	21,448
Recruits (000s)	2015		18,002	
	2016	2,025	25,082	57,599
	2017	1,987	25,127	57,671
	2018	1,998	25,098	57,604
	2019	2,012	25,135	57,853
Jan 1 biomass (mt)	2015		32,010	
	2016	20,253	29,246	38,238
	2017	18,461	27,540	36,619
	2018	17,176	25,874	34,572
	2019	14,865	24,704	34,543
Catch (mt)	2015		3,683	
	2016		3,024	
	2017	1,880	3,006	4,132
	2018	1,906	2,859	3,812
	2019	1,778	2,764	3,750

## FIGURES




Figure A1. Map of geographic boundaries for three black sea bass stocks in United States.







Figure A2. Black sea bass sex ratio at length and proportion female at age from NEFSC spring and winter surveys, 1984-2015.




Figure A3. Black sea bass von Bertalanffy growth curve, all regions and sexes combined.




Figure A4. Black sea bass maturity at length from NEFSC spring and winter bottom trawl survey data, 1989-2015.



Figure A5. Black sea bass age at maturity from NEFSC spring and winter bottom trawl survey data, 1989-2015.




Figure A6. Frequency of natural mortality estimates from a variety of methods (see also Table A2).




Figure A7 . Commercial landing time series for the North Atlantic stock of black sea bass.




Figure A8. Black sea bass commercial landings by gear type, 1989-2015.

### VIMS Black Sea Bass Age 1 index




Figure A9 Black sea bass age one indices of abundance from VIMS spring trawl survey in Chesapeake Bay, 1989-2013.

### MD Black Sea Bass Age 1 index




Figure A10. MD DNR trawl survey index of black sea bass age one indices of abundance in Maryland coastal bays, 1989-2015.

### Delaware Black Sea Bass Age 1 index




Figure A11. DE DFW trawl survey indices of black sea bass age one abundance in Delaware Bay, 1989-2015.

### New Jersey Black Sea Bass Trawl Survey Index




Figure A12. NJ DEM trawl survey indices ( $\pm$  95% CI) of black sea bass abundance, age 1-8+ collected in coastal New Jersey, 1989-2016.

### NY Black Sea Bass Age 1 index




Figure A13. NY DEC spring trawl survey indices of black sea bass age one abundance collected in Peconic Bay, Long Island, 1989-2016.

### CT Spring Black Sea Bass Trawl Survey Index




Figure A14. CT DEEP spring trawl survey indices ( $\pm$  95% CI) of black sea bass, ages 1-8+, collected in Long Island Sound, 1989-2015.

### RI Spring Black Sea Bass Trawl Survey Index




Figure A15. RI DEM spring trawl survey indices ( $\pm$  95% CI) of black sea bass, ages 1-8+, collected in Narragansett Bay, 1989-2015.

### MA Spring Black Sea Bass Trawl Survey Index




Figure A16. MA DMF spring trawl survey indices ( $\pm$  95% CI) of black sea bass, ages 1-8+, collected in south coastal Massachusetts, 1989-2015.



NEAMAP Spring North Black Sea Bass Trawl Survey Index




Figure A17. NEAMAP spring trawl survey indices ( $\pm$  95% CI bounded by 0) of black sea bass abundance, ages 1-8+ collected in coastal strata north of Hudson Canyon, 2008-2015.

NEAMAP Spring South Black Sea Bass Trawl Survey Index




Figure A18. NEAMAP spring trawl survey indices ( $\pm$  95% CI) of black sea bass abundance, ages 1-8+ collected in coastal strata south of Hudson Canyon, 2008-2015.




Figure A19. NEFSC winter trawl survey indices ( $\pm$  95% CI) of black sea bass abundance, ages 1-8+ collected in offshore strata north of Hudson Canyon 1992-2007.




Figure A20. NEFSC winter trawl survey indices ( $\pm$  95% CI) of black sea bass abundance, ages 1-8+ collected in offshore strata south of Hudson Canyon 1992-2007.




Figure A21. NEFSC spring trawl survey mean number per tow ( $\pm$  95% CI), 1969-2016, from offshore strata. Bigelow indices are not calibrated to the Albatross series.




Figure A22. NEFSC spring trawl survey indices ( $\pm$  95% CI) of black sea bass abundance, ages 1-8+ collected in offshore strata north of Hudson Canyon, 1989-2008 (*Albatross*) and 2009-2016 (*Bigelow*).

### NEFSC Spring South Black Sea Bass Trawl Survey Index




Figure A23. NEFSC spring trawl survey indices ( $\pm$  95% CI) of black sea bass abundance, ages 1-8+ collected in offshore strata south of Hudson Canyon, 1989-2008 (*Albatross*) and 2009-2016 (*Bigelow*).




Figure A24. Recreational catch per angler for black sea bass caught from NY north, 1989-2015.




Figure A25. Recreational catch per angler for black sea bass caught from NJ south, 1989-2015.




Figure A26. Recreational catch per angler for black sea bass caught from NC to ME, 1989-2015.




Figure A27. Commercial black sea bass Landings (mt), Maine-Cape Hatteras,NC, 1989-2015.




Figure A28. Percent of black sea bass commercial landings by market category, 1989-2015.




Figure A29. Mean length (cm  $\pm$  95% CI) of commercial landings black sea bass samples by market category, season and year.




Figure A30. Comparison of black sea bass commercial landings length (cm) samples, by half year, collected north and south of Hudson Canyon. Length distributions in each panel represent the sum of samples in years common between areas.






Figure A31. Annual sampling intensity for black sea bass sampled from the commercial fishery, 1989-2015. Intensity measured as the number of metric tons per length sampled. Larger values imply poorer sampling intensity.

Market Category	Qtr	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	
Unclassified	1,2										■		■																■
Small	1,2																												
Medium	1,2																												
Large	1,2																												
Jumbo	1,2		■				■			■																			
Unclassified	3,4				■	■	■	■	■			■	■	■	■	■							■	■	■				■
Small	3,4															■		■											
Medium	3,4		■		■																								
Large	3,4												■	■															
Jumbo	3,4		■		■	■	■	■	■				■	■	■														

Figure A32. Black sea bass commercial length samples by market category and season (qtr 1,2 is Jan-Jun, qtr 3,4 is Jul-Dec) and year. Black squares in the matrix represent cells with no samples available and substitution required.

Region/gear	Qtr	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	
North trawl	1,2		A	A	A	A	A								B														
North non-trawl	1,2	I	I	I	D	F	I	D	D	D	D	F	D	D	D	C	C	C	C			C							
North Trawl	3,4	C	A	A	A	A					E	E																	
North non-trawl	3,4	C	A	A	A	A	A	A	A	A	D	F	D	D	D		C						E	E	E				
South trawl	1,2				A	A	A	H				I																	
South non-trawl	1,2	I				F	I	I	I	D	D	I	D		E				E										
South trawl	3,4	C	A	A	A	A					G	C	G																
South non-trawl	3,4	I	I	I	I	F	I	I	I	I	I	I		E												C			

- A average all pre-1998
- B combined regions only
- C combined month only
- D combined region/month
- E other half year
- F adjacent year average
- G other region
- H other gear
- I combined annual/regions/gear

Figure A33. Sources of black sea bass commercial length sample by year, area and region. Cells without a letter indicate data available for that cell.




Figure A34. Black sea bass landings (number of fish in 000s) by area, 1989-2015.




Figure A35. Annual mean length (cm  $\pm$  2 SD) from expanded black sea bass commercial landings length frequencies for northern region, by season and gear category, 1989-2015.




Figure A36. Annual mean length (cm  $\pm$  2 SD) from expanded commercial black sea bass landings length frequencies for South region, by season and gear category, 1989-2015.




Figure A37. Annual mean black sea bass length (cm  $\pm$  2 SD) from expanded commercial landings length frequencies for region, by season and gear category, 1989-2015.





Figure A38. Annual mean black sea bass length (cm  $\pm$  2 SD) from expanded commercial discard length frequencies for region, by season 1989-2015.




Figure A39. Reason for discarding black sea bass in commercial fisheries, 2010-2014.




Figure A40. Relationship between black sea bass catch estimated by MRFSS survey vs MRIP survey, 2004-2011.






Figure A41. Black sea bass recreational landings (number) and weight (mt) coastwide, 1989-2015.




Figure A42. Black sea bass recreational landings (number) by year and region.




Figure A43. Mean length ( $\pm 2$  SD) of black sea bass recreational landings, Maine to Cape Hatteras, NC, 1989-2015.







Figure A44. Mean length (cm  $\pm$  2 SD) of black sea bass recreational landings by region and season, 1989-2015.




Figure A45. Black sea bass recreational discard mortalities (15% release mortality) by number and weight (mt) stock-wide, 1989-2015.




Figure A46. Black sea bass recreational discard mortalities (15% release mortality) by area, 1989-2015.




Figure A47. Mean length (cm) of black sea bass recreational discards by region.




Figure A48. Total black sea bass catch (mt) by region, 1989-2015.




Figure A49. Total black sea bass catch (mt) by fishery and disposition for the northern region, 1989-2015.




Figure A50. Total black sea bass catch (mt) by fishery and disposition for South region, 1989-2015.




Figure A51. Mean length ( $\pm 2$  SD) from black sea bass total expanded catch, by area, 1989-2015.





Figure A52. Mean weight at age (kg) for black sea bass by region, 1989-2015.





- ind.total
- ⊗ BIGELOW
- ▣ VIMS
- ⊠ MD
- ⊕ DE
- ⊖ NJ
- \* REC CPA
- ⊠ NEFSC winter
- ▽ NY
- ◇ CT
- × RI
- + MADMF
- △ NEAMAP
- NEFSC spring Alb

Figure A53. Final Root Mean Square Error (RMSE) values for black sea bass survey indices in final **Overall** single area model.




Figure A54. Fit diagnostics for the trawl catch in final black sea bass **Overall** single area model.

### Fleet 2 Catch (non-trawl)




Figure A55. Fit diagnostics for the non-trawl catch in final black sea bass **Overall** single area model.




Figure A56. Results of stock number, fishing mortality, total biomass, spawning biomass, exploitable biomass and recruitment from the black sea bass **Overall** single area model.




Figure A57. Retrospective pattern in average F estimates from black sea bass ASAP model for **Overall** single area model. Mohn's rho = -0.11




Figure A58. Retrospective pattern in spawning stock biomass (mt) estimates from black sea bass ASAP model for **Overall** single area model. Mohn's rho = 0.09




Figure A59. Retrospective pattern in total biomass (mt) estimates from black sea bass ASAP model for **Overall** single area model. Mohn's rho = 0.08




Figure A60. Retrospective pattern in total stock number estimates from black sea bass ASAP model for **Overall** single area model. Mohn's rho = 0.08




Figure A61. Retrospective pattern in exploitable biomass (mt) estimates from black sea bass ASAP model for **Overall** single area model. Mohn's rho = 0.11




Figure A62. Retrospective pattern in age 1 stock number estimates from black sea bass ASAP model for **Overall** single area model. Mohn's rho = 0.06




Figure A63. ASAP MCMC estimates of black sea bass F mult ( $\pm$  90% CI) from **Overall** single area model.




Figure A64. MCMC distribution of 2015 black sea bass Fmult estimates from **Overall** single area model.






Figure A65. ASAP MCMC estimates of black sea bass spawning stock biomass (MT  $\pm$  90% CI) from **Overall** single area model.




Figure A66. MCMC distribution of 2015 black sea bass spawning stock biomass (mt) estimates from **Overall** single area model.




Figure A67. ASAP MCMC estimates of black sea bass total biomass (mt  $\pm$  90% CI) from **Overall** single area model.




Figure A68. MCMC distribution of 2015 black sea bass total biomass (mt) estimates from **Overall** single area model.





Figure A69. Distribution of objective function components contribution to total likelihood for black sea bass **North** area model.



- ◆ ind.total
- \* bigelow
- ⊠ REC CPA
- ▽ NY
- ◇ CT
- × RI
- + MADMF
- △ NEAMAP
- NEFSC spring Alb

Figure A70. Final Root Mean Square Error (RMSE) values for survey indices in final black sea bass **North** area model.




Figure A71. Selectivity at age from trawl fleet in final black sea bass **North** area model.




Figure A72. Selectivity at age from non-trawl fleet in final black sea bass **North** area model.

### Fleet 1 Catch (trawl)




Figure A73. Fit diagnostics for the trawl catch in final black sea bass **North** area model.

### Fleet 2 Catch (non-trawl)




Figure A74. Fit diagnostics for the non-trawl catch in final black sea bass **North** area model.






Figure A75. Fits to 1989-2002 trawl catch proportions at age in final black sea bass **North** area model.




Figure A76. Fits to 2003-2015 trawl catch proportions at age and 1989 non-trawl catch proportion at age in final black sea bass **North** area model.




Figure A77. Fits to 1990-2004 non-trawl catch proportions at age in final black sea bass **North** area model.




Figure A78. Fits to 2005-2015 non-trawl catch proportions at age in final black sea bass **North** area model.

### Age Comp Residuals for Catch by Fleet 1 (trawl)




Figure A79. Trawl catch age composition residuals in final black sea bass **North** area model.

### Age Comp Residuals for Catch by Fleet 2 (non-trawl)




Figure A80. Non-trawl catch age composition residuals in final black sea bass **North** area model.




Figure A81. Fit diagnostics for the NEFSC spring trawl survey 1989-2008 (*Albatross*) in final black sea bass **North** area model.

### Age Comp Residuals for Index 1 (NEFSC spring Alb)




Figure A82. Age composition residuals for the NEFSC spring trawl survey 1989-2008 (*Albatross*) in final black sea bass **North** area model.



### Index 2 (NEAMAP)




Figure A83. Fit diagnostics for the NEAMAP trawl survey in final black sea bass **North** area model.

### Age Comp Residuals for Index 2 (NEAMAP)




Figure A84. Age composition residuals for the NEAMAP trawl survey in final black sea bass **North** area model.

### Index 3 (MADMF)




Figure A85. Fit diagnostics for the MADMF trawl survey in final black sea bass **North** area model.

### Age Comp Residuals for Index 3 (MADMF)




Figure A86. Age composition residuals for the MADMF trawl survey in final black sea bass North area model.

### Index 4 (RI)




Figure A87. Fit diagnostics for the RI trawl survey in final black sea bass **North** area model.

### Age Comp Residuals for Index 4 (RI)




Figure A88. Age composition residuals for the RI trawl survey in final black sea bass **North** area model.

### Index 5 (CT)




Figure A89. Fit diagnostics for the CT trawl survey in final black sea bass **North** area model.

### Age Comp Residuals for Index 5 (CT)




Figure A90. Age composition residuals for the CT trawl survey in final black sea bass **North** area model.



### Index 6 (NY)




Figure A91. Fit diagnostics for the NY trawl survey in final black sea bass **North** area model.

Index 7 (REC CPA)




Figure A92. Fit diagnostics for the Rec CPA in final black sea bass **North** area model.

### Age Comp Residuals for Index 7 (REC CPA)




Figure A93. Age composition residuals for the Rec CPA in final black sea bass **North** area model.

### Index 8 (bigelow)




Figure A94. Fit diagnostics the NEFSC spring trawl survey, 2009-2015 (*Bigelow*) in final black sea bass **North** area model.

### Age Comp Residuals for Index 8 (bigelow)




Figure A95. Age composition residuals for the NEFSC spring trawl survey, 2009-2015 (*Bigelow*) in final black sea bass **North** area model.




Figure A96. Index catchability estimates from final black sea bass **North** area model.




Figure A97. Comparison of age at maturity and average selectivity from final black sea bass **North** area model.




Figure A98. Results of stock number, fishing mortality, total biomass, spawning biomass, exploitable biomass and recruitment from the black sea bass **North** area model.






Figure A99. Retrospective pattern in average F estimates from black sea bass ASAP model for **North** area. Mohn's rho = 0.70.




Figure A100. Retrospective pattern in spawning stock biomass (mt) estimates from black sea bass ASAP model for **North** area. Mohn's rho = -0.39




Figure A101. Retrospective pattern in January 1 biomass (mt) estimates from black sea bass ASAP model for **North** area. Mohn's rho = -0.38




Figure A102. Retrospective pattern in exploitable biomass (mt) estimates from black sea bass ASAP model for **North** area. Mohn's rho = -0.33




Figure A103. Retrospective pattern in total stock number estimates from black sea bass ASAP model for **North** area. Mohn's rho = -0.42




Figure A104. Retrospective pattern in age 1 (recruits) estimates from black sea bass ASAP model for **North** area. Mohn's rho = -0.48




Figure A105. ASAP MCMC estimates of black sea bass  $F_{mult}$  ( $\pm 90\%$  CI) from **North** area model.




Figure A106. MCMC distribution of 2015 black sea bass  $F$  mult estimates from **North** area model.




Figure A107. ASAP MCMC estimates of black sea bass spawning stock biomass (mt  $\pm$  90% CI) from **North** area model.




Figure A108. MCMC distribution of 2015 black sea bass spawning stock biomass (mt) estimates from **North** area model.




Figure A109. ASAP MCMC estimates of black sea bass Jan 1 biomass (mt  $\pm$  90% CI) from **North** area model.




Figure A110. MCMC distribution of 2015 black sea bass total biomass (mt) estimates from **North** area model.




Figure A110a. Black sea bass median fishing mortality and SSB (000s mt)  $\pm$  90% CI and fishing mortality and total biomass (000s mt) from base estimates and retro-adjusted values for North and South area.





Figure A111. Distribution of objective function components contribution to total likelihood for black sea bass **South** area model.





- ◆ ind.total
- ◆ BIGELOW
- \* REC CPA
- ⊠ NEFSC winter
- ▽ VIMS
- ◇ MD
- × DE
- + NJ
- △ NEAMAP
- NEFSC spring Alb

Figure A112. Final Root Mean Square Error (RMSE) values for survey indices in final black sea bass **South** area model.




Figure A113. Selectivity at age from trawl fleet in final black sea bass **South** area model.




Figure A114. Selectivity at age from non-trawl fleet in final black sea bass **South** area model.

### Fleet 1 Catch (trawl)




Figure A115. Fit diagnostics for the trawl catch in final black sea bass **South** area model.

### Fleet 2 Catch (non-trawl)




Figure A116. Fit diagnostics for the non-trawl catch in final black sea bass **South** area model.




Figure A117. Fits to 1989-2002 trawl catch proportions at age in final black sea bass **South** area model.




Figure A118. Fits to 2003-2015 trawl catch and 1989 non-trawl catch proportions at age in final black sea bass **South** area model.




Figure A119. Fits to 1990-2004 trawl catch proportions at age in final black sea bass **South** area model.






Figure A120. Fits to 2005-2015 trawl catch proportions at age in final black sea bass **South** area model.

### Age Comp Residuals for Catch by Fleet 1 (trawl)




Figure A121. Age composition residuals of proportion at age from black sea bass trawl catch in final **South** area model.

### Age Comp Residuals for Catch by Fleet 2 (non-trawl)




Figure A122. Age composition residuals of proportion at age from non-trawl black sea bass catch in final **South** area model.

### Index 1 (NEFSC spring Alb)




Figure A123. Fit diagnostics for NEFSC spring 1989-2008 (*Albatross*) trawl survey in final black sea bass **South** area model.

### Age Comp Residuals for Index 1 (NEFSC spring Alb)




Figure A124. Age composition residuals for NEFSC spring 1989-2008 (*Albatross*) trawl survey in final black sea bass **South** area model.

### Index 2 (NEAMAP)




Figure A125. Fit diagnostics for NEAMAP trawl survey in final black sea bass **South** area model.

### Index 3 (NJ)




Figure A126. Age composition residuals for NEAMAP trawl survey in final black sea bass **South** area model.

### Age Comp Residuals for Index 3 (NJ)




Figure A127. Fit diagnostics for NJ trawl survey in final black sea bass **South** area model.



### Index 4 (DE)




Figure A128. Fit diagnostics for DE trawl survey in final black sea bass **South** area model.

### Index 5 (MD)




Figure A129. Fit diagnostics for MD trawl survey in final black sea bass **South** area model.

### Index 6 (VIMS)




Figure A130. Fit diagnostics for VIMS trawl survey in final black sea bass **South** area model.

### Index 7 (NEFSC winter)




Figure A131. Fit diagnostics for NEFSC winter trawl survey in final black sea bass **South** area model.

### Age Comp Residuals for Index 7 (NEFSC winter)




Figure A132. Age composition residuals for NEFSC winter trawl survey in final black sea bass **South** area model.

### Index 8 (REC CPA)




Figure A133. Fit diagnostics for Rec CPA in final black sea bass **South** area model.

### Age Comp Residuals for Index 8 (REC CPA)




Figure A134. Age composition residuals for Rec CPA in final black sea bass **South** area model.

### Index 9 (BIGELOW)




Figure A135. Fit diagnostics for NEFSC spring trawl survey (*Bigelow*) 2008-2015 in final black sea bass **South** area model.



### Age Comp Residuals for Index 9 (BIGELOW)




Figure A136. Age composition residuals for NEFSC spring trawl survey (*Bigelow*) 2008-2015 in final black sea bass **South** area model.




Figure A137. Index catchability estimates from final black sea bass **South** area model.




Figure A138. Comparison of maturity and average selectivity at age in final black sea bass **South** area model.








Figure A139. Results of stock number, fishing mortality, total biomass, spawning biomass, exploitable biomass and recruitment from the black sea bass **South** area model.




Figure A140. Retrospective pattern in average F estimates from black sea bass ASAP model for **South** area. Mohn's rho = -0.42




Figure A141. Retrospective pattern in spawning stock biomass (mt) estimates from black sea bass ASAP model for **South** area. Mohn's rho = 0.63




Figure A142. Retrospective pattern in total biomass (mt) estimates from black sea bass ASAP model for **South** area. Mohn's rho = 0.60




Figure A143. Retrospective pattern in exploitable biomass (mt) estimates from black sea bass ASAP model for **South** area. Mohn's rho = 0.55




Figure A144. Retrospective pattern in total stock number estimates from black sea bass ASAP model for **South** area. Mohn's rho = 0.70




Figure A145 Retrospective pattern in age 1 abundance (recruits) estimates from black sea bass ASAP model for **South** area. Mohn's rho = 0.87




Figure A146. ASAP MCMC estimates of black sea bass F mult ( $\pm$  90% CI) from **South** area model.




Figure A147. MCMC distribution of 2015 black sea bass F mult estimates from **South** area model.






Figure A148. ASAP MCMC estimates of black sea bass spawning stock biomass (mt  $\pm$  90% CI) from **South** area model.




Figure A149. MCMC distribution of 2015 black sea bass spawning stock biomass (mt) estimates from **South** area model.




Figure A150. ASAP MCMC estimates of black sea bass total biomass (mt  $\pm$  90% CI) from **South** area model.




Figure A151. MCMC distribution of 2015 black sea bass total biomass (mt) estimates from **South** area model.




Figure A152. Comparison of black sea bass ASAP model results for base (Combined) area model and Area Exchange model, by region. Base model shown in black, area exchange model in red.




Figure A153. Comparison of black sea bass ASAP model results for Combined area model, Overall single area model and area exchange model. Base (Combined) model shown in blue, Area Exchange model in red and single Overall model in black.




Figure A154. Exploitable biomass (mt) estimates from black sea bass ASAP **North** area model examining sensitivity to indices used in the model.




Figure A155. Total biomass (mt) estimates from black sea bass ASAP **North** area model examining sensitivity to indices used in the model.




Figure A156. Spawning stock biomass (mt) estimates from black sea bass ASAP **North** area model examining sensitivity to indices used in the model.




Figure A157. Fishing mortality estimates from black sea bass ASAP **North** area model examining sensitivity to indices used in the model.




Figure A158. Recruitment (age 1) estimates from black sea bass ASAP **North** area model examining sensitivity to indices used in the model.




Figure A159. Exploitable biomass (mt) estimates from black sea bass ASAP **South** area model examining sensitivity to indices used in the model.




Figure A160. Spawning stock biomass (mt) estimates from black sea bass ASAP **South** area model examining sensitivity to indices used in the model.




Figure A161. Total biomass (mt) estimates from black sea bass ASAP **South** area model examining sensitivity to indices used in the model.






Figure A162. Fishing mortality estimates from black sea bass ASAP **South** area model examining sensitivity to indices used in the model.




Figure A163. Recruitment (age 1) estimates from black sea bass ASAP **South** area model examining sensitivity to indices used in the model.




Figure A164. Likelihood profile of varying M estimates in regional black sea bass ASAP models.







Figure A 165. Sensitivity of CV for non-trawl fleet on the resulting black sea bass spawning biomass and fishing mortality estimates.




Figure A166. Time series trajectory of black sea bass SSB and F relative to biological reference points.




Figure A167. Time series trajectory of black sea bass total biomass and F relative to biological reference points.




Figure A168. Time series of black sea bass total catch (mt) with 3 year projections at F<sub>MSY</sub>. Projected catch based on the Overall single area model, the Combined area model and the Combined area model adjusted for retrospective patterns in each area.




Figure A169. Time series of black sea bass SSB (mt) with 3 year projections at F<sub>MSY</sub>. Projected SSB based on the Overall single area model, the Combined area model and the Combined area model adjusted for retrospective patterns in each area. SSB<sub>MSY proxy</sub> reference point shown.




Figure A170. Time series of black sea bass Total Biomass (mt) with 3 year projections at  $F_{MSY}$ . Projected Biomass based on the Overall single area model, the Combined area model and the Combined area model adjusted for retrospective patterns in each area.  $B_{MSYproxy}$  reference point shown.




Figure A171. Time series of black sea bass total catch (mt) with 3 year projections at  $F_{Sq}$ . Projected catch based on the Overall single area model, the Combined area model and the Combined area model adjusted for retrospective patterns in each area.




Figure A172. Time series of black sea bass SSB (mt) with 3 year projections at  $F_{SQ}$ . Projected SSB based on the Overall single area model, the Combined area model and the Combined area model adjusted for retrospective patterns in each area.  $SSB_{MSY proxy}$  reference point shown.




Figure A173. Time series of black sea bass Total Biomass (mt) with 3 year projections at  $F_{SQ}$ . Projected Biomass based on the Overall single area model, the Combined area model and the Combined area model adjusted for retrospective patterns in each area.  $B_{MSY proxy}$  reference point shown.

## **Appendix A**

The following information is the result of requests for additional information made by CIE reviewers during the SARC.






Figure A174. Retrospective pattern of black sea bass fishing mortality from Two Area model combined.




Figure A175. Retrospective pattern of black sea bass SSB (mt) from Two Area model combined.




Figure A176. Retrospective pattern of black sea bass age 1 recruitment from Two Area model combined.




Figure A177. Comparison of black seas bass SSB from the ASAP Two Area model (combined) and Stock Synthesis exploratory model.




Figure A178. Comparison of black seas bass fishing mortality SSB from the ASAP Two Area model (combined) and Stock Synthesis exploratory model.




Figure A179. Comparison of black seas bass recruitment SSB from the ASAP Two Area model (combined) and Stock Synthesis exploratory model.




Figure A180. Black sea bass likelihood profiles by component across varying natural mortality values using North ASAP model.




Figure A181. Black sea bass likelihood profiles by component across varying natural mortality values using South ASAP model.




Figure A182. Black sea bass likelihood profiles by component across varying natural mortality values using Two Area combined ASAP model.




Figure A183. Black sea bass survey standardized indices by area and age groups.

Table A48. Summary of black sea bass survey index rank, CVs, CV multipliers and residual mean square error for overall and regional models.

<b>NORTH</b>						<b>SOUTH</b>					
		Index						Index			
Rank	Index	avg. CV	model CV	CV multiplier	RMSE	Rank	Index	avg. CV	model CV	CV multiplier	RMSE
1	REC CPA	-	0.40		0.74	1	REC CPA	-	0.35		0.68
2	NEFSC spring Alb	0.65	0.70	1.08	1.49	2	NEFSC spring Alb	0.41	0.70	1.71	1.20
3	BIGELOW	0.38	0.65	1.71	1.06	3	BIGELOW	0.36	0.70	1.94	1.45
3	NEAMAP	0.48	0.30	0.63	1.16	3	NEFSC winter	0.26	0.70	2.69	1.11
4	MADMF	0.36	0.50	1.39	1.60	4	NJ	0.32	0.50	1.56	1.36
4	RI	0.53	0.80	1.51	1.16	5	NEAMAP (age 1)	0.22	0.40	1.82	0.77
4	CT	0.31	0.60	1.94	1.41	5	DE	-	0.60		1.38
5	NY	0.46	0.60	1.30	1.88	5	MD	-	0.60		1.16
						5	VIMS	0.18	0.60	3.33	1.13

<b>OVERALL</b>					
		Index			
Rank	Index	avg. CV	model CV	CV multiplier	RMSE
1	REC CPA	-	0.5		0.47
2	NEFSC spring Alb	0.38	0.6	1.58	1.23
3	BIGELOW	0.30	0.4	1.33	1.25
3	NEFSC winter	0.25	0.7	2.80	1.00
3	NEAMAP	-	0.4		0.72
4	MADMF	0.36	0.5	1.39	1.97
4	RI	0.53	0.8	1.51	0.96
4	CT	0.31	0.6	1.94	1.85
5	NY	0.46	0.8	1.74	1.65
4	NJ	0.32	0.5	1.56	1.16
5	DE	-	0.6		1.48
5	MD	-	0.4		1.67
5	VIMS	0.18	0.6	3.33	1.31

Ranked by:  
 Area coverage  
 Time series length  
 Ages represented

## Retrospective analysis for Black sea bass Stock Synthesis model ‘run\_164’

Additional analyses requested by SARC 63 review panel Gavin Fay<sup>1</sup>

<sup>1</sup>*Department of Fisheries Oceanography, University of Massachusetts Dartmouth,  
gfay@umassd.edu*

November 30, 2016

On 11/29/16, the Review panel requested to see retrospective analyses for the ‘base’ version of the SS black sea bass model.

Retrospective analyses were conducted for the ‘run\_164’ model detailed in WG documents Appendix X. Model peels using data only up to 2008 did not include the Bigelow survey indices or lengths. Note that for some peels some surveys will have a very small number of years of data (e.g. Bigelow, NEAMAP). An eight year peel was used to reflect the same extent of model/data years as the ASAP assessment models (the SS model used the 2016 spring survey data).

Table 1. Mohn’s  $\rho$  statistics for spawning biomass (SSB), fishing mortality (F), and recruitment (Recruits) based on an eight year peel.

<b>Quantity</b>	<b>Mohn’s <math>\rho</math></b>
SSB	-0.073
F	0.008
Recruits	-0.128






Figure A184. Retrospective analysis for black sea bass spawning biomass in SS exploratory model results




Figure A185. Retrospectives for time series of black sea bass fishing mortality in SS exploratory model results.




Figure A186. Retrospectives for time series of numbers of black sea bass Age 0 recruits (Recruitment in Table 1) in SS exploratory model results.




Figure A187. Retrospectives for time series of black sea bass relative spawning biomass (B/B0) in SS exploratory model results.




Figure A188. Retrospectives for time series of black sea bass F relative to F40% in SS exploratory model results.